

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/073646 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number:
PCT/US2004/004914
- (22) International Filing Date: 19 February 2004 (19.02.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/448,663 19 February 2003 (19.02.2003) US
- (71) Applicant (for all designated States except US): **UNIVERSITY OF ROCHESTER** [US/US]; 601 Elmwood Avenue, Box 706, Rochester, NY 14642 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **KYRKANIDES, Stephanos** [US/US]; 1963 Harris Road, Penfield, NY 14526 (US). **TALLENTS, Ross, H.** [US/US]; 1333 Lake Road, Rochester, NY 14620 (US).
- (74) Agents: **HUIZENGA, David, E.** et al.; Needle & Rosenberg, P.C., Suite 1000, 999 Peachtree Street, Atlanta, GA 30309-3915 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TREATMENT OF PAIN THROUGH EXPRESSION OF OPIOID RECEPTORS

(57) Abstract: Disclosed are compositions and methods related to nucleic acid constructs containing a HUMOR encoding element. These constructs can be used in the treatment of pain.

WO 2004/073646 A2

TREATMENT OF PAIN THROUGH EXPRESSION OF OPIOID RECEPTORS

[01] This application claims priority of United States Provisional Application No. 60/448,663, filed on February 19, 2003, which is herein incorporated by reference in its entirety.

I. BACKGROUND

5 [02] Tissue injury and nerve damage caused by trauma, infection, arthritis or iatrogenic procedures can produce inflammation, spontaneous pain and hyperalgesia (Levine JD and Taiwo YO, *Anesth Prog* 1990;37:133-35; Goelet P, et al., *Nature*. 1986;322(6078):419-22). Furthermore, patients affected by temporomandibular joint and/or masticatory muscle (orofacial) pain often suffer because dental, surgical and/or pharmacologic therapies do not consistently give adequate symptom
10 relief. In fact, according to Public Health Services (PHS) estimates, there are more than 50 million Americans who experience chronic pain with 45% seeking medical care at some point in their lives. It is also estimated that 40% of pain patients never receive adequate relief. Lipton et al. (Lipton JA, et al., *JADA* 1993;124:115-21.) reported that 22% of the population in the United States experienced at least one episode of orofacial pain in the last six months. It is estimated that approximately \$80
15 billion is spent annually to treat pain and that 40% of that is to treat craniofacial pain (Bonica JJ. Preface et al., eds. *Advances in Pain Research Therapy*, Vol 3. New York: Raven Press; 1973:v-vii). To date, pharmacological approaches still dominate the clinical pain arena, with only modest efforts being directed towards the development of new innovative treatment regimes for the management of pain. Disclosed are vectors and methods for reducing pains, such as myalgic and
20 arthralgic pains, and such as those in the orofacial region.

II. SUMMARY

[03] As embodied and broadly described herein, disclosed herein, in one aspect, are vector constructs that comprise sequence encoding a polypeptide for treating pain. Also disclosed are methods for treating pain by expressing the μ -opioid receptor protein in nerve cells.

25 [04] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive.

III. BRIEF DESCRIPTION OF THE DRAWINGS

[05] Figure 1 shows the human μ -opioid transient transfection in N2a cells. Figure 1A shows that immunocytochemistry reveals expression of HUMOR driven by the cytomegalovirus
30 CMV promoter in neuronal cells using monoclonal antibody. Figure 1B shows mock transfected cells.

[05] Figure 2 shows FIV(lacZ) local administration to the temporomandibular joint (TMJ) area of the face. Figure 2A shows FIV was injected at the right TMJ with 10^8 infectious particles per mL, which receives sensory innervation from the mandibular division of the trigeminal nerve. Sagittal(B) and horizontal (C) sections of the right gasserion (trigeminal) ganglion display X-gal positive neuronal cellbodies that were transduced following FIV(lacZ) injection in the TMJ.

[06] Figure 3 shows a representation of a lentiviral system containing the HUMOR gene. The 3-vector FIV(HUMOR) system. The FIV(HUMOR) lentiviral system is comprised of 3 vectors: Packaging vector providing the packaging instructions in trans, - VSV-G envelop vector (VSV-G sequence in SEQ ID NO:54) providing the envelop instructions in trans, - and FIV(HUMOR) vector containing the therapeutic gene.

[07] Figure 4 shows a schematic of an exemplary LIV vector carrying a HUMOR cassette.

[08] Figure 5 shows FIV(lacZ) injection (a total of 5×10^6 infectious particles) to the right TMJ resulted in widespread infection of hard as well as soft tissues of the joint. (A) Sagittal TMJ sections analyzed by β -galactosidase immunohistochemistry and counter-stained by nuclear fast red revealed expression of the reporter gene lacZ in the hypertrophic zone of the condyle, primarily comprised of cartilaginous cells, (B) as well as in the meniscus, endothelial cells and perivascular osteocytes. Panel (C) depicts TMJ sections from a saline injected animal.

c=condyle; d=disk; m=muscle; v=vessel.

[09] Figure 6 shows the development of the control FIV(Δ' lac) vector with inactive β -galactosidase gene. (A) The reporter gene lacZ was inactivated after deletion of a critical *placZ* DNA fragment containing the β -galactosidase gene transcription initiation site by restriction enzyme-mediated excision and re-ligation of the backbone vector. (B) The structure of mutated FIV(Δ' lac) and wild type FIV(lacZ) viral vectors were confirmed by PCR following transient transfection into the murine cell line NIH 3T3. The presence of viral DNA in cells was detected by a 444 bp DNA band utilizing the "FIV" primers (as depicted in panel A). The complete structure of lacZ gene was confirmed by a 1.7 kb DNA band utilizing the lacZ primers (depicted as UP, LP in panel A). In the case of the mutated FIV(Δ' lac), there was lack of the 1.7 kb DNA band as the annealing site for the lower primer LP was excised. (C) Deletion of the lacZ transcription initiation sequence in the FIV(Δ' lac) resulted in inactivation of the β -galactosidase

reporter gene as demonstrated by the lack of X-gal staining compared to (D) the FIV(lacZ) vector.

[10] Figure 7 shows FIV(lacZ) and FIV(Δ' lac) injections (5×10^6 infectious particles) in the right TMJ of mice resulted in successful infection of primary sensory neurons located in the ipsilateral trigeminal ganglion. The animals' left side TMJ was not treated (A) The presence of backbone FIV DNA in the right trigeminal ganglia ipsilateral to FIV injections was detected by a 444 bp DNA band in lanes 1 and 3, utilizing the "FIV" primers (as depicted in panel A), suggesting successful transduction of the trigeminal sensory neurons by FIV vectors. Lanes 2 and 4 do not display any viral DNA as they represent left side ganglia. (B) The inactive form of β -galactosidase gene in transduced neurons was detected by the absence of the 1.7 kb DNA band (lane 1) compared with the wild type gene (lane 3). Lanes 2 and 4 do not display any viral DNA as they represent left side ganglia. (C) The successful extraction of genomic DNA from left and right ganglia was confirmed by PCR utilizing primers designed for the murine housekeeping gene G3PDH (385 bp).

[11] Figure 8 shows injection of FIV(lacZ) in the right TMJ (5×10^6 infectious particles) resulted in successful transduction of primary sensory neurons with the reporter gene β -galactosidase in trigeminal ganglia ipsilateral to the treated joint. (A) β -galactosidase expression was detected by X-gal histochemistry in sagittal sections of right-side ganglion (4X), (B) primarily at its posterior and posterolateral region (20X). (C) Injection of FIV(Δ' lac) did not result in β -galactosidase expression. (D) The X-gal staining was confirmed with immunocytochemistry employing antibodies raised against bacterial β -galactosidase following FIV(lacZ) injection compared to (E) FIV(Δ' lac) treatment.

[12] Figure 9 shows The neuronal cell line N2 α was infected with HIV(HUMOR) [a Lenti virus]. Total RNA was extracted from infected and control cells. The levels of HUMOR and G3PDH transcript were assessed by RT-PCR. Minimal amounts of HUMOR were detected in naïve cells (C1, C2), as well as cells infected with the HIV(lacZ) virus (L1, L2). In contrast, HUMOR was readily detected in cells infected with HIV(HUMOR). As a control, the housekeeping gene G3PDH transcript was detected in all samples analyzed.

IV. DETAILED DESCRIPTION

[13] Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that the disclosure is not limited to specific synthetic methods or specific recombinant biotechnology methods unless otherwise specified, or to particular reagents unless otherwise specified, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

[14] Disclosed are the components to be used to prepare the disclosed compositions as well as the compositions themselves to be used within the methods disclosed herein. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutation of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a particular μ -opioid receptor vector is disclosed and discussed and a number of modifications that can be made to a number of molecules including the μ -opioid receptor vector are discussed, specifically contemplated is each and every combination and permutation of the μ -opioid receptor vector and the modifications that are possible unless specifically indicated to the contrary. Thus, if a class of molecules A, B, and C are disclosed as well as a class of molecules D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited each is individually and collectively contemplated meaning combinations, A-E, A-F, B-D, B-E, B-F, C-D, C-E, and C-F are considered disclosed. Likewise, any subset or combination of these is also disclosed. Thus, for example, the subgroup of A-E, B-F, and C-E would be considered disclosed. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the disclosed methods.

A. Definitions

[15] As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

[16] Ranges can be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also herein disclosed as "about" that particular value in addition to the value itself. For example, if the value "10" is disclosed, then "about 10" is also disclosed. It is also understood that when a value is disclosed that "less than or equal to" the value, "greater than or equal to the value" and possible ranges between values are also disclosed, as appropriately understood by the skilled artisan. For example, if the value "10" is disclosed the "less than or equal to 10" as well as "greater than or equal to 10" as well as "less than" and "greater than" 10 are also disclosed. It is also understood that the throughout the application, data is provided in a number of different formats, and that this data, represents endpoints and starting points, and ranges for any combination of the data points. For example, if a particular data point "10" and a particular data point 15 are disclosed, it is understood that greater than, greater than or equal to, less than, less than or equal to, and equal to 10 and 15 are considered disclosed as well as between 10 and 15.

[17] In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

[18] "Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

[19] "Primers" are a subset of probes which are capable of supporting some type of enzymatic manipulation and which can hybridize with a target nucleic acid such that the enzymatic manipulation can occur. A primer can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art which do not interfere with the enzymatic manipulation.

[20] "Probes" are molecules capable of interacting with a target nucleic acid, typically in a sequence specific manner, for example through hybridization. The hybridization of nucleic

acids is well understood in the art and discussed herein. Typically a probe can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art.

[21] Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains. The references disclosed are also individually and specifically incorporated by reference herein at least for the material contained in them that is discussed in the sentence in which the reference is relied upon.

B. Compositions and methods

[22] The μ -opioid receptor (HUMOR) is a key component of the intrinsic anti-nociceptive pathway in mammals: descending bulbospinal serotonergic and noradrenergic neuronal projections exert anti-nociceptive effects via release of endogenous opioids, which in turn activate μ -opioid receptors present on the presynaptic membrane of the primary sensory neurons. Pain stimulus travels through the nerve to the brain through activation of nociceptors. The activation of μ -opioid receptors through binding of opioids interrupts the transmission of the pain signal. Mammals release endogenous opioids when under pain assault and billions are spent each year in the pharmaceutical industry to treat pain through the administration of opioids and opioid like molecules that target the μ -opioid receptors. Disclosed herein are compositions and methods for the treatment of pain, which do not require the administration of μ -opioid receptor targeted molecules or utilize lower effective amounts of opioid receptor targeted molecules. The disclosed methods involve the over-expression of μ -opioid receptors, which can make the nerve cell more receptive to endogenous opioid molecules or to opioids or opioid like molecules administered as a pharmaceutical. Over expression of the μ -opioid receptors can occur through simulation of endogenous opioid receptor genes or through transgenic therapy that delivers a construct encoding the opioid receptor.

[23] Receptor up-regulation is designed to result in circumventing the observed desensitization following prolonged opioid drug administration, which in part occurs as a decrease in receptor expression. Furthermore, the strategy can take advantage of the existing intrinsic anti-nociceptive mechanism by ensuring adequate μ -opioid receptor presence at the site of the central processing of pain. This adequate receptor presence is consistent with heightened sensitivity of patients to drugs administered exogenously, which is consistent with requiring

smaller doses of opioid analgesics, such as Ultram, Fentanyl, and Darvon, which otherwise commonly result to pathologic addiction.

[24] Disclosed are compositions and methods to target the expression of opioid receptors, such as the μ -opioid receptors, such as human opioid receptors, to sensory neurons innervating regions that can experience pain, such as orofacial regions that experience nociception. Disclosed are compositions and methods for targeting opioid receptors, such as the μ -opioid receptors, expression in sensory orofacial neurons. Also disclosed are compositions and methods for targeting opioid receptors, such as the μ -opioid receptors, expression in any sensory neuron. For example, the compositions and methods can be used in any sensory neuron, wherein the sensory neuron processes pain or other "input" signals from peripheral tissues (e.g., joints, amputated limbs, extracted or endodontically treated teeth), as well as vital organs. For example, disclosed are vectors, such as feline immunodeficiency lentiviral vectors (FIV), rAAV vectors, HSV Amplicon, and liposomes for delivery of the opioid receptor DNA. Administration of the vectors peripherally to infect those sensory neurons, such as those innervating the orofacial region, can be performed. For example, the vectors can be delivered at the point of pain, for example, an extremity, by for example, injection into the extremity. Disclosed are vectors, such as FIV, rAAV, HSV Amplicon, and liposomes, capable of stably transducing terminally differentiated cells, including neurons.

a) Nervous system

[25] The nervous system can be divided into two parts: central and peripheral. The central nervous system consists of the encephalon or brain and the medulla spinalis or spinal cord. These two parts, the brain and the spinal cord are continuous with one another at the level of the upper border of the atlas vertebra. The peripheral nervous system consists of a series of nerves, which connect the central nervous system to all of the tissues in the body. Nerves also are often grouped as cerebrospinal and sympathetic. However, since the two groups are intimately connected and closely intermingled these distinctions are not absolute. Nerve cells can also be classified as efferent or afferent nerves. Efferent nerve cells are nerve cells that transmit signals from the brain to the periphery and afferent nerve cells are nerve cells that transmit signals from the periphery to the brain.

[26] Neurons act as pain pathways and these pathways include peripheral, spinal, and supraspinal elements. The peripheral part of the system includes the primary afferent sensory neurons. These neurons are called nociceptors, and can be found throughout the body, such as

in the skin, muscle, connective tissue, the cardiac system, and abdominal and thoracic viscera. Nociceptors are unencapsulated nerve endings that detect thermal, mechanical, or chemical stimuli, and are thus, not small molecule receptors. Nociceptors can be thinly myelinated or unmyelinated nerve fibers. The thinly myelinated variety are termed A-delta fibers and the
5 unmyelinated variety are termed C-polymodal fibers. The primary functional difference between A and C delta fibers is that A-delta fibers are rapidly conducting and C delta fibers are slowly conducting. This means that A delta fibers transmit sensations perceived as fast, sharp, well-localized pricking pain, and C-polymodal fibers transmit feeling via thermal, mechanical, and chemical stimuli transmitting sensations perceived as dull, aching, burning, poorly localized
10 pain.

[27] Most A-delta and the C-polymodal afferent fibers enter the dorsal horn of the spinal cord by way of the dorsal nerve roots and their ganglia. Wide dynamic range neurons receive nociceptive and non-nociceptive input from the skin, muscle, and viscera. This convergence can account for visceral referred pain. Impulses are then transmitted to the brain
15 by the spinal thalamic tract (STT). Near the thalamus, the STT bifurcates into the neospinothalamic tract and the paleospinothalamic tract, projecting to the thalamus, hypothalamus, periaqueductal gray matter (PAG) in the brain stem. The thalamus processes sensory input is projected to the cerebral cortex, basal ganglia, and limbic system. Descending pathways conduct transmission from the brain to the spinal cord control and modify afferent
20 sensory input.

[28] Nociception can be thought of as the detection of tissue damage by nociceptors. Modulation of nociception occurs peripherally, spinally, and supraspinally. Tissue damage is associated with the release of chemical mediators, such as serotonin, histamine, bradykinin, cytokines, prostaglandins, and leukotrienes, which produce inflammation, and occurs in the
25 peripheral system. The pain transmission is modulated by these events and this lowers excitability threshold of the nociceptor threshold so that stimuli normally non-painful stimuli become painful. This is called nociceptor sensitization. Two other substances that sensitize nociceptors are substance P and glutamate, which can be released from nerve terminals.

[29] The signals from the nociceptors are processed in the dorsal horn of the spine. Repetitive, convergent input from A-delta and C polymodal fibers at the dorsal horn can result
30 in a state where less stimulation is required for the generation of a pain response. This is known

as the wind-up phenomenon, and is thought to be initiated by the release of substance P and the excitatory amino acids glutamate and aspartate.

[30] The brain also signals the spinal cord to modulate the pain response. The PAG region of the brainstem contains high concentrations of opioid receptors, and sends projections to the rostral medulla and eventually to the dorsal root inhibiting ascending pain impulses. Thus, the activation of the opioid receptors interrupts the transmission of the pain signal. Descending pathways can also stimulate spinal nociceptive transmission as well.

b) Pain

[31] Pain is typically classified into two categories: nociceptive pain (somatic pain) and neuropathic pain. Nociceptive pain is pain that is sensed after some type of trauma. The nociceptive pain is sensed by the "nociceptor" sensory fibers which are connected to the nervous system. After an injury to a muscle, soft tissue (ligaments, tendons), bones, joints, or skin (or other organs), these sensory fibers are stimulated which causes a transmission of a signal through an afferent neuron to the brain. Nociceptive pain is often characterized as a deep aching, throbbing, gnawing, or sore sensation. Common examples of nociceptive pain include: pain after trauma (e.g. a car accident or a fall), postoperative pain, and arthritis pain. Nociceptive pain is usually localized and gets better with healing.

[32] Neuropathic pain is pain caused by damage to nerve tissue. Neuropathic pain is often characterized as burning, severe shooting pains, and/or persistent numbness or tingling. Common examples of neuropathic pain related to back pain include sciatica, pain that travels from the spine down the arm, and pain that persists after back surgery.

[33] It is thought that in some cases prolonged nociceptive pain may progress to neuropathic pain, and a patient may have both nociceptive and neuropathic pain at the same time. Pain is also often classified as acute pain or chronic pain. Acute pain is characterized as pain where the amount of pain directly correlates with the level and duration of tissue damage. Acute pain therefore, provides a protective reflex, such as the reflex to move your hand immediately if you touch a sharp object. This type of pain is a symptom of injured or diseased tissue, so that when the underlying problem is cured the pain goes away. Acute pain is a form of nociceptive pain. Chronic pain on the other hand, does not correlate with the severity of the insult, and therefore, typically will not serve a protective function. Prolonged damage to tissues, i.e. knee pain or tooth ache, will eventually result in plastic (non reversible) changes in the neurons that process pain from that area, which now facilitate either allodynia and/or

hyperalgesia. Chronic pain is born following these plastic neuronal changes, whereby the neurons are now "sick" and pain will occur even in the absence of peripheral stimulus (e.g., amputated limbs, extracted teeth). In fact, its basis is neuropathic now, and neurons continuously send pain messages to the brain even though there is no continuing tissue damage.

5 Neuropathic pain is a form of chronic pain.

(1) Anatomy of orofacial pain

[34] The mandibular division of the trigeminal nerve provides sensory innervation to the TMJ and masticatory muscles. The cell bodies of these primary sensory neurons are located in the inferior portion of the trigeminal ganglion extending their unmyelinated (C-fibers) or
 10 thinly myelinated (A δ -fibers) peripheral projections to structures of the face and jaws. More specifically, nociceptive innervation to the temporomandibular joint (TMJ) is primarily provided by the auriculotemporal nerve of the mandibular division of the trigeminal nerve (Sessle BJ, Hu JW (1991). *Can J Physiol Pharmacol* 69: 617-626). A δ and C nerve fibers, whose cell bodies are located in the posterolateral part of the trigeminal ganglion (Yoshino K, et
 15 al. (1998). *Arc Oral Biol*; 43: 679-686), project distally and terminate as non-encapsulated free nerve endings dispersed throughout the posterolateral part of the TMJ capsule (Bernick S (1962). *Oral Surg* 15:488-492; Thilander B (1964). *Acta Odont Scan* 22:151-156; Frommer J, Monroe CW (1966). *J Dent Res* 45:1762-1766; Klineberg I (1971). *Ann Royal Coll Surg Engl* 49:268-288), the posterior band of the meniscus and the posterior attachment (Dressen D, et al.
 20 (1990). *Acta Anat* 139:154-160; Kido MA, et al. (1991). *Arch Oral Biol* 36:397-400, Kido MA, et al. (1993). *J Dent Res* 72:592-598; Wink CS, et al. (1992). *J Oral Maxillofac Surg* 50:334-337). Inflammation, injury or other agents may cause excitation of the free and unspecialized nerve endings of the unmyelinated C-fibers, which are predominately involved in the transmission of nociception from the TMJ, muscles of mastication as well as the pulp of
 25 teeth. The central projections enter the brain stem via the ventrolateral pons, descend caudally as the dorsolateral trigeminal tract and synapse with second order sensory neurons at the substantia gelatinosa of the subnucleus caudalis of the descending trigeminal nucleus (medullary dorsal horn). Second order sensory neurons extend projections to the nucleus proprius, followed by subsequent projections to the intermedial gray, and then to the reticular formation
 30 of the brain stem, and through the intralaminar nuclei of the thalamus project wide spread connections into the cortex. The ascending sensory neural architecture is also susceptible to an intrinsic opioid-releasing anti-nociceptive descending system, the inhibitory effects of which

are mediated by opioid receptors expressed in the presynaptic membrane of the primary sensory neurons. Although pain is initially elicited at a peripheral site, it is further centrally modulated, i.e. in the brain, enhanced or attenuated, therefore making this aforementioned central processing of pain a major component in sensory orofacial nociception.

5 [35] In the quest for developing new therapies for orofacial pain, gene therapy appears to be an emerging treatment method (Kuboki T, et al. (1999). *Arc Oral Biol* 44: 701-709; Pohl M, Braz J (2001). *Eur J Pharmacol* 429: 39-48; Baum BJ, et al. (2002). *JADA* 133: 35-44). For example, it has been previously suggested that delivery of antisense oligonucleotides developed against nociceptive genes to appropriate tissues may offer alternatives in designing novel
10 treatments for pain management (Wu CL, et al. (2001). *Anesthesiology* 95: 216-240).

 [36] Disclosed herein, transfer of anti-nociceptive genes to sensory trigeminal neurons innervating the orofacial region can be achieved after injection of lentiviral vectors at the painful site, such as the TMJ, resulting in their uptake by free nerve endings and retrograde transport to the sensory cells' nuclei. Previous studies demonstrated axonal retrograde transport
15 of horseradish peroxidase from the TMJ to the central nervous system (Romfh JH, et al. (1979). *Exp Neurol* 65: 99-106; Capra NF (1987). *Somatosensory Res* 4: 201-213) including the trigeminal ganglia (Yoshino K, et al. (1998). *Arc Oral Biol*; 43: 679-686). In evaluating the employment of lentiviral vectors as the basis for TMJ gene therapy, as an example, VSV-G pseudotyped feline immunodeficiency viral vectors (FIV) were used. These vectors are capable
20 of stably transducing dividing, growth arrested as well as post-mitotic cells, as it is capable of transgene integration into the host's genome (Poeschla EM, et al. (1998). *Nature Med* 4: 354-357). VSV-G pseudotyping of viral vectors confers a broad range of host specificity, including human and murine cells, as infection is mediated by the interaction of the viral envelope protein and a phospholipid component of the cell membrane leading to membrane-fusion mediated
25 entry (Burns JC, et al. (1993). *Proc Natl Acad Sci USA* 90: 8033-8037; Carneiro FA, et al. (2002). *J Virol* 76: 3756-3764). Therefore, FIV vectors can mediate sustained gene expression in non-dividing terminally differentiated trigeminal sensory neurons, a property unique to lentiviral vectors.

c) Current pharmacologic agents in the management of pain

30 [37] Non-steroidal anti-inflammatory drugs (NSAID's) are often utilized as the first line of agents for the management of pain. NSAID's primarily exert their pain-killing effects by inhibiting the production of prostanoids and attenuating peripheral inflammatory conditions that

may be responsible for pain elicitation. Alternatively, corticosteroids may be utilized with peripheral routes of action. In contrast, exogenously administered opioid drugs (morphine) mimic the effects of the endogenous opioids by crossing the blood brain barrier (BBB). Similarly, tricyclic antidepressants that cross the BBB have been also employed in cases of chronic pain by inhibiting the reuptake of serotonin and norepinephrine. However, each of these four classes of drugs is characterized by significant side effects that prohibit their long term use as well as often show unfavorable treatment outcomes.

d) Opioid receptors and mechanism of action

[38] Opioid analgesics have been used for pain management for hundreds of years. Opium itself consists of the dried latex from the unripe fruit of the opium poppy *Papaver somniferum*. Morphine is isolated from opium. Opioid receptors exist in the spinal and supraspinal regions of the nervous systems. Opioids can modulate neuronal transmission by binding to opioid receptors in the dorsal-root ganglia, the central terminals of primary afferent neurons (LaMotte C, et al., Brain Res 1976;112:407-12; Fields HL, et al., Nature 1980;284:351-3) and peripheral sensory-nerve fibers and their terminals (Stein C, et al., Proc Natl Acad Sci U S A 1990;87:5935-9; Hassan AHS, et al., Neuroscience 1993;55:185-95.. The dorsal-root ganglia and trigeminal ganglion (Xie GX, et al., Life Sciences 1999; 64:2029-37; Li JL, et al., Brain Res 1998; 794:347-52.) contain messenger RNA (mRNA) for opioid receptors (Schafer M, et al., Eur J Pharmacol 1995;279:165-9; Mansour A, et al., Brain Res 1994;643:245-65) and primary afferent nerves mediate the peripheral antinociceptive effects of morphine (Bartho L, et al., Naunyn Schmiedebergs Arch Pharmacol 1990;342:666-70). The presence of endogenous opioids and their receptors are able to produce a potent antinociception. Opioids increase potassium currents and decrease calcium currents in the cell bodies of sensory neurons (Werz MA, Macdonald RL., Neurosci Lett 1983;42:173-8; Schroeder JE, et al., Neuron 1991;6:13-20), both of which can lead to the inhibition of neuronal firing and transmitter release. A similar process occurring throughout the neuron, can explain why opioids attenuate both the excitability of the peripheral nociceptive terminals and the propagation of action potentials (Andreev N, et al., Neuroscience 1994;58:793-8; Russell NJW, et al., Neurosci Lett 1987;76:107-12). At central nerve terminals, (Yaksh TL, et al., Nature 1980;286:155-7) opioids inhibit the calcium-dependent release of excitatory, pro-inflammatory compounds (e.g., substance P) from peripheral sensory-nerve endings, (Yaksh TL., Brain Res 1988 458:319-24)

which contribute to the anti-inflammatory actions of opioids (Barber A, Gottschlich R. et al., Med Res Rev 1992;12:525-62).

[39] There are three known opioid receptors, μ , κ , and δ -opioid receptors. μ -Receptors are further subdivided into at least two subclasses, $\mu 1$ and $\mu 2$ -receptors. The body produces opioid like molecules, called endogenous opioids, such as endorphins, enkephalins, and dynorphins. μ -receptors are known to mediate analgesia, respiratory depression, bradycardia, nausea/vomiting, and decreased gastrointestinal tone.

[40] δ -receptors mediate supraspinal analgesia, and κ -receptors mediate dysphoria and tachycardia. As pain impulses ascend through the spinal and supraspinal nervous system, activation of the opioid receptors inhibits these impulses and inhibits the transmission of pain from the dorsal horn. In addition, opioid analgesics prevent the presynaptic release of pain mediators such as Substance P into the spinal cord region.

[41] All animals, such as mammals, such as human, contain opioid receptors. It is understood that there are homologs for the various opioid receptors across animal species. A number of different opioid receptor sequences are set forth in the SEQ IDS, including μ -opioid receptors. The human μ -opioid receptor is referred to herein as HUMOR. It is understood that if a particular statement or reference is made regarding HUMOR that this statement is equally applicable to homologous receptors, unless specifically indicated otherwise.

[42] Opioid analgesics are typically grouped into three classes: naturally occurring (morphine, codeine); semi-synthetic morphine derivatives (hydromorphone, oxycodone, hydrocodone); and synthetic. Synthetic opioid analgesics include morphinan derivatives (levorphanol); methadone derivatives (methadone, propoxyphene); benzomorphan derivatives (pentazocine); and phenylpiperidine derivatives (meperidine, fentanyl, sufentanil, alfentanil, remifentanil). Tramadol is an opioid analgesic that also inhibits the reabsorption of norepinephrine and serotonin.

[43] Another way to classify opioid analgesics is as agonists, partial agonists, mixed agonists/antagonists, and antagonists based on their interactions at the opioid receptors. μ , and κ opioid-receptors are stimulated by agonists. Partial agonists have reduced μ -opioid receptor activity, and mixed agonists/antagonists only stimulate certain μ and κ -opioid receptors. Antagonists bind μ and κ -opioid receptors but do not stimulate the receptor activity.

[44] Some agonists are Morphine, Hydromorphone, Oxymorphone, Codeine, Oxycodone, Hydrocodone, Dihydrocodeine, Methadone, Meperidine, Fentanyl, Sufentanil, Alfentanil, and Remifentanil. An example of a partial agonist is Buprenorphine. Pentazocine, Nalbuphine, and Butorphanol are examples of mixed agonists/antagonists. Examples of
5 antagonists are Naloxone and Nalmefene. It is understood that one way to classify opioid receptors is by which molecules act as antagonists and which act as agonists, for example. Thus, a receptor can be defined as "a receptor for which morphine is an agonist."

[45] There are a number of adverse side effects that can occur when taking opioid analgesics, such as CNS effects, such as sedation, confusion, and respiratory depression.
10 Gastrointestinal adverse effects include nausea, vomiting, and constipation. Involuntary muscular contractions (twitching) known as myoclonus, bradycardia, and hypotension, can also occur. Lastly, physical and psychological dependence can also occur upon use or prolonged use of opioid analgesics. Thus there is a significant need for the disclosed compositions and methods, which reduce or eliminate the need for opioid analgesics in many indications.

15 **e) μ -opioid receptor therapy**

[46] The disclosed approach for the management of pain involves the targeted expression of opioid receptor(s) such as the μ -opioid receptor in the primary neurons innervating the areas, such as orofacial areas, experiencing pain, resulting in these same neurons becoming more susceptible to the intrinsic opioid anti-nociceptive mechanisms. Disclosed are
20 compositions and methods for treating pain. The compositions comprise an opioid receptor that is expressed from a vector. Typically these compositions will be delivered to at the point of pain. It is thought that their expression, increases the efficiency of the body's own opioid like molecules and decreases pain.

[47] Disclosed herein, the cDNA for a human μ -opioid receptor (HUMOR) is set
25 forth in SEQ ID NO:2. The μ -opioid receptor (Raynor K, et al., J Pharmacol Exp Ther. 1995; 272:423-8) has been placed into a vector herein and expressed in primary fibroblasts as well as cells of the N2a neuronal cell line (Figure 1). Transduction and stable expression of μ -opioid receptor in neurons can be accomplished by employing VSV-G pseudotyped immunodeficiency viral vectors (FIV).

30 [48] The expression of the μ -opioid receptor in the neurons at the point of pain in certain embodiments requires transduction in a non-dividing cell such as a neuron. This can be

accomplished using a transduction mechanism, such as lipofection or encapsulation methods, or via viral vector systems that function with cell division, such as lentiviruses, such as the FIV virus, or adeno-associated viruses, rAAV vectors, HSV Amplicon, and liposomes.

[49] It has been previously shown that this FIV system is capable, due to its lentiviral properties, of infecting terminally differentiated cells, including neurons. Disclosed are methods for administering vectors, such as the FIV(μ -opioid receptor) vector, peripherally at the site of pain. The neurons innervating that specific site and mediating the nociceptive signals are infected and stably transduced. These vectors, including vectors expressing lacZ and the μ -opioid receptor, can transduce nerve cells in vivo, in mice, through injection at the periphery.

[50] Disclosed herein is the stable expression of a reporter gene, the lacZ gene, in neurons located in the appropriate region of the trigeminal ganglion following peripheral injection of FIV(lacZ) in the area of the TMJ (Figure 2), as well as a variety of expression vectors containing the μ -opioid receptor, such as the human μ -opioid receptor.

[51] Disclosed are vectors wherein the vector includes sequence encoding the μ -opioid receptor gene. Also disclosed are vectors, wherein a μ -opioid receptor gene has been cloned in an FIV vector. Disclosed are methods comprising administering the disclosed vectors to cells, including cells involved in transmitting pain signals, such as nerve cells in the orofacial regions, related to for example, pain from TMJ and the masseter muscle.

[52] Also disclosed are transgenic mice that have been stably transfected with the disclosed vectors. These mice can be used, for example, as models of pain and the testing of therapeutics.

f) Mouse model of experimental nociception

[53] The majority of models evaluate reflex increases in jaw muscle activity, activating putative nociceptive pathways by the injection of algescic substances. Broton and Sessle (Botton JG, Sessle BJ, et al., Arch Oral Biol 1988;33:741-47.) placed hypertonic saline, potassium chloride and histamine into the TMJ of cats and found increase in electromyographic (EMG) activity in the temporal, digastric and genioglossus muscles. This suggests that the products of tissue injury and or inflammation are capable of producing pain within the TMJ, causing excitation of sensory afferents and reflex muscle activity. Tambeli et al. (Tambeli CH, et al., *J Dent Res* 1997; 76: (Special Issue) abstr# 1263.) injected mustard oil unilaterally into the TMJ and the results were similar to Broton and Sessle (Botton JG, Sessle BJ, et al., Arch Oral

Biol 1988;33:741-47.). Increases in masticatory muscle activity have been also demonstrated by injection of excitatory amino acids into the TMJ area. Cairns et al. (Cairns BE, et al., J Neurosci 18;1998:8056-64. Cairns BE, et al., J Neurophysiol 1999; 81: 1966-69.) injected glutamate into the TMJ area of rats and in a dose dependent manner induced a prolonged
5 increase in EMG activity by the excitation of nociceptors (A delta and C fibers) (Cairns BE, et al., J Neurophysiol 2001; 86: 782-90). Cairns et al. (Cairns BE, et al., J Neurophysiol 85;2001:2446-54.) placed glutamate into the masseter muscle of the rat and human and evoked higher masseter muscle activity in female than males of both species. These results indicate that peripheral insult can produce pain behavior and changes in resting muscle activity when
10 algogenic substances are injected into the TMJ and masticatory muscles. Disclosed are behavioral models to assess orofacial pain from the TMJ and masseter muscle via application of glutamate into the TMJ and masticatory muscles. These models can include mice, disclosed herein, which have had vectors encoding μ -opioid receptor stably transduced. Resistance to jaw opening and EMG activity can serve as behavioral measures of pain.

15 C. Compositions

1. opioid receptors

[54] There are typically considered three classes of opioid receptor μ , δ and κ . Genes encoding for these receptors have been cloned (Evans et al (1992) Science 258 1952; Kieffer et al (1992) Proc.Natl.Acad.Sci.USA 89 12048; Chen et al (1993) Mol.Pharmacol. 44 8; and
20 Minami et al (1993) FEBS Lett. 329 291 all of which are herein incorporated by reference for material related to opioid receptors and there sequence). In addition, an orphan receptor was identified which has a high degree of homology to the known opioid receptors and based on structural grounds it is considered a receptor called ORL1 (opioid receptor-like) (Mollereau et al (1994) FEBS Lett. 341 33, herein incorporated by reference for material related to opioid
25 receptors and there sequence). Since the cloned receptors function as opioid receptors, by for example interacting with pertussis toxin-sensitive G-proteins, all of the cloned opioid receptors possess the same general structure which includes an extracellular N-terminal region, seven transmembrane domains and intracellular C-terminal tail structure. Evidence obtained from pharmacokinetic and activity data indicate there are subtypes of each receptor and other types,
30 such as less well-characterized opioid receptors, such as ϵ , λ , ι , ζ , which are known. One way of characterizing the different receptor subtypes for μ -, δ - and κ -receptors is through different post-translational modifications of the gene product (glycosylation, palmitoylation,

phosphorylation, etc). Also receptor dimerization to form homomeric and heteromeric complexes or from interaction of the gene product with associated proteins such as RAMPs can effect function, and thus represent another way to characterize the receptors. Different opioids have different affinity for the different opioid receptors. For example, μ -morphine, δ -leukenkephalin metenkephalin, κ -dynorphin, β -endorphin, have different affinities for the various opioid receptors.

a) μ -Receptor subtypes

[55] The MOR-1 gene, encoding for one form of the μ -receptor, shows approximately 50-70% homology to the genes encoding for the δ -(DOR-1), κ -(KOR-1) and orphan (ORL1) receptors. Two different splice variants of the MOR-1 gene have been cloned, and they differ by 8 amino acids in the C-terminal tail which are either present or not. The splice variants exhibit differences in their rate of onset and recovery from agonist-induced internalization but their pharmacology does not appear to differ in ligand binding assays. A MOR-1 knockout mouse has been made and the mouse does not respond to morphine, by failing to alleviate pain, and by failing to exhibit positive reinforcing properties or an ability to induce physical dependence in the absence of the MOR-1 gene. This indicates that at least in this species, morphine's analgesia is not mediated through δ - or κ -receptors. (Matthes et al (1996) Nature 383 818).

[56] The μ receptor is divided into the $\mu 1$ and $\mu 2$ groups. The division occurs because of binding and pharmacology activity studies which indicate, for example, that naloxazone and naloxonazine abolish the binding of radioligands to the $\mu 1$ -site, and in vivo studies showed that naloxazone selectively blocked morphine-induced antinociception but did not block morphine-induced respiratory depression or the induction of morphine dependence, indicating different types of μ -receptor (Ling et al (1984) Science 226 462 and Ling et al (1985) J.Pharmacol.Exp.Ther. 232 149). Subsequent work in other laboratories has failed to confirm this classification.

[57] Peptide sequences of the human and mouse μ receptor are set forth in SEQ ID Nos 1 and 3 respectively.

[58] There is also data consistent with a third form of μ receptor where analogues of morphine with substitutions at the 6 position (e.g. morphine-6b-glucuronide, heroin and 6-acetyl morphine) are agonists, but with which morphine itself does not interact (Rossi et al

(1996) Neuroscience Letters 216 1, herein incorporated by reference for material at least related to opioid receptors and their function and structure). Antinociception tests on mice show that morphine does not exhibit cross tolerance with morphine-6b-glucuronide, heroin or 6-acetyl morphine. Furthermore, in mice of the CXBX strain morphine is a poor antinociceptive agent
 5 whereas morphine-6b-glucuronide, heroin and 6-acetyl morphine are all potently antinociceptive. Heroin and morphine-6-glucuronide, but not morphine, still produce antinociception in MOR-1 knockout mice in which the disruption in the MOR-1 gene was engineered in exon-1 (Schuller et al (1999) Nature Neuroscience 2 151). Furthermore, all three agonists were ineffective as antinociceptive agents, in MOR-1 knockout mice in which exon-2,
 10 not exon-1, had been disrupted. This indicates that the antinociceptive actions of heroin and morphine-6-glucuronide in the exon-1 MOR-1 mutant mice are mediated through a receptor produced from an alternative transcript of the MOR-1 gene differing from the MOR-1 gene product, the μ -opioid receptor, in the exon-1 region.

b) δ -Receptor subtypes

15 [59] Only one δ -receptor gene has been cloned (DOR-1), but overlapping subdivisions of δ -receptor have been proposed ($\delta 1/\delta 2$ and $\delta cx/\delta ncx$) on the basis of in vivo and in vitro pharmacological experiments.

[60] The δ receptor subclasses arise from pharmacological studies. Results from in vivo rodent studies are shown in Table 1.

20 [61] Table 1.

	Agonist	Competitive antagonist	Non-competitive antagonist
$\delta 1$	DPDPE / DADLE	BNTX (7-benzylidenenaltrexone)	DALCE ([D-Ala2, D-Leu5]enkephalyl-Cys)
$\delta 2$	Deltorphin II / DSLET	Naltriben	5'-NTII (naltrindole 5'-isothiocyanate)

[62] There are a number of different ligands for the opioid receptors which differentially bind one or more receptors. Examples of these ligands are shown in Table 2.

Receptor type	μ -Receptor	δ -Receptor	κ -Receptor	ORL ₁
---------------	-----------------	--------------------	--------------------	------------------

Selective agonists	endomorphin-1 endomorphin-2 DAMGO	[D-Ala ²]-deltorphin I [D-Ala ²]-deltorphin II DPDPE SNC 80	enadoline U-50488 U-69593	nociceptin / OFQAc-RYYRWK-NH ₂ *
Selective antagonists CTAP naltrindole TIPP	yICI 174864 nor	binaltorphimine	Selective antagonists CTAP naltrindole TIPP	None as yet**
Radioligands	[³ H]	DAMGO [³ H]	[³ H]-enadoline [³ H]-U69593	[³ H]-nociceptin

[63] Table 2

[64] The pharmacological properties of the cloned DOR-1 receptor are somewhere between those predicted for either the $\delta 1$ or $\delta 2$ subtypes. Mouse and human recombinant receptors both bind DPDPE and deltorphin II, which can displace of [³H]-diprenorphine. This is different than either a $\delta 1$ or $\delta 2$ classification (Law et al (1994) J.Pharmacol.Exp.Ther. 271 1686). [³H]-diprenorphine, binding to the mouse recombinant receptor, however, is more highly displaced by naltriben than BNTX, consistent with it being $\delta 2$ like.

[65] Opioid receptors have also been indicated to be in complex μ -receptors and κ -receptors. For example, one type of δ receptor subtypes complexes, $\delta c x$, and another appears not to complex, $\delta n c x$ (Rothman et al (1993) In: Handbook Exp.Pharmacol. Ed. A. Herz 104/1 p217).

c) κ -Receptor

[66] The cloned κ -Receptor has the sequence set forth in SEQ ID NO: 5, which represents an example of a κ -receptor.

d) The orphan opioid receptor

[67] The orphan receptor has been identified in three species: rat, mouse and man, all having a greater than 90% identity with each other. This receptor is typically referred to as ORL-1 for orphan receptor like 1. The endogenous peptide agonist for ORL1 is known as nociceptin or orphanin FQ. While the ORL1 receptor has structural homology to orphan receptors it does not have pharmacological homology. Non-selective ligands that exhibit high affinity for all μ -, κ - and δ -receptors, have very low affinity for the ORL1 receptor. Comparison

of the deduced amino-acid sequences of the four receptors highlights structural differences consistent with the lack of coligand binding. The trans-membrane regions are conserved near their top in the μ -, κ - and δ -receptors, but are altered in ORL1. Site-directed mutants of ORL1 towards the traditional receptors (rat) are able to bind the traditional receptor's ligands. For example, benzomorphan bremazocine binds ORL1 by changing Ala213 in TM5 to the conserved Lys of μ , κ and δ , or by changing the Val-Gln-Val276-278 sequence of TM6 to the conserved Ile-His-Ile motif (Meng et al (1996) J.Biol.Chem. 271 32016). There are also a number of splice variants of the ORL1 receptor, XOR (Wang et al (1994) FEBS Lett. 348 75) with a long form (XOR1L) containing an additional 28 amino acids in the third extracellular loop and in the homologous receptor from mouse, KOR-3, five splice variants have been reported to date (Pan et al (1998) FEBS Lett. 435 65).

e) Endogenous Ligands

[68] In mammals the endogenous opioid peptides are mainly derived from four precursors: pro-opiomelanocortin, pro-enkephalin, pro-dynorphin and pro-nociceptin/orphanin FQ. Nociceptin/orphanin FQ is processed from pro-nociceptin/orphanin FQ and is the endogenous ligand for the ORL1-receptor; it has little affinity for the μ -, δ - and κ -receptors. Table 3 sets forth endogenous ligands for the opioid receptors. These peptides bind μ , δ - and κ -receptors with different affinity, and have negligible affinity for ORL1-receptors, but none binds exclusively to one opioid receptor type. β -endorphin is equiactive at μ - and δ -receptors with much lower affinity for κ -receptors; the post-translational product, N-acetyl- β -endorphin, has very low affinity for any of the opioid receptors. [Met]- and [Leu]enkephalin have high affinities for δ -receptors, ten-fold lower affinities for μ -receptors and negligible affinity for κ -receptors. Other products of processing of pro-enkephalin, which are N-terminal extensions of [Met]enkephalin, have a decreased preference for the δ -receptor with some products, e.g. metorphamide displaying highest affinity for the μ -receptor. The opioid fragments of pro-dynorphin, particularly dynorphin A and dynorphin B, have high affinity for κ -receptors but also have significant affinity for μ - and δ -receptors.

[69] Endomorphin-1 and endomorphin-2 are putative products of an as yet unidentified precursor, that have been proposed to be the endogenous ligands for the μ -receptor where they are highly selective. The endomorphins are amidated tetrapeptides and are structurally unrelated to the other endogenous opioid peptides (Table 3). Although the study of the cellular localization of these peptides is at an early stage, endomorphin-2 is found in discrete

regions of rat brain, some of which are known to contain high concentrations of μ -receptors. Endomorphin-2 is also present in primary sensory neurones and the dorsal horn of the spinal cord where it could function to modulate nociceptive input.

[70] In comparison to the mainly non-selective mammalian opioid peptides (the exceptions being the endomorphins), amphibian skin contains two families of D-amino acid-containing peptides that are selective for μ - or δ -receptors. Dermorphin is a μ -selective heptapeptide Tyr-D-Ala-Phe-Gly-Tyr-Pro-Ser-NH₂ without significant affinity at δ - and κ -receptors. In contrast, the deltorphins - deltorphin (dermenkephalin; Tyr-D-Met-Phe-His-Leu-Met-Asp-NH₂), [D-Ala²]-deltorphan I and [D-Ala²]-deltorphan II (Tyr-D-Ala-Phe-Xaa-Val-Val-Gly-NH₂, where Xaa is Asp or Glu respectively) - are highly selective for δ -opioid receptors. Table 3 shows a variety of endogenous opioid receptor molecules.

[71] Table 3.

Precursor	Endogenous peptide	Amino acid sequence
Pro-opiomelanocortin	β -Endorphin	YGGFMTSEKSQTPLVTL-FKNAIIKNAYKKGE
Pro-enkephalin	[Met]enkephalin [Leu]enkephalin Metorphamide	YGGFM YGGFL YGGFMRF YGGFMRGL YGGFMRRV-NH ₂
Pro-dynorphin	Dynorphin A Dynorphin A(1-8) Dynorphin B α -neoendorphin β -neoendorphin	YGGFLRRIRPKLKWDNQ YGGFLRRI YGGFLRRQFKVVT YGGFLRKYPK YGGFLRKYP
Pro-nociceptin / OFQ	Nociceptin	FGGFTGARKSARKLANQ
Pro-endomorphin*	Endomorphin-1 Endomorphin-2	YPWF-NH ₂ YPFF-NH ₂

[72] Opioid receptor activation produces a wide array of cellular responses (Table 4). For example, there are Direct G-protein $\beta\gamma$ or α subunit-mediated effects, such as activation of an inwardly rectifying potassium channel, inhibition of voltage operated calcium channels (N, P, Q and R type), inhibition of adenylyl cyclase, Responses of unknown intermediate mechanism, activation of PLA₂, activation of PLC β (possibly direct G protein $\beta\gamma$ subunit activation), activation of MAPKinase, activation of large conductance calcium-activated potassium channels, activation of L type voltage operated calcium channels, inhibition of T type voltage operated calcium channels, and direct inhibition of transmitter exocytosis. There are also

responses in other effector pathways, such as activation of voltage-sensitive potassium channels (activation of PLA2), inhibition of M channels (activation of PLA2), inhibition of the hyperpolarisation-activated cation channel (Ih) (reduction in cAMP levels following inhibition of adenylyl cyclase), elevation of intracellular free calcium levels (activation of PLCb, activation of L type voltage operated calcium conductance), potentiation of NMDA currents (activation of protein kinase C), inhibition of transmitter release (inhibition of adenylyl cyclase, activation of potassium channels and inhibition of voltage operated calcium channels), decreases in neuronal excitability (activation of potassium channels), increases in neuronal firing rate (inhibition of inhibitory transmitter release - disinhibition), and changes in gene expression (long-term changes in adenylyl cyclase activity, elevation of intracellular calcium levels, activation of cAMP response element binding protein (CREB)).

2. Compositions for treating pain

[73] Disclosed are constructs and vectors for expressing one or more opioid receptors in a cell, such as a nerve cell, such as a peripheral nerve cell. As discussed herein, opioid receptors are typically expressed in the spinal or supraspinal nerve cells, and the periphery. typically do not express these receptors. The disclosed compositions and methods are designed to express the opioid receptors in nerve cells which are damaged or transmitting because of trauma, but which do not have endogenous opioid receptors or insufficient numbers of endogenous receptors to react to the endogenous opioid like molecules, typically in the periphery of the nerve cell. Thus, the expression of the opioid receptors in the nerve cell near the point of pain, will typically increase the amount of opioid receptors in this area and thus, increase the responsiveness to endogenous opioid like molecules. By expression of the opioid receptors, the sensation of pain can be reduced, not by administration of opioid analgesics, but rather by more efficient use of endogenous opioid like compounds. It is understood, however, that opioids, opioid like molecules, and/or other pain alleviating molecules can be added in addition to the disclosed opioid receptors.

[74] Disclosed are methods wherein administration occurs in the intra-articular region of the jaw. The results shown herein demonstrated that intra-articular injection of FIV(lacZ) resulted in successful gene transfer to articular TMJ surfaces as well as the joint meniscus. Thus, disclosed are methods, wherein the administration of the disclosed vectors, results in delivery to the articular TMJ surfaces and the joint meniscus.

[75] Nociceptive innervation to the temporomandibular joint (TMJ) is primarily provided by the auriculotemporal nerve of the mandibular division of the trigeminal nerve (Sessle & Wu, 1991). A δ and C nerve fibers, whose cell bodies are located in the posterolateral part of the trigeminal ganglion (Yoshino et al., 1998), project distally and terminate as non-encapsulated free nerve endings dispersed throughout the posterolateral part of the TMJ capsule (Bernick, 1962; Thilander, 1964; Frommer & Monroe, 1966; Klineberg, 1971), the posterior band of the meniscus and the posterior attachment (Dressen et al., 1990; Kido et al., 1991, 1993; Wink et al., 1992). Transfer of anti-nociceptive genes to sensory trigeminal neurons innervating the orofacial region can be achieved after injection of lentiviral vectors at the painful site, such as the TMJ, resulting in their uptake by free nerve endings and retrograde transport to the sensory cells' nuclei. Previous studies demonstrated axonal retrograde transport of horseradish peroxidase from the TMJ to the central nervous system (Romfh et al., 1979; Carpa, 1987) including the trigeminal ganglia (Yoshino et al., 1998).

[76] Disclosed are constructs capable of expressing any of the opioid receptor gene products.

[77] Disclosed are constructs capable of expressing opioid receptors, such as the μ -opioid receptor gene product. The μ -opioid receptor construct allows for synthesis of μ -opioid receptor protein.

[78] The μ -opioid receptor construct typically comprises three parts: 1) a promoter, 2) the μ -opioid receptor coding sequence, and 3) polyA tail. The poly A tail can be from the bovine growth hormone or any polyA tail including synthetic poly A tails. The Bovine growth hormone poly A tail carries elements that not only increase expression, but also increase stability of any gene construct. These three parts can be integrated into any vector delivery system, which is capable of transducing terminally differentiated cells, such as nerve cells.

[79] The promoter can be any promoter, such as those discussed herein. It is understood as discussed herein that there are functional variants of opioid receptors, such as the μ -opioid receptor protein which can be made. In certain embodiments the promoter is going to be a cell specific promoter, such as a nerve cell specific promoter, such as the neuron specific enolase promoter. Other promoters are disclosed herein.

[80] The promoter can be any promoter, such as those discussed herein. It is understood as discussed herein that there are functional variants of opioid receptors, such as the

μ -opioid receptor protein which can be made. In certain embodiments the promoter is going to be a cell specific promoter, such as a nerve cell specific promoter, such as the neuron specific enolase promoter.

[81] μ -opioid receptor cDNA can be obtained from the American Tissue Culture
5 Collection. (American Tissue Culture Collection, Manassas, VA 20110-2209; μ -opioid receptor ATCC#. Raynor K, et al., Characterization of the cloned human mu opioid receptor. J Pharmacol Exp Ther. 1995; 272:423-8.)

[82] Also disclosed are constructs encoding for the human or mouse μ -opioid receptor, as well as the β -galactosidase reporter gene (*lacZ*).

10 [83] Disclosed are nucleic acids comprising sequence encoding μ -opioid receptor. Also disclosed are nucleic acids, wherein the nucleic acid further comprises a promoter sequence, wherein the μ -opioid receptor has at least 80% identity to the sequence set forth in SEQ ID NO:2 or 4, wherein the μ -opioid receptor has at least 85% identity to the sequence set forth in SEQ ID NO:1 or 3, wherein the μ -opioid receptor has at least 90% identity to
15 sequence set forth in SEQ ID NO:1 or 3, wherein the μ -opioid receptor has at least 95% identity to the sequence set forth in SEQ ID NO:1 or 3, and/or wherein the μ -opioid receptor has the sequence set forth in SEQ ID NO: 1 or 3.

[84] Also disclosed are vectors comprising the disclosed nucleic acids. Also disclosed are cells comprising the disclosed nucleic acids and vectors. Any cell can be targeted
20 with the disclosed constructs. However, nerve cells, for example, are terminally differentiated. This means that they are no longer dividing. The state of a mature non-dividing nerve cell can define terminally differentiated cells. In terms of differentiated/stable transduction, nerve cells thus represent attractive targets because once DNA is integrated, there is a very low probability that it will not remain in the cell.

25 [85] Also disclosed are non-human mammals comprising the disclosed nucleic acids, vectors, and cells disclosed herein.

[86] Also disclosed are methods of providing μ -opioid receptor in a cell comprising transfecting the cell with the nucleic acids.

[87] Also disclosed are method of delivering the disclosed compositions, wherein the
30 transfection occurs in vitro or in vivo.

[88] Disclosed are methods of making a transgenic organism comprising administering the disclosed nucleic acids, vectors and/or cells.

[89] Disclosed are methods of making a transgenic organism comprising transfecting a lentiviral vector to the organism at during a perinatal stage of the organism's development.

5 Strategies of producing genetically engineered pluripotent, such as embryonic, stem cells, can be performed with the disclosed compositions to produce engineered cells and organisms as discussed herein. Furthermore by cloning strategies can be used to produce desired organisms, which carry one or more of the disclosed compositions.

[90] Also disclosed are methods of treating a subject having pain comprising
10 administering any of the disclosed compounds and compositions.

[91] Delivery of the disclosed constructs to terminally differentiated cells is also disclosed. Disclosed is a pseudotyped feline immunodeficiency virus (FIV) for μ -opioid receptor delivery to terminally differentiated cells. Stable expression of the therapeutic gene aids prolonged expression, enhancing treatment efficacy and contributing to long-term
15 therapeutic outcomes. The backbone FIV system has been shown to effectively incorporate, due to its lentiviral properties, the transgene of interest into the host's genome, allowing for stable gene expression (Poeschla et al., 1998). Disclosed herein is stable expression of the reporter gene *lacZ* in N2a cells, following perinatal systemic FIV(*lacZ*) administration.

[92] In certain embodiments the constructs become an integrated product with the
20 genome of the host. For example, lentiviruses, such as HIV and LIV, have the characteristic of transfecting the therapeutic gene into the host chromosome, thus forming an integrated product. In certain embodiments, the requirement is that the vectors allow for expression in the periphery of the cell, such as the nerve cell, and/or at or near the point of pain. The contrast to integrated products is episomal products which can also be produced using, for example, HSV
25 and AV vectors. Thus, transient expression can be beneficial. The optimal time of expression is correlated with the amount of product produced and amount that is needed. For example, in certain embodiments, expression for at least 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 45, 60, 90, 120, 150, or 180 days is desirable.

[93] A model system for the study of these vectors is a mouse that is knockout mouse
30 deficient in μ -opioid receptor.

[94] Stable transduced are cells where a nucleic acid has been integrated into the cell genome.

3. Delivery of the compositions to cells

[95] Delivery can be applied, in general, via local or systemic routes of administration. Local administration includes virus injection directly into the region or organ of interest, versus intravenous (*IV*) or intraperitoneal (*IP*) injections (systemic) aiming at viral delivery to multiple sites and organs via the blood circulation. Previous research on the effects of local administration demonstrated gene expression limited to the site/organ of the injection, which did not extend to the rest of the body (Daly et al., 1999a; Kordower et al., 1999). Furthermore, previous studies have demonstrated successful global gene transfer to multiple tissues and organs in rodents and primates following viral *IV* and *IP* injections (Daly et al., 1999b; Tarntal et al., 2001; McCormack et al., 2001; Lipschutz et al., 2001). Disclosed herein *IP* injection of FIV(lacZ) in mice of adult (3 months old) as well as of perinatal age (P4) resulted in global transfer and expression of the reporter gene lacZ in brain, liver, spleen and kidney. Also disclosed, the levels of expression achieved via *IP* injections were superior to those acquired following local administration directly into the liver.

[96] There are a number of compositions and methods which can be used to deliver nucleic acids to cells, either in vitro or in vivo. These methods and compositions can largely be broken down into two classes: viral based delivery systems and non-viral based delivery systems. For example, the nucleic acids can be delivered through a number of direct delivery systems such as, electroporation, lipofection, calcium phosphate precipitation, plasmids, viral vectors, viral nucleic acids, phage nucleic acids, phages, cosmids, or via transfer of genetic material in cells or carriers such as cationic liposomes. Appropriate means for transfection, including viral vectors, chemical transfectants, or physico-mechanical methods such as electroporation and direct diffusion of DNA, are described by, for example, Wolff, J. A., et al., Science, 247, 1465-1468, (1990); and Wolff, J. A. Nature, 352, 815-818, (1991). Such methods are well known in the art and readily adaptable for use with the compositions and methods described herein. In certain cases, the methods will be modified to specifically function with large DNA molecules. Further, these methods can be used to target certain diseases and cell populations by using the targeting characteristics of the carrier.

a) Nucleic acid based delivery systems

[97] Transfer vectors can be any nucleotide construction used to deliver genes into cells (e.g., a plasmid), or as part of a general strategy to deliver genes, e.g., as part of recombinant retrovirus or adenovirus (Ram et al. Cancer Res. 53:83-88, (1993)).

5 [98] As used herein, plasmid or viral vectors are agents that transport the disclosed nucleic acids, such as the μ -opioid receptor construct into the cell without degradation and include a promoter yielding expression of the μ -opioid receptor encoding sequences in the cells into which it is delivered. In some embodiments the vectors for the μ -opioid receptor constructs are derived from either a virus, retrovirus, or lentivirus. Viral vectors can be, for example,
10 Adenovirus, Adeno-associated virus, Herpes virus, Vaccinia virus, Polio virus, AIDS virus, neuronal trophic virus, Sindbis and other RNA viruses, including these viruses with the HIV backbone, and lentiviruses. Also preferred are any viral families which share the properties of these viruses which make them suitable for use as vectors. Retroviruses include Murine Maloney Leukemia virus, MMLV, and retroviruses that express the desirable properties of
15 MMLV as a vector. Retroviral vectors are able to carry a larger genetic payload, i.e., a transgene, such as, the disclosed μ -opioid receptor constructs or marker gene, than other viral vectors, and for this reason are a commonly used vector. However, they are not as useful in non-proliferating cells. Adenovirus vectors are relatively stable and easy to work with, have high titers, and can be delivered in aerosol formulation, and can transfect non-dividing cells.
20 Pox viral vectors are large and have several sites for inserting genes, they are thermostable and can be stored at room temperature. A preferred embodiment is a viral vector, which has been engineered so as to suppress the immune response of the host organism, elicited by the viral antigens. Preferred vectors of this type will carry coding regions for Interleukin 8 or 10.

[99] Viral vectors can have higher transaction (ability to introduce genes) abilities
25 than chemical or physical methods to introduce genes into cells. Typically, viral vectors contain, nonstructural early genes, structural late genes, an RNA polymerase III transcript, inverted terminal repeats necessary for replication and encapsidation, and promoters to control the transcription and replication of the viral genome. When engineered as vectors, viruses typically have one or more of the early genes removed and a gene or gene/promotor cassette is
30 inserted into the viral genome in place of the removed viral DNA. Constructs of this type can carry up to about 8 kb of foreign genetic material. The necessary functions of the removed early

genes are typically supplied by cell lines which have been engineered to express the gene products of the early genes in trans.

(1) Retroviral Vectors

[100] A retrovirus is an animal virus belonging to the virus family of Retroviridae, including any types, subfamilies, genus, or tropisms. Retroviral vectors, in general, are described by Verma, I.M., Retroviral vectors for gene transfer. In Microbiology-1985, American Society for Microbiology, pp. 229-232, Washington, (1985), which is incorporated by reference herein. Examples of methods for using retroviral vectors for gene therapy are described in U.S. Patent Nos. 4,868,116 and 4,980,286; PCT applications WO 90/02806 and WO 89/07136; and Mulligan, (Science 260:926-932 (1993)); the teachings of which are incorporated herein by reference.

[101] A retrovirus is essentially a package which has packed into it nucleic acid cargo. The nucleic acid cargo carries with it a packaging signal, which ensures that the replicated daughter molecules will be efficiently packaged within the package coat. In addition to the package signal, there are a number of molecules which are needed in cis, for the replication, and packaging of the replicated virus. Typically a retroviral genome, contains the gag, pol, and env genes which are involved in the making of the protein coat. It is the gag, pol, and env genes which are typically replaced by the foreign DNA that it is to be transferred to the target cell. Retrovirus vectors typically contain a packaging signal for incorporation into the package coat, a sequence which signals the start of the gag transcription unit, elements necessary for reverse transcription, including a primer binding site to bind the tRNA primer of reverse transcription, terminal repeat sequences that guide the switch of RNA strands during DNA synthesis, a purine rich sequence 5' to the 3' LTR that serve as the priming site for the synthesis of the second strand of DNA synthesis, and specific sequences near the ends of the LTRs that enable the insertion of the DNA state of the retrovirus to insert into the host genome. The removal of the gag, pol, and env genes allows for about 8 kb of foreign sequence to be inserted into the viral genome, become reverse transcribed, and upon replication be packaged into a new retroviral particle. This amount of nucleic acid is sufficient for the delivery of a one to many genes depending on the size of each transcript. It is preferable to include either positive or negative selectable markers along with other genes in the insert.

[102] Since the replication machinery and packaging proteins in most retroviral vectors have been removed (gag, pol, and env); the vectors are typically generated by placing them into

a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals.

(2) Adenoviral Vectors

[103] The construction of replication-defective adenoviruses has been described (Berkner et al., *J. Virology* 61:1213-1220 (1987); Massie et al., *Mol. Cell. Biol.* 6:2872-2883 (1986); Haj-Ahmad et al., *J. Virology* 57:267-274 (1986); Davidson et al., *J. Virology* 61:1226-1239 (1987); Zhang "Generation and identification of recombinant adenovirus by liposome-mediated transfection and PCR analysis" *BioTechniques* 15:868-872 (1993)). The benefit of the use of these viruses as vectors is that they are limited in the extent to which they can spread to other cell types, since they can replicate within an initial infected cell, but are unable to form new infectious viral particles. Recombinant adenoviruses have been shown to achieve high efficiency gene transfer after direct, in vivo delivery to airway epithelium, hepatocytes, vascular endothelium, CNS parenchyma and a number of other tissue sites (Morsy, *J. Clin. Invest.* 92:1580-1586 (1993); Kirshenbaum, *J. Clin. Invest.* 92:381-387 (1993); Roessler, *J. Clin. Invest.* 92:1085-1092 (1993); Moullier, *Nature Genetics* 4:154-159 (1993); La Salle, *Science* 259:988-990 (1993); Gomez-Foix, *J. Biol. Chem.* 267:25129-25134 (1992); Rich, *Human Gene Therapy* 4:461-476 (1993); Zabner, *Nature Genetics* 6:75-83 (1994); Guzman, *Circulation Research* 73:1201-1207 (1993); Bout, *Human Gene Therapy* 5:3-10 (1994); Zabner, *Cell* 75:207-216 (1993); Caillaud, *Eur. J. Neuroscience* 5:1287-1291 (1993); and Ragot, *J. Gen. Virology* 74:501-507 (1993)). Recombinant adenoviruses achieve gene transduction by binding to specific cell surface receptors, after which the virus is internalized by receptor-mediated endocytosis, in the same manner as wild type or replication-defective adenovirus (Chardonnet and Dales, *Virology* 40:462-477 (1970); Brown and Burlingham, *J. Virology* 12:386-396 (1973); Svensson and Persson, *J. Virology* 55:442-449 (1985); Seth, et al., *J. Virol.* 51:650-655 (1984); Seth, et al., *Mol. Cell. Biol.* 4:1528-1533 (1984); Varga et al., *J. Virology* 65:6061-6070 (1991); Wickham et al., *Cell* 73:309-319 (1993)).

[104] A viral vector can be one based on an adenovirus which has had the E1 gene removed and these virions are generated in a cell line such as the human 293 cell line. In another preferred embodiment both the E1 and E3 genes are removed from the adenovirus genome.

5

(3) Adeno-associated viral vectors

[105] Another type of viral vector is based on an adeno-associated virus (AAV). This defective parvovirus is a preferred vector because it can infect many cell types and is nonpathogenic to humans. AAV type vectors can transport about 4 to 5 kb and wild type AAV is known to stably insert into chromosome 19. Vectors which contain this site specific
10 integration property are preferred. An especially preferred embodiment of this type of vector is the P4.1 C vector produced by Avigen, San Francisco, CA, which can contain the herpes simplex virus thymidine kinase gene, HSV-tk, and/or a marker gene, such as the gene encoding the green fluorescent protein, GFP.

[106] In another type of AAV virus, the AAV contains a pair of inverted terminal
15 repeats (ITRs) which flank at least one cassette containing a promoter which directs cell-specific expression operably linked to a heterologous gene. Heterologous in this context refers to any nucleotide sequence or gene which is not native to the AAV or B19 parvovirus.

[107] Typically the AAV and B19 coding regions have been deleted, resulting in a safe, noncytotoxic vector. The AAV ITRs, or modifications thereof, confer infectivity and site-
20 specific integration, but not cytotoxicity, and the promoter directs cell-specific expression. United states Patent No. 6,261,834 is herein incorporated by reference for material related to the AAV vector.

[108] The vectors of the present invention thus provide DNA molecules which are capable of integration into a mammalian chromosome without substantial toxicity.

[109] The inserted genes in viral and retroviral usually contain promoters, and/or
25 enhancers to help control the expression of the desired gene product. A promoter is generally a sequence or sequences of DNA that function when in a relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of RNA polymerase and transcription factors, and can contain upstream elements and response
30 elements.

(4) Lentiviral vectors

[01] The vectors can be lentiviral vectors, including but not limited to, SIV vectors, HIV vectors or a hybrid construct of these vectors, including viruses with the HIV backbone. These vectors also include first, second and third generation lentiviruses. Third generation lentiviruses have lentiviral packaging genes split into at least 3 independent plasmids or
5 constructs. Also vectors can be any viral family that shares the properties of these viruses which make them suitable for use as vectors. Lentiviral vectors are a special type of retroviral vector which are typically characterized by having a long incubation period for infection. Furthermore, lentiviral vectors can infect non-dividing cells. Lentiviral vectors are based on the nucleic acid backbone of a virus from the lentiviral family of viruses. Typically, a lentiviral vector contains
10 the 5' and 3' LTR regions of a lentivirus, such as SIV and HIV. Lentiviral vectors also typically contain the Rev Responsive Element (RRE) of a lentivirus, such as SIV and HIV.

(a) Feline immunodeficiency viral vectors

[110] One type of vector that the disclosed constructs can be delivered in is the VSV-G pseudotyped Feline Immunodeficiency Virus system developed by Poeschla *et al.* (1998). This
15 lentivirus has been shown to efficiently infect dividing, growth arrested as well as post-mitotic cells. Furthermore, due to its lentiviral properties, it allows for incorporation of the transgene into the host's genome, leading to stable gene expression. This is a 3-vector system, whereby each confers distinct instructions: the FIV vector carries the transgene of interest and lentiviral apparatus with mutated packaging and envelope genes. A vesicular stomatitis virus G-
20 glycoprotein vector (VSV-G; Burns *et al.*, 1993) contributes to the formation of the viral envelope *in trans*. The third vector confers packaging instructions *in trans* (Poeschla *et al.*, 1998). FIV production is accomplished *in vitro* following co-transfection of the aforementioned vectors into 293-T cells. The FIV-rich supernatant is then collected, filtered and can be used directly or following concentration by centrifugation. Titers routinely range between $10^4 - 10^7$
25 bfu/ml..

(5) Packaging vectors

[111] As discussed above, retroviral vectors are based on retroviruses which contain a number of different sequence elements that control things as diverse as integration of the virus, replication of the integrated virus, replication of un-integrated virus, cellular invasion, and
30 packaging of the virus into infectious particles. While the vectors in theory could contain all of their necessary elements, as well as an exogenous gene element (if the exogenous gene element is small enough) typically many of the necessary elements are removed. Since all of the

packaging and replication components have been removed from the typical retroviral, including lentiviral, vectors which will be used within a subject, the vectors need to be packaged into the initial infectious particle through the use of packaging vectors and packaging cell lines.

Typically retroviral vectors have been engineered so that the myriad functions of the retrovirus are separated onto at least two vectors, a packaging vector and a delivery vector. This type of system then requires the presence of all of the vectors providing all of the elements in the same cell before an infectious particle can be produced. The packaging vector typically carries the structural and replication genes derived from the retrovirus, and the delivery vector is the vector that carries the exogenous gene element that is preferably expressed in the target cell. These types of systems can split the packaging functions of the packaging vector into multiple vectors, e.g., third-generation lentivirus systems. Dull, T. et al., "A Third-generation lentivirus vector with a conditional packaging system" J. Virol 72(11):8463-71 (1998)

[112] Retroviruses typically contain an envelope protein (env). The Env protein is in essence the protein which surrounds the nucleic acid cargo. Furthermore cellular infection specificity is based on the particular Env protein associated with a typical retrovirus. In typical packaging vector/delivery vector systems, the Env protein is expressed from a separate vector than for example the protease (pro) or integrase (in) proteins.

(6) Packaging cell lines

[113] The vectors are typically generated by placing them into a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals. One type of packaging cell line is a 293 cell line.

(7) Large payload viral vectors

[114] Molecular genetic experiments with large human herpesviruses have provided a means whereby large heterologous DNA fragments can be cloned, propagated and established in cells permissive for infection with herpesviruses (Sun et al., Nature genetics 8: 33-41, 1994; Cotter and Robertson. Curr Opin Mol Ther 5: 633-644, 1999). These large DNA viruses (herpes simplex virus (HSV) and Epstein-Barr virus (EBV), have the potential to deliver fragments of human heterologous DNA > 150 kb to specific cells. EBV recombinants can

maintain large pieces of DNA in the infected B-cells as episomal DNA. Individual clones carried human genomic inserts up to 330 kb appeared genetically stable. The maintenance of these episomes requires a specific EBV nuclear protein, EBNA1, constitutively expressed during infection with EBV. Additionally, these vectors can be used for transfection, where large amounts of protein can be generated transiently in vitro. Herpesvirus amplicon systems are also being used to package pieces of DNA > 220 kb and to infect cells that can stably maintain DNA as episomes.

[115] Other useful systems include, for example, replicating and host-restricted non-replicating vaccinia virus vectors.

10 b) Non-nucleic acid based systems

[116] The disclosed compositions can be delivered to the target cells in a variety of ways. For example, the compositions can be delivered through electroporation, or through lipofection, or through calcium phosphate precipitation. The delivery mechanism chosen will depend in part on the type of cell targeted and whether the delivery is occurring for example in vivo or in vitro.

[117] Thus, the compositions can comprise, in addition to the disclosed constructs or vectors for example, lipids such as liposomes, such as cationic liposomes (e.g., DOTMA, DOPE, DC-cholesterol) or anionic liposomes. Liposomes can further comprise proteins to facilitate targeting a particular cell, if desired. Administration of a composition comprising a compound and a cationic liposome can be administered to the blood afferent to a target organ or inhaled into the respiratory tract to target cells of the respiratory tract. Regarding liposomes, see, e.g., Brigham et al. *Am. J. Resp. Cell. Mol. Biol.* 1:95-100 (1989); Felgner et al. *Proc. Natl. Acad. Sci USA* 84:7413-7417 (1987); U.S. Pat. No.4,897,355. Furthermore, the compound can be administered as a component of a microcapsule that can be targeted to specific cell types, such as macrophages, or where the diffusion of the compound or delivery of the compound from the microcapsule is designed for a specific rate or dosage.

[118] In the methods described above which include the administration and uptake of exogenous DNA into the cells of a subject (i.e., gene transduction or transfection), delivery of the compositions to cells can be via a variety of mechanisms. As one example, delivery can be via a liposome, using commercially available liposome preparations such as LIPOFECTIN, LIPOFECTAMINE (GIBCO-BRL, Inc., Gaithersburg, MD), SUPERFECT (Qiagen, Inc. Hilden, Germany) and TRANSFECTAM (Promega Biotec, Inc., Madison, WI), as well as other

liposomes developed according to procedures standard in the art. In addition, the nucleic acid or vector of this invention can be delivered *in vivo* by electroporation, the technology for which is available from Genetronics, Inc. (San Diego, CA) as well as by means of a SONOPORATION machine (ImaRx Pharmaceutical Corp., Tucson, AZ).

5 [119] The materials can be in solution, suspension (for example, incorporated into microparticles, liposomes, or cells). These can be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter, et al., Bioconjugate Chem., 2:447-451, (1991); Bagshawe, K.D., Br. J. Cancer, 60:275-281, (1989); Bagshawe, et al., Br.
10 J. Cancer, 58:700-703, (1988); Senter, et al., Bioconjugate Chem., 4:3-9, (1993); Battelli, et al., Cancer Immunol. Immunother., 35:421-425, (1992); Pietersz and McKenzie, Immunolog. Reviews, 129:57-80, (1992); and Roffler, et al., Biochem. Pharmacol, 42:2062-2065, (1991)). These techniques can be used for a variety of other specific cell types. Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to
15 colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells *in vivo*. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., Cancer Research, 49:6214-6220, (1989); and Litzinger and Huang, Biochimica et Biophysica Acta, 1104:179-187,
20 (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways serve a variety of functions, such as nutrient uptake, removal of
25 activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, DNA and Cell Biology
30 10:6, 399-409 (1991)).

[120] Nucleic acids that are delivered to cells which are to be integrated into the host cell genome, typically contain integration sequences. These sequences are often viral related

sequences, particularly when viral based systems are used. These viral integration systems can also be incorporated into nucleic acids which are to be delivered using a non-nucleic acid based system of deliver, such as a liposome, so that the nucleic acid contained in the delivery system can be come integrated into the host genome.

5 [121] Other general techniques for integration into the host genome include, for example, systems designed to promote homologous recombination with the host genome. These systems typically rely on sequence flanking the nucleic acid to be expressed that has enough homology with a target sequence within the host cell genome that recombination between the vector nucleic acid and the target nucleic acid takes place, causing the delivered nucleic acid to
10 be integrated into the host genome. These systems and the methods necessary to promote homologous recombination are known to those of skill in the art.

c) *In vivo/ex vivo*

[122] As described herein, the compositions can be administered in a pharmaceutically acceptable carrier and can be delivered to the subject's cells *in vivo* and/or *ex vivo* by a variety
15 of mechanisms well known in the art (e.g., uptake of naked DNA, liposome fusion, intramuscular injection of DNA via a gene gun, endocytosis and the like).

[123] If *ex vivo* methods are employed, cells or tissues can be removed and maintained outside the body according to standard protocols well known in the art. The compositions can be introduced into the cells via any gene transfer mechanism, such as, for example, calcium
20 phosphate mediated gene delivery, electroporation, microinjection or proteoliposomes. The transduced cells can then be infused (e.g., in a pharmaceutically acceptable carrier) or homotopically transplanted back into the subject per standard methods for the cell or tissue type. Standard methods are known for transplantation or infusion of various cells into a subject.

[124] If *in vivo* delivery methods are performed the methods can be designed to deliver
25 the nucleic acid constructs directly to a particular cell type, via any delivery mechanism, such as intra-peritoneal injection of a vector construct. In this type of delivery situation, the nucleic acid constructs can be delivered to any type of tissue, for example, brain or neural or muscle. The nucleic acid constructs can also be delivered such that they generally deliver the nucleic acid constructs to more than one type of cell. This type of delivery can be accomplished, by for
30 example, injecting the constructs intraperitoneally into the flank of the organism. (See Example 2 and figures 8-10). In certain delivery methods, the timing of the delivery is monitored. For

example, the nucleic acid constructs can be delivered at the perinatal stage of the recipient's life or at the adult stage.

[125] The various vectors delivering the opioid receptors, such as the m-opioid receptor can be delivered to differentiated cells. For example, cells that are quiescent can be targeted with the disclosed vectors in certain embodiments. For example, nerve cells, which are no longer dividing, or are dividing very slowly, can be transfected with the disclosed compositions in certain embodiments. The nucleic acids can be delivered peripherally in certain embodiments and can be delivered by injection at a site distal to the body of the cell. For example, pain may be initiated at a point in the foot of an organism, but the body of the nerve transmitting the pain signal will be located at or near the spinal cord. In certain embodiments, the compositions can be delivered at the foot, transfecting the distal part of the nerve, including the most distal part of the nerve. Transfection, can take place along the full length of the cell, however. In certain embodiments, the vectors are delivered by injection at a site distal to a nerve body, or, for example, at the point of the pain with regard to where the body of the nerve is located.

4. Expression systems

[126] The nucleic acids that are delivered to cells typically contain expression controlling systems. For example, the inserted genes in viral and retroviral systems usually contain promoters, and/or enhancers to help control the expression of the desired gene product. A promoter is generally a sequence or sequences of DNA that function when in a relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of RNA polymerase and transcription factors, and can contain upstream elements and response elements.

a) Promoters and Enhancers

[127] Preferred promoters controlling transcription from vectors in mammalian host cells can be obtained from various sources, for example, the genomes of viruses such as: polyoma, Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis-B virus and most preferably cytomegalovirus, or from heterologous mammalian promoters, e.g. beta actin promoter. The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication (Fiers et al., Nature, 273: 113 (1978)). The immediate early promoter of the human cytomegalovirus is conveniently

obtained as a HindIII E restriction fragment (Greenway, P.J. et al., Gene 18: 355-360 (1982)). Of course, promoters from the host cell or related species also are useful herein.

[128] Enhancer generally refers to a sequence of DNA that functions at no fixed distance from the transcription start site and can be either 5' (Laimins, L. et al., Proc. Natl. Acad. Sci. 78: 993 (1981)) or 3' (Lusky, M.L., et al., Mol. Cell Bio. 3: 1108 (1983)) to the transcription unit. Furthermore, enhancers can be within an intron (Banerji, J.L. et al., Cell 33: 729 (1983)) as well as within the coding sequence itself (Osborne, T.F., et al., Mol. Cell Bio. 4: 1293 (1984)). They are usually between 10 and 300 bp in length, and they function in cis. Enhancers function to increase transcription from nearby promoters. Enhancers also often contain response elements that mediate the regulation of transcription. Promoters can also contain response elements that mediate the regulation of transcription. Enhancers often determine the regulation of expression of a gene. While many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, -fetoprotein and insulin), typically one will use an enhancer from a eukaryotic cell virus for general expression. Preferred examples are the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

[129] The promoter and/or enhancer can be specifically activated either by light or specific chemical events which trigger their function. Systems can be regulated by reagents such as tetracycline and dexamethasone. There are also ways to enhance viral vector gene expression by exposure to irradiation, such as gamma irradiation, or alkylating chemotherapy drugs.

[130] In certain embodiments the promoter and/or enhancer region can act as a constitutive promoter and/or enhancer to maximize expression of the region of the transcription unit to be transcribed. In certain constructs the promoter and/or enhancer region be active in all eukaryotic cell types, even if it is only expressed in a particular type of cell at a particular time. A preferred promoter of this type is the CMV promoter (650 bases). Other preferred promoters are SV40 promoters, cytomegalovirus (full length promoter), and retroviral vector LTR.

[131] It has been shown that all specific regulatory elements can be cloned and used to construct expression vectors that are selectively expressed in specific cell types such as melanoma cells. The glial fibrillary acidic protein (GFAP) promoter has been used to selectively express genes in cells of glial origin.

[132] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) can also contain sequences necessary for the termination of transcription which can affect mRNA expression. These regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding tissue factor protein. The 3' untranslated regions also include transcription termination sites. It is preferred that the transcription unit also contains a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in expression constructs is well established. It is preferred that homologous polyadenylation signals be used in the transgene constructs. In certain transcription units, the polyadenylation region is derived from the SV40 early polyadenylation signal and consists of about 400 bases. It is also preferred that the transcribed units contain other standard sequences alone or in combination with the above sequences improve expression from, or stability of, the construct.

b) Constitutive promoters

[133] In certain embodiments the promoters are constitutive promoters. This can be any promoter that causes transcription regulation in the absence of the addition of other factors. Examples of this type of promoter are the CMV promoter and the beta actin promoter, as well as others discussed herein. In certain embodiments the promoter can consist of fusions of one or more different types of promoters. For example, the regulatory regions of the CMV promoter and the beta actin promoter are well known and understood, examples, of which are disclosed herein. Parts of these promoters can be fused together to, for example, produce a CMV-beta actin fusion promoter. It is understood that this type of promoter has a CMV component and a beta actin component. These components can function independently as promoters, and thus, are themselves considered beta actin promoters and CMV promoters. A promoter can be any portion of a known promoter that causes promoter activity. It is well understood that many promoters, including the CMV and Beta Actin promoters have functional domains which are understood and that these can be used as a beta actin promoter or CMV promoter. Furthermore, these domains can be determined. There are many CMV promoter variations that exist, as well as beta actin promoters, and fusion promoters. These promoters can be compared, and for example, functional regions delineated, as described herein. Furthermore, each of these sequences can function independently or together in any combination to provide a promoter region for the disclosed nucleic acids.

c) Non-constitutive promoters

[134] The promoters can also be non-constitutive promoters, such as cell specific promoters. These are promoters that are turned on at specific time in development or stage or a particular type of cell, such as a cardiac cell, or neural cell, or a bone cell. Some examples of cell specific promoters are, the neural enolase specific promoter, (NSE) the COL1A1 procollagen promoter, and the CD11b promoter (PBMC-microglia/macrophage/monocyte specific promoter.

[02] It is understood that the recombinant systems can be expressed in a tissue-specific manner. It is understood that tissue specific expression can occur due to the presence of a tissue-specific promoter. Typically, proteins under control of a tissue-specific promoter are transcribed when the promoter becomes active by virtue of being present in the tissue for which it is specific. Therefore, all cells can encode for a particular gene without global expression. As such, labeled proteins can be shown to be present in certain tissues without expression in other nearby tissues that may complicate results or expression of proteins in tissues where expression may be detrimental to the host. Disclosed are methods wherein the cre recombinase is under the control of the EIIA promoter, a promoter specific for breast tissue, such as the WAP promoter, a promoter specific for ovarian tissue, such as the ACTB promoter, or a promoter specific for bone tissue, such as osteocalcin. Any tissues specific promoter can be used. Promoters specific for prostate, testis, and neural are also disclosed. Examples of some tissue-specific promoters include but are not limited to MUC1, EIIA, ACTB, WAP, bHLH-EC2, HOXA-1, Alpha-fetoprotein (AFP), opsin, CR1/2, Fc- γ -Receptor 1 (Fc- γ -R1), MMTVD-LTR, the human insulin promoter, Pdha-2, rat neuron-specific enolase. For example, use of the AFP promoter creates specificity for the liver. Another example, HOXA-1 is a neuronal tissue specific promoter, and as such, proteins expressed under the control of HOXA-1 are only expressed in neuronal tissue. (All of which are herein incorporated by reference at least for the sequence of the promoters and related sequences.)

[135] Other cell specific promoters can be found in (Sutcliffe, J.G. (1988), *Ann. Rev. Neuroscience* 11, 157-198). For example, when transfecting nerve cells, there are a variety of nerve specific promoters, such as the neuron specific enolase promoter. Other examples of neuron specific promoters would be the Tau promoter, Synapsin I (Hoesche, C., Sauerwald, A., et al., (1993) *J. Biol. Chem.* 268, 26494-26502. and II (Chin, L.-S et al., (1994), *J. Biol. Chem.* 269, 18507-18513) promoters, the amino acid decarboxylase (AADC) (Albert, V., et al., (1992),

Proc. Natl. Acad. Sci. 89, 12053-12057) and FE65 (Faraonio, R., et al., (1994), *Nucl. Acids Res.* 22, 4876-4883) promoters. Other nerve specific promoters include, the promoter for the WT1 gene (Fraizer, G, et al., (1994), *J. Biol. Chem.* 269, 8892-8900), neurofilament light chain promoter (Yazdanbakhsh, K., et al., (1993) *Nucl. Acids Res.* 21, 455-461), and the glial fibrillary acidic protein, (Kaneko, R. & Sueoka, N. (1993) *Proc. Natl. Acad. Sci.* 90, 4698-4702). (All of which are herein incorporated by reference at least for the sequence of the promoters and related sequences.)

[136] Expression of the transgene can be targeted selectively to neurons by cloning a neuron specific promoter, such as the NSE promoter as disclosed herein (Liu H. et al., *Journal of Neuroscience*. 23(18):7143-54, 2003); tyrosine hydroxylase promoter (Kessler MA. et al., *Brain Research. Molecular Brain Research*. 112(1-2):8-23, 2003); myelin basic protein promoter (Kessler MA. et al *Biochemical & Biophysical Research Communications*. 288(4):809-18, 2001); glial fibrillary acidic protein promoter (Nolte C. et al., *GLIA*. 33(1):72-86, 2001); neurofilaments gene (heavy, medium, light) promoters (Yaworsky PJ. et al., *Journal of Biological Chemistry*. 272(40):25112-20, 1997) (All of which are herein incorporated by reference at least for the sequence of the promoters and related sequences.) The NSE promoter is disclosed in Peel AL. et al., *Gene Therapy*. 4(1):16-24, 1997) (SEQ ID NO:69) (pTR-NT3myc; Powell Gene Therapy Center, University of Florida, Gainesville FL).

d) Markers

[137] The viral vectors can include nucleic acid sequence encoding a marker product. This marker product is used to determine if the gene has been delivered to the cell and once delivered is being expressed. Preferred marker genes are the *E. Coli* lacZ gene, which encodes β -galactosidase, and green fluorescent protein.

[138] In some embodiments the marker can be a selectable marker. Examples of suitable selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase, neomycin, neomycin analog G418, hydromycin, and puromycin. When such selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell can survive if placed under selective pressure. There are two widely used distinct categories of selective regimes. The first category is based on a cell's metabolism and the use of a mutant cell line which lacks the ability to grow independent of a supplemented media. Two examples are: CHO DHFR- cells and mouse LTK- cells. These cells lack the ability to grow without the addition of such nutrients as thymidine or hypoxanthine. Because these cells lack

certain genes necessary for a complete nucleotide synthesis pathway, they cannot survive unless the missing nucleotides are provided in a supplemented media. An alternative to supplementing the media is to introduce an intact DHFR or TK gene into cells lacking the respective genes, thus altering their growth requirements. Individual cells which were not transformed with the DHFR or TK gene will not be capable of survival in non-supplemented media.

[139] The second category is dominant selection which refers to a selection scheme used in any cell type and does not require the use of a mutant cell line. These schemes typically use a drug to arrest growth of a host cell. Those cells which have a novel gene would express a protein conveying drug resistance and would survive the selection. Examples of such dominant selection use the drugs neomycin, (Southern P. and Berg, P., *J. Molec. Appl. Genet.* 1: 327 (1982)), mycophenolic acid, (Mulligan, R.C. and Berg, P. *Science* 209: 1422 (1980)) or hygromycin, (Sugden, B. et al., *Mol. Cell. Biol.* 5: 410-413 (1985)). The three examples employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid) or hygromycin, respectively. Others include the neomycin analog G418 and puramycin.

e) Post transcriptional regulatory elements

[140] The disclosed vectors can also contain post-transcriptional regulatory elements. Post-transcriptional regulatory elements can enhance mRNA stability or enhance translation of the transcribed mRNA. An exemplary post-transcriptional regulatory sequence is the WPRE sequence isolated from the woodchuck hepatitis virus. (Zufferey R, et al., "Woodchuck hepatitis virus post-transcriptional regulatory element enhances expression of transgenes delivered by retroviral vectors," *J Virol*; 73:2886-92 (1999)). Post-transcriptional regulatory elements can be positioned both 3' and 5' to the exogenous gene, but it is preferred that they are positioned 3' to the exogenous gene.

f) Transduction efficiency elements

[141] Transduction efficiency elements are sequences that enhance the packaging and transduction of the vector. These elements typically contain polypurine sequences. An example of a transduction efficiency element is the ppt-cts sequence that contains the central polypurine tract (ppt) and central terminal site (cts) from the FIV. These sequences are in the disclosed FIV sequences herein. Each retrovirus and lentivirus can have there own ppt-cts.

g) 3' untranslated regions

[142] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) can also contain sequences necessary for the termination of transcription which can affect mRNA expression. These 3' untranslated regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding the exogenous
5 gene. The 3' untranslated regions also include transcription termination sites. The transcription unit also can contain a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in expression constructs is well established. Homologous polyadenylation signals can be used in the transgene constructs. In an embodiment
10 of the transcription unit, the polyadenylation region is derived from the SV40 early polyadenylation signal and consists of about 400 bases. Transcribed units can contain other standard sequences alone or in combination with the above sequences improve expression from, or stability of, the construct.

5. Sequence similarities

[143] It is understood that as discussed herein the use of the terms homology and
15 identity mean the same thing as similarity. Thus, for example, if the use of the word homology is used between two non-natural sequences it is understood that this is not necessarily indicating an evolutionary relationship between these two sequences, but rather is looking at the similarity or relatedness between their nucleic acid sequences. Many of the methods for determining
20 homology between two evolutionarily related molecules are routinely applied to any two or more nucleic acids or proteins for the purpose of measuring sequence similarity regardless of whether they are evolutionarily related or not.

[144] In general, it is understood that one way to define any known variants and
derivatives or those that might arise, of the disclosed genes and proteins herein, is through
25 defining the variants and derivatives in terms of homology to specific known sequences. This identity of particular sequences disclosed herein is also discussed elsewhere herein. In general, variants of genes and proteins herein disclosed typically have at least, about 70, 71, 72, 73, 74,
75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99
percent homology to the stated sequence or the native sequence. Those of skill in the art readily
30 understand how to determine the homology of two proteins or nucleic acids, such as genes. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

[145] Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol Biol.* 48: 443 (1970), by the search for similarity
5 method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by inspection.

[146] The same types of homology can be obtained for nucleic acids by for example
10 the algorithms disclosed in Zuker, M. *Science* 244:48-52, 1989, Jaeger et al. *Proc. Natl. Acad. Sci. USA* 86:7706-7710, 1989, Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 which are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in certain instances the results of these various methods can differ, but the skilled artisan understands if identity is found with
15 at least one of these methods, the sequences would be said to have the stated identity, and be disclosed herein.

[147] For example, as used herein, a sequence recited as having a particular percent homology to another sequence refers to sequences that have the recited homology as calculated by any one or more of the calculation methods described above. For example, a first sequence
20 has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using the Zuker calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by any of the other calculation methods. As another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to
25 have 80 percent homology to the second sequence using both the Zuker calculation method and the Pearson and Lipman calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by the Smith and Waterman calculation method, the Needleman and Wunsch calculation method, the Jaeger calculation methods, or any of the other calculation methods. As yet another example, a first sequence has 80 percent homology,
30 as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using each of calculation methods (although, in practice, the different calculation methods will often result in different calculated homology percentages).

6. Hybridization/selective hybridization

[148] The term hybridization typically means a sequence driven interaction between at least two nucleic acid molecules, such as a primer or a probe and a gene. Sequence driven interaction means an interaction that occurs between two nucleotides or nucleotide analogs or nucleotide derivatives in a nucleotide specific manner. For example, G interacting with C or A interacting with T are sequence driven interactions. Typically sequence driven interactions occur on the Watson-Crick face or Hoogsteen face of the nucleotide. The hybridization of two nucleic acids is affected by a number of conditions and parameters known to those of skill in the art. For example, the salt concentrations, pH, and temperature of the reaction all affect whether two nucleic acid molecules will hybridize.

[149] Parameters for selective hybridization between two nucleic acid molecules are well known to those of skill in the art. For example, in some embodiments selective hybridization conditions can be defined as stringent hybridization conditions. For example, stringency of hybridization is controlled by both temperature and salt concentration of either or both of the hybridization and washing steps. For example, the conditions of hybridization to achieve selective hybridization can involve hybridization in high ionic strength solution (6X SSC or 6X SSPE) at a temperature that is about 12-25°C below the T_m (the melting temperature at which half of the molecules dissociate from their hybridization partners) followed by washing at a combination of temperature and salt concentration chosen so that the washing temperature is about 5°C to 20°C below the T_m. The temperature and salt conditions are readily determined empirically in preliminary experiments in which samples of reference DNA immobilized on filters are hybridized to a labeled nucleic acid of interest and then washed under conditions of different stringencies. Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA hybridizations. The conditions can be used as described above to achieve stringency, or as is known in the art. (Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. *Methods Enzymol.* 1987:154:367, 1987 which is herein incorporated by reference for material at least related to hybridization of nucleic acids). A preferable stringent hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization and washing, if desired, can be reduced accordingly as the degree of complementarity desired is decreased, and further, depending upon the G-C or A-T richness of any area wherein variability

is searched for. Likewise, stringency of hybridization and washing, if desired, can be increased accordingly as homology desired is increased, and further, depending upon the G-C or A-T richness of any area wherein high homology is desired, all as known in the art.

[150] Another way to define selective hybridization is by looking at the amount
5 (percentage) of one of the nucleic acids bound to the other nucleic acid. For example, in some embodiments selective hybridization conditions would be when at least about, 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 percent of the limiting nucleic acid is bound to the non-limiting nucleic acid. Typically, the non-limiting primer is in for example, 10 or 100 or 1000 fold excess. This type of
10 assay can be performed at under conditions where both the limiting and non-limiting primer are for example, 10 fold or 100 fold or 1000 fold below their k_d , or where only one of the nucleic acid molecules is 10 fold or 100 fold or 1000 fold or where one or both nucleic acid molecules are above their k_d .

[151] Another way to define selective hybridization is by looking at the percentage of
15 primer that gets enzymatically manipulated under conditions where hybridization is required to promote the desired enzymatic manipulation. For example, in some embodiments selective hybridization conditions would be when at least about, 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 percent of the primer is enzymatically manipulated under conditions which promote the enzymatic
20 manipulation, for example if the enzymatic manipulation is DNA extension, then selective hybridization conditions would be when at least about 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 percent of the primer molecules are extended. Preferred conditions also include those suggested by the manufacturer or indicated in the art as being appropriate for the enzyme performing the
25 manipulation.

[152] Just as with homology, it is understood that there are a variety of methods herein disclosed for determining the level of hybridization between two nucleic acid molecules. It is understood that these methods and conditions can provide different percentages of hybridization between two nucleic acid molecules, but unless otherwise indicated meeting the parameters of
30 any of the methods would be sufficient. For example if 80% hybridization was required and as long as hybridization occurs within the required parameters in any one of these methods it is considered disclosed herein.

[153] It is understood that those of skill in the art understand that if a composition or method meets any one of these criteria for determining hybridization either collectively or singly it is a composition or method that is disclosed herein.

7. Nucleic acids

5 [154] There are a variety of molecules disclosed herein that are nucleic acid based, including for example the nucleic acids that encode, for example μ -opioid receptor, or functional nucleic acids. The disclosed nucleic acids can be made up of for example, nucleotides, nucleotide analogs, or nucleotide substitutes. Non-limiting examples of these and other molecules are discussed herein. It is understood that for example, when a vector is
10 expressed in a cell, that the expressed mRNA will typically be made up of A, C, G, and U. Likewise, it is understood that if, for example, an antisense molecule is introduced into a cell or cell environment through for example exogenous delivery, it is advantageous that the antisense molecule be made up of nucleotide analogs that reduce the degradation of the antisense molecule in the cellular environment.

15 [155] A nucleotide is a molecule that contains a base moiety, a sugar moiety and a phosphate moiety. Nucleotides can be linked together through their phosphate moieties and sugar moieties creating an internucleoside linkage. The base moiety of a nucleotide can be adenin-9-yl (A), cytosin-1-yl (C), guanin-9-yl (G), uracil-1-yl (U), and thymin-1-yl (T). The sugar moiety of a nucleotide is a ribose or a deoxyribose. The phosphate moiety of a nucleotide
20 is pentavalent phosphate. A non-limiting example of a nucleotide would be 3'-AMP (3'-adenosine monophosphate) or 5'-GMP (5'-guanosine monophosphate).

[156] A nucleotide analog is a nucleotide which contains some type of modification to either the base, sugar, or phosphate moieties. Modifications to nucleotides are well known in the art and would include for example, 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine,
25 xanthine, hypoxanthine, and 2-aminoadenine as well as modifications at the sugar or phosphate moieties.

[157] Nucleotide substitutes are molecules having similar functional properties to nucleotides, but which do not contain a phosphate moiety, such as peptide nucleic acid (PNA). Nucleotide substitutes are molecules that will recognize nucleic acids in a Watson-Crick or
30 Hoogsteen manner, but which are linked together through a moiety other than a phosphate

moiety. Nucleotide substitutes are able to conform to a double helix type structure when interacting with the appropriate target nucleic acid.

[158] It is also possible to link other types of molecules (conjugates) to nucleotides or nucleotide analogs to enhance for example, cellular uptake. Conjugates can be chemically
5 linked to the nucleotide or nucleotide analogs. Such conjugates include but are not limited to lipid moieties such as a cholesterol moiety. (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989,86, 6553-6556),

[159] A Watson-Crick interaction is at least one interaction with the Watson-Crick face of a nucleotide, nucleotide analog, or nucleotide substitute. The Watson-Crick face of a
10 nucleotide, nucleotide analog, or nucleotide substitute includes the C2, N1, and C6 positions of a purine based nucleotide, nucleotide analog, or nucleotide substitute and the C2, N3, C4 positions of a pyrimidine based nucleotide, nucleotide analog, or nucleotide substitute.

[160] A Hoogsteen interaction is the interaction that takes place on the Hoogsteen face of a nucleotide or nucleotide analog, which is exposed in the major groove of duplex DNA. The
15 Hoogsteen face includes the N7 position and reactive groups (NH₂ or O) at the C6 position of purine nucleotides.

a) Sequences

[161] There are a variety of sequences related to μ -opioid receptor and promoter sequences. These sequences and others are herein incorporated by reference in their entireties as
20 well as for individual subsequences contained therein. It is understood that there are numerous Genbank accession sequences related to μ -opioid receptor, all of which are incorporated by reference herein.

[162] One particular sequence set forth in SEQ ID NO:2, which is a sequence for human μ -opioid receptor cDNA, is used herein, as an example, to exemplify the disclosed
25 compositions and methods. It is understood that the description related to this sequence is applicable to any sequence related to μ -opioid receptor unless specifically indicated otherwise. Those of skill in the art understand how to resolve sequence discrepancies and differences and to adjust the compositions and methods relating to a particular sequence to other related sequences. Primers and/or probes can be designed for any of the sequences disclosed herein
30 given the information disclosed herein and that known in the art.

[163] It is also understood for example that there are numerous vectors that can be used to create the μ -opioid receptor construct nucleic acids.

b) Primers and probes

[164] Disclosed are compositions including primers and probes, which are capable of interacting with, for example, the μ -opioid receptor construct nucleic acids, as disclosed herein. In certain embodiments the primers are used to support DNA amplification reactions. Typically the primers will be capable of being extended in a sequence specific manner. Extension of a primer in a sequence specific manner includes any methods wherein the sequence and/or composition of the nucleic acid molecule to which the primer is hybridized or otherwise associated directs or influences the composition or sequence of the product produced by the extension of the primer. Extension of the primer in a sequence specific manner therefore includes, but is not limited to, PCR, DNA sequencing, DNA extension, DNA polymerization, RNA transcription, or reverse transcription. Techniques and conditions that amplify the primer in a sequence specific manner are preferred. In certain embodiments the primers are used for the DNA amplification reactions, such as PCR or direct sequencing. It is understood that in certain embodiments the primers can also be extended using non-enzymatic techniques, where for example, the nucleotides or oligonucleotides used to extend the primer are modified such that they will chemically react to extend the primer in a sequence specific manner. Typically the disclosed primers hybridize with, for example, the μ -opioid receptor construct nucleic acid, or region of the μ -opioid receptor construct nucleic acids or they hybridize with the complement of the μ -opioid receptor construct nucleic acids or complement of a region of the μ -opioid receptor construct nucleic acids.

8. Peptides

a) Protein variants

[165] As discussed herein there are numerous variants of the μ -opioid receptor protein that are known and herein contemplated. In addition, to the known functional species and allelic variants of μ -opioid receptor there are derivatives of the μ -opioid receptor protein which also function in the disclosed methods and compositions. Protein variants and derivatives are well understood to those of skill in the art and in can involve amino acid sequence modifications. For example, amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional or deletional variants. Insertions include amino

and/or carboxyl terminal fusions as well as intrasequence insertions of single or multiple amino acid residues. Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Immunogenic fusion protein derivatives, such as those described in the examples, are made by fusing a polypeptide

5 sufficiently large to confer immunogenicity to the target sequence by cross-linking in vitro or by recombinant cell culture transformed with DNA encoding the fusion. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site within the protein molecule. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in

10 the DNA encoding the protein, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example M13 primer mutagenesis and PCR mutagenesis. Amino acid substitutions are typically of single residues, but can occur at a number of different locations at once; insertions usually will be on

15 the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs, i.e. a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof can be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could

20 produce secondary mRNA structure. Substitutional variants are those in which at least one residue has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Tables 4 and 5 and are referred to as conservative substitutions.

[166] TABLE 4: Amino Acid Abbreviations

Amino Acid	Abbreviations
alanine	AlaA
allosoleucine	Aile
arginine	ArgR
asparagine	AsnN
aspartic acid	AspD
cysteine	CysC
glutamic acid	GluE
glutamine	GlnK
glycine	GlyG
histidine	HisH
isoleucine	IleI
leucine	LeuL

Amino Acid	Abbreviations
lysine	LysK
phenylalanine	PheF
proline	ProP
pyroglutamic acidp	Glu
serine	SerS
threonine	ThrT
tyrosine	TyrY
tryptophan	TrpW
valine	ValV

TABLE 5: Amino Acid Substitutions	
Original Residue Exemplary Conservative Substitutions, others are known in the art.	
Ala	ser
Arg	lys, gln
Asn	gln; his
Asp	glu
Cys	ser
Gln	asn, lys
Glu	asp
Gly	pro
His	asn; gln
Ile	leu; val
Leu	ile; val
Lys	arg; gln;
Met	Leu; ile
Phe	met; leu; tyr
Ser	thr
Thr	ser
Trp	tyr
Tyr	trp; phe
Val	ile; leu

[167] Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those in Table 2, i.e., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the protein properties will be those in which (a) a hydrophilic residue, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an

electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine, in this case, (e) by increasing the number of sites for sulfation and/or glycosylation.

5 [168] For example, the replacement of one amino acid residue with another that is biologically and/or chemically similar is known to those skilled in the art as a conservative substitution. For example, a conservative substitution would be replacing one hydrophobic residue for another, or one polar residue for another. The substitutions include combinations such as, for example, Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe,
10 Tyr. Such conservatively substituted variations of each explicitly disclosed sequence are included within the mosaic polypeptides provided herein.

 [169] Substitutional or deletional mutagenesis can be employed to insert sites for N-glycosylation (Asn-X-Thr/Ser) or O-glycosylation (Ser or Thr). Deletions of cysteine or other labile residues also may be desirable. Deletions or substitutions of potential proteolysis sites,
15 e.g. Arg, is accomplished for example by deleting one of the basic residues or substituting one by glutamyl or histidyl residues.

 [170] Certain post-translational derivatizations are the result of the action of recombinant host cells on the expressed polypeptide. Glutamyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues.
20 Alternatively, these residues are deamidated under mildly acidic conditions. Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the o-amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco pp 79-86 (1983)), acetylation of the N-
25 terminal amine and, in some instances, amidation of the C-terminal carboxyl.

 [171] It is understood that one way to define the variants and derivatives of the disclosed proteins herein is through defining the variants and derivatives in terms of homology/identity to specific known sequences. For example, SEQ ID NO:1 sets forth a particular sequence of μ -opioid receptor. Specifically disclosed are variants of these and other
30 proteins herein disclosed which have at least, 70% or 75% or 80% or 85% or 90% or 95% homology to the stated sequence. Those of skill in the art readily understand how to determine

the homology of two proteins. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

[172] Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison can be conducted by the local homology
5 algorithm of Smith and Waterman *Adv. Appl. Math.* 2: 482 (1981), by the homology
alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search
for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85: 2444 (1988),
by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA
in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr.,
10 Madison, WI), or by inspection.

[173] The same types of homology can be obtained for nucleic acids by for example
the algorithms disclosed in Zuker, M. *Science* 244:48-52, 1989, Jaeger et al. *Proc. Natl. Acad.
Sci. USA* 86:7706-7710, 1989, Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 which are
herein incorporated by reference for at least material related to nucleic acid alignment.

15 [174] It is understood that the description of conservative mutations and homology can
be combined together in any combination, such as embodiments that have at least 70%
homology to a particular sequence wherein the variants are conservative mutations.

[175] As this specification discusses various proteins and protein sequences it is
understood that the nucleic acids that can encode those protein sequences are also disclosed.
20 This would include all degenerate sequences related to a specific protein sequence, i.e. all
nucleic acids having a sequence that encodes one particular protein sequence as well as all
nucleic acids, including degenerate nucleic acids, encoding the disclosed variants and
derivatives of the protein sequences. Thus, while each particular nucleic acid sequence may not
be written out herein, it is understood that each and every sequence is in fact disclosed and
25 described herein through the disclosed protein sequence. For example, one of the many nucleic
acid sequences that can encode the protein sequence set forth in SEQ ID NO:3 is set forth in
SEQ ID NO:4. Another nucleic acid sequence that encodes the same protein sequence set forth
in SEQ ID NO:3 is set forth in SEQ ID NO:8. In addition, for example, a disclosed
conservative derivative of SEQ ID NO:3 is shown in SEQ ID NO: 9, where the valine (V) at
30 position 21 is changed to an isoleucine (I). It is understood that for this mutation, all of the
nucleic acid sequences that encode this particular derivative of the SEQ ID NO:3 polypeptide
are also disclosed. It is also understood that while no amino acid sequence indicates what

particular DNA sequence encodes that protein within an organism, where particular variants of a disclosed protein are disclosed herein, the known nucleic acid sequence that encodes that protein in the particular organism from which that protein arises is also known and herein disclosed and described.

5 [176] It is understood that there are numerous amino acid and peptide analogs which can be incorporated into the disclosed compositions. For example, there are numerous D amino acids or amino acids which have a different functional substituent than the amino acids shown in Table 1 and Table 2. The opposite stereo isomers of naturally occurring peptides are disclosed, as well as the stereo isomers of peptide analogs. These amino acids can readily be
10 incorporated into polypeptide chains by charging tRNA molecules with the amino acid of choice and engineering genetic constructs that utilize, for example, amber codons, to insert the analog amino acid into a peptide chain in a site specific way (Thorson et al., *Methods in Molec. Biol.* 77:43-73 (1991), Zoller, *Current Opinion in Biotechnology*, 3:348-354 (1992); Ibba, *Biotechnology & Genetic Engineering Reviews* 13:197-216 (1995), Cahill et al., *TIBS*,
15 14(10):400-403 (1989); Benner, *TIB Tech*, 12:158-163 (1994); Ibba and Hennecke, *Bio/technology*, 12:678-682 (1994) all of which are herein incorporated by reference at least for material related to amino acid analogs).

[177] Molecules can be produced that resemble peptides, but which are not connected via a natural peptide linkage. For example, linkages for amino acids or amino acid analogs can
20 include $\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{S}-$, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$ (cis and trans), $-\text{COCH}_2-$, $-\text{CH}(\text{OH})\text{CH}_2-$, and $-\text{CHH}_2\text{SO}-$ (These and others can be found in Spatola, A. F. in *Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins*, B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983); Spatola, A. F., *Vega Data* (March 1983), Vol. 1, Issue 3, *Peptide Backbone Modifications* (general review); Morley, *Trends Pharm Sci* (1980) pp. 463-468;
25 Hudson, D. et al., *Int J Pept Prot Res* 14:177-185 (1979) ($-\text{CH}_2\text{NH}-$, CH_2CH_2-); Spatola et al. *Life Sci* 38:1243-1249 (1986) ($-\text{CH H}_2-\text{S}$); Hann J. *Chem. Soc Perkin Trans. I* 307-314 (1982) ($-\text{CH}-\text{CH}-$, cis and trans); Almquist et al. *J. Med. Chem.* 23:1392-1398 (1980) ($-\text{COCH}_2-$); Jennings-White et al. *Tetrahedron Lett* 23:2533 (1982) ($-\text{COCH}_2-$); Szelke et al. *European Appln*, EP 45665 CA (1982): 97:39405 (1982) ($-\text{CH}(\text{OH})\text{CH}_2-$); Holladay et al. *Tetrahedron Lett* 24:4401-4404 (1983) ($-\text{C}(\text{OH})\text{CH}_2-$); and Hruby *Life Sci* 31:189-199 (1982) ($-\text{CH}_2-\text{S}-$);
30 each of which is incorporated herein by reference. A particularly preferred non-peptide linkage is $-\text{CH}_2\text{NH}-$. It is understood that peptide analogs can have more than one atom between the bond atoms, such as β -alanine, γ -aminobutyric acid, and the like.

[178] Amino acid analogs and analogs and peptide analogs often have enhanced or desirable properties, such as, more economical production, greater chemical stability, enhanced pharmacological properties (half-life, absorption, potency, efficacy, etc.), altered specificity (e.g., a broad-spectrum of biological activities), reduced antigenicity, and others.

5 [179] D-amino acids can be used to generate more stable peptides, because D amino acids are not recognized by peptidases and such. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) can be used to generate more stable peptides. Cysteine residues can be used to cyclize or attach two or more peptides together. This can be beneficial to constrain peptides into
10 particular conformations. (Rizo and Gierasch Ann. Rev. Biochem. 61:387 (1992), incorporated herein by reference).

[180]

9. Pharmaceutical carriers/Delivery of pharmaceutical products

[181] As described above, the compositions can also be administered *in vivo* in a
15 pharmaceutically acceptable carrier. By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material can be administered to a subject, along with the nucleic acid or vector, without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. The carrier would naturally be selected to minimize any
20 degradation of the active ingredient and to minimize any adverse side effects in the subject, as would be well known to one of skill in the art.

[182] The compositions can be administered orally, parenterally (e.g., intravenously), by intramuscular injection, by intraperitoneal injection, transdermally, extracorporeally, topically or the like, including topical intranasal administration or administration by inhalant.
25 As used herein, "topical intranasal administration" means delivery of the compositions into the nose and nasal passages through one or both of the nares and can comprise delivery by a spraying mechanism or droplet mechanism, or through aerosolization of the nucleic acid or vector. Administration of the compositions by inhalant can be through the nose or mouth via delivery by a spraying or droplet mechanism. Delivery can also be directly to any area of the
30 respiratory system (e.g., lungs) via intubation. The exact amount of the compositions required will vary from subject to subject, depending on the species, age, weight and general condition of the subject, the severity of the allergic disorder being treated, the particular nucleic acid or

vector used, its mode of administration and the like. Thus, it is not possible to specify an exact amount for every composition. However, an appropriate amount can be determined by one of ordinary skill in the art using only routine experimentation given the teachings herein.

[183] Parenteral administration of the composition, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution of suspension in liquid prior to injection, or as emulsions. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system such that a constant dosage is maintained. See, e.g., U.S. Patent No. 3,610,795, which is incorporated by reference herein.

[184] The materials can be in solution, suspension (for example, incorporated into microparticles, liposomes, or cells). These can be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter, et al., *Bioconjugate Chem.*, 2:447-451, (1991); Bagshawe, K.D., *Br. J. Cancer*, 60:275-281, (1989); Bagshawe, et al., *Br. J. Cancer*, 58:700-703, (1988); Senter, et al., *Bioconjugate Chem.*, 4:3-9, (1993); Battelli, et al., *Cancer Immunol. Immunother.*, 35:421-425, (1992); Pietersz and McKenzie, *Immunolog. Reviews*, 129:57-80, (1992); and Roffler, et al., *Biochem. Pharmacol.*, 42:2062-2065, (1991)). Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells *in vivo*. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., *Cancer Research*, 49:6214-6220, (1989); and Litzinger and Huang, *Biochimica et Biophysica Acta*, 1104:179-187, (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of

receptor-mediated endocytosis has been reviewed (Brown and Greene, *DNA and Cell Biology* 10:6, 399-409 (1991)).

a) **Pharmaceutically Acceptable Carriers**

[185] The compositions, including antibodies, can be used therapeutically in
5 combination with a pharmaceutically acceptable carrier.

[186] Suitable carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy* (19th ed.) ed. A.R. Gennaro, Mack Publishing Company, Easton, PA 1995. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to render the formulation isotonic. Examples of the pharmaceutically-acceptable
10 carrier include, but are not limited to, saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more preferably from about 7 to about 7.5. Further carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, liposomes or microparticles. It will be apparent to those persons skilled in
15 the art that certain carriers can be more preferable depending upon, for instance, the route of administration and concentration of composition being administered.

[187] Pharmaceutical carriers are known to those skilled in the art. These most typically would be standard carriers for administration of drugs to humans, including solutions such as sterile water, saline, and buffered solutions at physiological pH. The compositions can
20 be administered intramuscularly or subcutaneously. Other compounds will be administered according to standard procedures used by those skilled in the art.

[188] Pharmaceutical compositions can include carriers, thickeners, diluents, buffers, preservatives, surface active agents and the like in addition to the molecule of choice. Pharmaceutical compositions can also include one or more active ingredients such as antimicrobial
25 agents, antiinflammatory agents, anesthetics, and the like.

[189] The pharmaceutical composition can be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Administration can be topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip, subcutaneous, intraperitoneal or
30 intramuscular injection. The disclosed antibodies can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally.

[190] Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives can also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

[191] Formulations for topical administration can include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

[192] Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be desirable.

[193] Some of the compositions can potentially be administered as a pharmaceutically acceptable acid- or base- addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

10. Chips and micro arrays

[194] Disclosed are chips where at least one address is the sequences or part of the sequences set forth in any of the nucleic acid sequences disclosed herein. Also disclosed are chips where at least one address is the sequences or portion of sequences set forth in any of the peptide sequences disclosed herein.

[195] Also disclosed are chips where at least one address is a variant of the sequences or part of the sequences set forth in any of the nucleic acid sequences disclosed herein. Also

disclosed are chips where at least one address is a variant of the sequences or portion of sequences set forth in any of the peptide sequences disclosed herein.

11. Computer readable mediums

[196] It is understood that the disclosed nucleic acids and proteins can be represented as a sequence consisting of the nucleotides of amino acids. There are a variety of ways to display these sequences, for example the nucleotide guanosine can be represented by G or g. Likewise the amino acid valine can be represented by Val or V. Those of skill in the art understand how to display and express any nucleic acid or protein sequence in any of the variety of ways that exist, each of which is considered herein disclosed. Specifically contemplated herein is the display of these sequences on computer readable mediums, such as, commercially available floppy disks, tapes, chips, hard drives, compact disks, and video disks, or other computer readable mediums. Also disclosed are the binary code representations of the disclosed sequences. Those of skill in the art understand what computer readable mediums. Thus, computer readable mediums on which the nucleic acids or protein sequences are recorded, stored, or saved.

[197] Disclosed are computer readable mediums comprising the sequences and information regarding the sequences set forth herein.

12. Kits

[198] Disclosed herein are kits that are drawn to reagents that can be used in practicing the methods disclosed herein. The kits can include any reagent or combination of reagent discussed herein or that would be understood to be required or beneficial in the practice of the disclosed methods. For example, the kits could include primers to perform the amplification reactions discussed in certain embodiments of the methods, as well as the buffers and enzymes required to use the primers as intended.

25 D. Methods of making the compositions

[199] The compositions disclosed herein and the compositions necessary to perform the disclosed methods can be made using any method known to those of skill in the art for that particular reagent or compound unless otherwise specifically noted.

[200] The disclosed viral vectors can be made using standard recombinant molecular biology techniques. Many of these techniques are illustrated in Maniatis (Maniatis et al., *"Molecular Cloning--A Laboratory Manual,"* (Cold Spring Harbor Laboratory, Latest edition)

and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989.

1. Nucleic acid synthesis

[201] For example, the nucleic acids, such as, the oligonucleotides to be used as
5 primers can be made using standard chemical synthesis methods or can be produced using enzymatic methods or any other known method. Such methods can range from standard enzymatic digestion followed by nucleotide fragment isolation (see for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) Chapters 5, 6) to purely synthetic methods, for example,
10 by the cyanoethyl phosphoramidite method using a Milligen or Beckman System Plus DNA synthesizer (for example, Model 8700 automated synthesizer of Milligen-Bioscience, Burlington, MA or ABI Model 380B). Synthetic methods useful for making oligonucleotides are also described by Ikuta et al., *Ann. Rev. Biochem.* 53:323-356 (1984), (phosphotriester and phosphite-triester methods), and Narang et al., *Methods Enzymol.*, 65:610-620 (1980),
15 (phosphotriester method). Protein nucleic acid molecules can be made using known methods such as those described by Nielsen et al., *Bioconjug. Chem.* 5:3-7 (1994).

2. Peptide synthesis

[202] One method of producing the disclosed proteins is to link two or more peptides or polypeptides together by protein chemistry techniques. For example, peptides or
20 polypeptides can be chemically synthesized using currently available laboratory equipment using either Fmoc (9-fluorenylmethyloxycarbonyl) or Boc (*tert*-butyloxycarbonyl) chemistry. (Applied Biosystems, Inc., Foster City, CA). One skilled in the art can readily appreciate that a peptide or polypeptide corresponding to the disclosed proteins, for example, can be synthesized by standard chemical reactions. For example, a peptide or polypeptide can be synthesized and
25 not cleaved from its synthesis resin whereas the other fragment of a peptide or protein can be synthesized and subsequently cleaved from the resin, thereby exposing a terminal group which is functionally blocked on the other fragment. By peptide condensation reactions, these two fragments can be covalently joined via a peptide bond at their carboxyl and amino termini, respectively, to form an antibody, or fragment thereof. (Grant GA (1992) *Synthetic Peptides: A*
30 *User Guide*. W.H. Freeman and Co., N.Y. (1992); Bodansky M and Trost B., Ed. (1993) *Principles of Peptide Synthesis*. Springer-Verlag Inc., NY (which is herein incorporated by reference at least for material related to peptide synthesis). Alternatively, the peptide or

polypeptide is independently synthesized *in vivo* as described herein. Once isolated, these independent peptides or polypeptides can be linked to form a peptide or fragment thereof via similar peptide condensation reactions.

[203] For example, enzymatic ligation of cloned or synthetic peptide segments allow
5 relatively short peptide fragments to be joined to produce larger peptide fragments, polypeptides or whole protein domains (Abrahmsen L et al., *Biochemistry*, 30:4151 (1991)). Alternatively, native chemical ligation of synthetic peptides can be utilized to synthetically construct large peptides or polypeptides from shorter peptide fragments. This method consists of a two step chemical reaction (Dawson et al. *Synthesis of Proteins by Native Chemical Ligation. Science*,
10 266:776-779 (1994)). The first step is the chemoselective reaction of an unprotected synthetic peptide--thioester with another unprotected peptide segment containing an amino-terminal Cys residue to give a thioester-linked intermediate as the initial covalent product. Without a change in the reaction conditions, this intermediate undergoes spontaneous, rapid intramolecular reaction to form a native peptide bond at the ligation site (Baggiolini M et al. (1992) *FEBS Lett.*
15 307:97-101; Clark-Lewis I et al., *J.Biol.Chem.*, 269:16075 (1994); Clark-Lewis I et al., *Biochemistry*, 30:3128 (1991); Rajarathnam K et al., *Biochemistry* 33:6623-30 (1994)).

[204] Alternatively, unprotected peptide segments are chemically linked where the bond formed between the peptide segments as a result of the chemical ligation is an unnatural (non-peptide) bond (Schnolzer, M et al. *Science*, 256:221 (1992)). This technique has been
20 used to synthesize analogs of protein domains as well as large amounts of relatively pure proteins with full biological activity (deLisle Milton RC et al., *Techniques in Protein Chemistry IV*. Academic Press, New York, pp. 257-267 (1992)).

3. Processes for making the compositions

[205] Disclosed are processes for making the compositions as well as making the
25 intermediates leading to the compositions. There are a variety of methods that can be used for making these compositions, such as synthetic chemical methods and standard molecular biology methods. It is understood that the methods of making these and the other disclosed compositions are specifically disclosed.

[206] Disclosed are nucleic acid molecules produced by the process comprising linking
30 in an operative way a promoter element and a μ -opioid receptor element.

[207] Disclosed are nucleic acid molecules produced by the process comprising linking in an operative way nucleic acid molecules comprising sequences set forth in SEQ ID NO:2 and SEQ ID NO:4.

[208] Also disclosed are nucleic acid molecules produced by the process comprising
5 linking in an operative way nucleic acid molecules comprising sequences having 80% identity to sequences set forth in SEQ ID NO:2 and SEQ ID NO:4.

[209] Also disclosed are nucleic acid molecules produced by the process comprising linking in an operative way nucleic acid molecules comprising sequences that hybridizes under stringent hybridization conditions to sequences set forth in SEQ ID NO:2 and SEQ ID NO:4.

10 [210] Disclosed are nucleic acid molecules produced by the process comprising linking in an operative way a nucleic acid molecule comprising a sequence encoding a μ -opioid receptor peptide and a sequence controlling an expression of the sequence encoding the μ -opioid receptor peptide.

[211] Disclosed are nucleic acid molecules produced by the process comprising linking
15 in an operative way a nucleic acid molecule comprising a sequence encoding a μ -opioid receptor peptide wherein the μ -opioid receptor peptide has 80% identity to the peptide set forth in SEQ ID NO:1 or 3 and a sequence controlling expression of the sequences encoding the peptide.

[212] Disclosed are nucleic acid molecules produced by the process comprising linking in an operative way a nucleic acid molecule comprising a sequence encoding a μ -opioid receptor
20 peptide wherein the μ -opioid receptor peptide has 80% identity to the peptides set forth in SEQ ID NO:1 or 3 and, wherein any change from the sequences set forth in SEQ ID NO:1 or 3 are conservative changes and a sequence controlling expression of the sequences encoding the peptide.

[213] Disclosed are cells produced by the process of transforming the cell with any of
25 the disclosed nucleic acids. Disclosed are cells produced by the process of transforming the cell with any of the non-naturally occurring disclosed nucleic acids.

[214] Disclosed are any of the disclosed peptides produced by the process of expressing any of the disclosed nucleic acids. Disclosed are any of the non-naturally occurring disclosed peptides produced by the process of expressing any of the disclosed nucleic acids.
30 Disclosed are any of the disclosed peptides produced by the process of expressing any of the non-naturally disclosed nucleic acids.

[215] Disclosed are animals produced by the process of transfecting a cell within the animal with any of the nucleic acid molecules disclosed herein. Disclosed are animals produced by the process of transfecting a cell within the animal any of the nucleic acid molecules disclosed herein, wherein the animal is a mammal. Also disclosed are animals produced by the process of transfecting a cell within the animal any of the nucleic acid molecules disclosed herein, wherein the mammal is mouse, rat, rabbit, cow, sheep, pig, or primate. Also disclosed are non human primates and non-human mammals.

[216] Also disclose are animals produced by the process of adding to the animal any of the cells disclosed herein.

E. Methods of using the compositions

1. Methods of using the compositions as research tools

[217] The disclosed compositions can be used in a variety of ways as research tools. For example, the disclosed compositions, the μ -opioid receptor constructs, and other nucleic acids, such as SEQ ID NOs:2 and 4 can be used to produce organisms, such as transgenic or knockout mice, which can be used as model systems for the study of pain.

2. Methods of gene modification and gene disruption

[218] The disclosed compositions and methods can be used for targeted gene disruption and modification in any animal that can undergo these events. Gene modification and gene disruption refer to the methods, techniques, and compositions that surround the selective removal or alteration of a gene or stretch of chromosome in an animal, such as a mammal, in a way that propagates the modification through the germ line of the mammal. In general, a cell is transformed with a vector which is designed to homologously recombine with a region of a particular chromosome contained within the cell, as for example, described herein. This homologous recombination event can produce a chromosome which has exogenous DNA introduced, for example in frame, with the surrounding DNA. This type of protocol allows for very specific mutations, such as point mutations, to be introduced into the genome contained within the cell. Methods for performing this type of homologous recombination are disclosed herein.

[219] One of the preferred characteristics of performing homologous recombination in mammalian cells is that the cells should be able to be cultured, because the desired recombination event occurs at a low frequency.

[220] Once the cell is produced through the methods described herein, an animal can be produced from this cell through either stem cell technology or cloning technology. For example, if the cell into which the nucleic acid was transfected was a stem cell for the organism, then this cell, after transfection and culturing, can be used to produce an organism which will contain the gene modification or disruption in germ line cells, which can then in turn be used to produce another animal that possesses the gene modification or disruption in all of its cells. In other methods for production of an animal containing the gene modification or disruption in all of its cells, cloning technologies can be used. These technologies generally take the nucleus of the transfected cell and either through fusion or replacement fuse the transfected nucleus with an oocyte which can then be manipulated to produce an animal. The advantage of procedures that use cloning instead of ES technology is that cells other than ES cells can be transfected. For example, a fibroblast cell, which is very easy to culture can be used as the cell which is transfected and has a gene modification or disruption event take place, and then cells derived from this cell can be used to clone a whole animal.

3. Therapeutic Uses

[221] Effective dosages and schedules for administering the compositions can be determined empirically, and making such determinations is within the skill in the art. The dosage ranges for the administration of the compositions are those large enough to produce the desired effect in which the symptoms disorder are effected. The dosage should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient, route of administration, or whether other drugs are included in the regimen, and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any contraindications. Dosage can vary, and can be administered in one or more dose administrations daily, for one or several days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products.

[222] Following administration of a disclosed composition, such as the disclosed constructs, for treating, inhibiting, or preventing pain, the efficacy of the therapeutic construct can be assessed in various ways well known to the skilled practitioner. For instance, one of ordinary skill in the art will understand that a composition, such as the disclosed constructs, disclosed herein is efficacious in treating pain or inhibiting or reducing the effects of pain in a subject by observing that the composition reduces the onset of the conditions associated with

these diseases. Furthermore, the amount of protein or transcript produced from the constructs can be analyzed using any diagnostic method. For example, it can be measured using polymerase chain reaction assays to detect the presence of construct nucleic acid or antibody assays to detect the presence of protein produced from the construct in a sample (e.g., but not limited to, blood or other cells, such as neural cells) from a subject or patient, or it can be measured by any of the methods disclosed herein for monitoring non-human pain, and through communication for human pain.

[223]

F. Examples

[224] It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

[225] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices and/or methods claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.), but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or near atmospheric.

1. Example 1 Orofacial pain

a) Methods and approaches

[226] Expression of the μ -opioid receptor in neurons involved in the central processing of orofacial pain will result in attenuation of nociception. FIV(μ -opioid receptor), a lentivirus capable of stably transducing terminally differentiated cells (neurons) with μ -opioid receptor, can be peripherally administered at orofacial sites, including the TMJ and the masseter muscle, exposed to nociceptive substances. The efficacy of the therapy can be assessed by assaying changes in muscle EMG as well as resistance to jaw opening.

**(1) Construct FIV(HUMOR) and confirm its ability to
transduce cells of neuronal origin**

[227] The opening reading frame of the HUMOR cDNA having the sequence set forth in SEQ ID NO:2 was (Raynor K, et al., *J Pharmacol Exp Ther.* 1995; 272:423-8) cloned into the pFIV vector using standard molecular biology methods generating pFIV(HUMOR) SEQID NO: 7 (See Figure 4). Constructs having two different promoters were made to drive the expression of HUMOR, the neuron specific enolase (NSE) promoter as well as the cytomegalovirus (CMV) promoter. The NSE promoter limits the expression of HUMOR selectively in neurons at constitutive levels, whereas the CMV promoter results in ubiquitous levels of expression independent of cell type. Clones were selected following DNA isolation and confirmation by multiple restriction digestions and direct DNA sequencing of both strands. pFIV(HUMOR) can be co-transfected with pPAK and pVSV-G vectors in 293H cells utilizing the Lipofectamine 2000® reagent (Gibco/BRL-Invitrogen) for virus production as previously described (Figure 3). Following 60 hours incubation, the supernatant will be collected, filtered and titered. Titers of 10^5 ip/ml are typical using this method. If required, the titers can be increased by ultra-centrifugation of the supernatant and re-suspension in normal saline.

[228] FIV(NSE-HUMOR) as well as FIV(CMV-HUMOR) efficacy can be tested on the N2a neuronal cell line, whereby cells can be seeded on 12 well plates and infected by FIV(HUMOR), followed by a 24 hour fresh media change. DNA, RNA and protein samples can be harvested using standard laboratory methods at 72 hours post infection. The presence of HUMOR gene copies in the infected samples can be assessed by Q-PCR, and the expression of HUMOR can be determined at the mRNA and protein levels by RT-PCR and western immunoblotting, respectively. In addition, other cells can be fixed by 4% paraformaldehyde for HUMOR protein expression detection and visualization by immunocytochemistry. Figure 1 or 2 demonstrates the expression of HUMOR in N2a cells following transient transfection and detection of its expression by immunocytochemistry utilizing a commercially available antibody raised against HUMOR (Research & Diagnostic Antibodies, Benicia CA Cat# AS-3942S). The FIV vector can infect and stably transduce sensory trigeminal neurons following local peripheral administration as demonstrated by the data in Figure 3. At the animal behavioral level, the targeted expression of HUMOR in orofacial sensory neurons can result in attenuation of nociceptive symptoms that readily can be measured as change (decrease) in EMG activity and decrease in resistance to jaw opening.

**(2) Targeted expression of μ -opioid receptor in orofacial
somatosensory neurons can attenuate orofacial pain following
administration of algesic substance**

[229] The feline immunodeficiency viral vectors FIV(NSE-HUMOR) and FIV(CMV-
5 HUMOR) can be employed in the transduction of trigeminal sensory neurons with the μ -opioid
receptor. Orofacial nociception can be quantitatively assessed by means of masticatory muscle
activity, as measured by electromyography (EMG), as well as resistance to mouth opening, as
measured by a digital force transducer. Table 6 summarizes the experimental conditions. In
brief, FIV(NSE-HUMOR) and FIV(CMV-HUMOR) can be injected (50 μ l of 10^7 ip/ml) in the
10 TMJ or masseter muscle of anesthetized animals, which can be returned to their cages after
recovery. Five weeks post-treatment, EMG bipolar percutaneous hook electrodes can be
inserted in the masseter and temporal muscles under anesthesia and an orthodontic Kobayashi
hook can be bonded to the mandibular incisors for attachment to the digital force gauge. The
animals can then be positioned in a custom made restraining device. Base line EMG and
15 resistance to jaw opening measurements can be taken for every animal. Four channels of
simultaneous EMG signal (right and left masseter and temporal muscles) and one channel for
the digital force gauge can be recorded. An A/D conversion card (NIO16E1, National
Instruments) for the EMG signal and a digital force gauge (FGF series, Kernco Instruments) can
be used for signal collection. A "blinded" observer can collect 10 seconds of EMG and digital
20 force measurements at 5, 10 and 15 mm of vertical opening five times simultaneously. Data can
be stored for later analysis. The appropriate animals can then be injected in the
temporomandibular joint and masseter muscle area with the algesic agent glutamate (2.5 μ mol
in saline) or vehicle solution (saline), and EMG and resistance to jaw opening recordings can be
taken. Data can be analyzed by two-way ANOVA with repeated measures. Student's t-test can
25 compare the two groups. These experiments replicate conditions seen in the human condition.
(Molin C., *Acta Odontol Scand* 1972; 30:485-99; Moller UM, et al., *Scand J Dent Res* 1984; 92:
64-83; Stohler CS, et al., *Helv Odontol Acta* 1985; 29:13-20; Stohler CS, et al., *Arch Oral Biol*
1988; 33: 175-82; Stohler CS. Temporomandibular disorders and related pain conditions. In:
Sessle, B.J., Bryant, P.S. and Dionne, R.A. Editors, 1995. Progress in pain research and
30 management Elsevier, Amsterdam, pp. 3-30; Lund JP, Stohler CS. Effect of pain on muscular
activity in temporomandibular disorders and related conditions. In: Stohler, C.S., Carlson, D.S.,
editors. Biological and Psychological aspects of orofacial pain, craniomandibular growth series.
Ann Arbor, MI: University of Michigan, 1994. pp. 75-91; Hoshio T. Senior Thesis, Division

of Orthodontics, Eastman Department of Dentistry, University of Rochester School of Medicine and Dentistry. Rochester, New York; Balkhi KM, et al., *J Orofacial Pain* 1993;7:89-97).

[230] HUMOR expression can be confirmed following the experimental recordings in deeply anesthetized animals that can be sacrificed via trans-cardial perfusion of 4% paraformaldehyde, and the brain stem, trigeminal ganglia, masseter muscle and TMJ can be excised and collected for further analysis. Specifically, we can characterize the expression of HUMOR in the central and peripheral tissues utilizing immunocytochemistry as depicted by our preliminary data. Cell type specificity can be accomplished by double immunofluorescence, as previously demonstrated by our laboratory. Potential deleterious effects of HUMOR expression (or over-expression in the CMV driven gene) to neurons or other cells can be examined by Hoechst staining and confirmed by TUNEL (in the case of cell death) as well as by the expression of neuron specific housekeeping genes, such as neuron specific enolase as well substance P (expressed selectively by sensory neurons).

G. TABLE 6 Summary of experimental conditions

ALGESIC AGENT	SITE of NOCICEPTION	TREATMENT	N
Glutamate	Temporomandibular Joint	FIV(NSE-HUMOR)	10 each
saline	Masseter muscle	FIV(CMV-HUMOR)	
		morphine	
		FIV(NSE-HUMOR)+opioid	
		FIV(CMV-HUMOR)+opioid	
		saline	

[231] Patients presenting with orofacial pain from TMJ disorders have been characterized as having decreased bite and chewing forces and limited jaw opening. For example, reduced grip strength (Norsdenskiold UM and Grimby G., *Scand J Rheumatol* 1993; 22: 14-9.), sustained jaw closing pain (Jow RW and Clark GT., *Arch Oral Biol* 1989; 34: 857-62.) and reduced bite strength (Molin C., *Acta Odontol Scand* 1972; 30:485-99) have all been reported in patients with muscle pain. The reduction in muscle force exertion associated with myalgia has been suggested to be due to reduced activity of agonist muscles and increased activity of antagonist muscles (Molin C., *Acta Odontol Scand* 1972; 30:485-99; Stohler CS, et al., *Helv Odontol Acta* 1985; 29:13-20; Stohler CS, et al., *Arch Oral Biol* 1988; 33: 175-82; Stohler CS. Temporomandibular disorders and related pain conditions. In: Sessle, B.J., Bryant, P.S. and Dionne, R.A. Editors, 1995. Progress in pain research and management Elsevier, Amsterdam, pp. 3-30; and Lund JP, Stohler CS. Effect of pain on muscular activity in temporomandibular disorders and related conditions. In: Stohler, C.S., Carlson, D.S., editors.

Biological and Psychological aspects of orofacial pain, craniomandibular growth series. Ann Arbor, MI: University of Michigan, 1994. pp. 75-91.) Keil et al., (Keil LJ, et al., *Pain* 2000; 85:333-43) have demonstrated that forelimb grip force reduction is a behavioral index of hyperalgesia in the carrageenan model of muscle hyperalgesia. This would translate to reduction of bite force and an increase in antagonist muscle activity in the orofacial region. These experiments have been done in humans (Lund JP, Stohler CS. Effect of pain on muscular activity in temporomandibular disorders and related conditions. In: Stohler, C.S., Carlson, D.S., editors. Biological and Psychological aspects of orofacial pain, craniomandibular growth series. Ann Arbor, MI: University of Michigan, 1994. pp. 75-91; Hoshio T. Senior Thesis, Division of Orthodontics, Eastman Department of Dentistry, University of Rochester School of Medicine and Dentistry. Rochester, New York; and Balkhi KM, et al., *J Orofacial Pain* 1993;7:89-97.). Hoshio has also demonstrated that compared to controls, patients with TMD demonstrate decreased bite (201 and 223 mV for the masseter muscles respectively in asymptomatic volunteers and 128 and 153 mV for symptomatic patients). Balkhi (Balkhi KM, et al., *J Orofacial Pain* 1993;7:89-97) demonstrated that chewing force was decreased in patients with pain (113 and 102 mV for deliberate right and left side chewing of gum masseter muscles respectively in asymptomatic volunteers and 85 and 83 mV for symptomatic patients). It has been demonstrated that there was increased jaw muscle activity of antagonists during jaw opening. These data clearly demonstrate that patients with TMD have similar characteristics and that decreased bite force and chewing activity are a reflection of somatic pain.

[232] Mice (C57/B6) can be employed to make the transgenic mice.

a) Injection of replication defective FIV(HUMOR) vectors

[233] The animals, such as pups, can be anesthetized with ketamine (60 mg/Kg) and xylazine (5 mg/Kg) IP. To verify the induction of surgical anesthesia, a toe is pinched in order to test for reflex withdrawal. One type of injection will be performed in two distinct areas. Under surgical plane of anesthesia the animals, such as mice, can be injected with 50 μ l of 10⁷ ip/ml of FIV(HUMOR) using a 1ml syringe with a 27^{1/2} gage needle directly into the temporomandibular joint and masseter muscle. The animals can be identified by ear punching. Animals can be held at the base of the tail with distal portion of tail situated on surface of nestlet, for example. Using a straight edge blade, ~7mm of distal tail can be removed, and the mouse can be placed in a cage and the tail specimen stored in a vial labeled by mouse ID# and sex. The mice are euthanised with sodium pentobarbital (200 mg/kg).

[234] Fixation by intracardial transfusion can be performed. Upon exposure of the heart, the right atrium can be clipped and the left ventricle can be catheterized with a 17 gage needle through which 50ml of 4% paraformaldehyde solution in phosphate buffered saline can be transfused into the animal. The liver, spleen, kidney and brain can be dissected and post-
5 fixed until sectioned for histology. The middle part of the cranium, including the cranial base (sphenoid, ethmoid, maxilla) can also be dissected, demineralized by immersion into an EDTA solution and section for histology.

2. Example 2: Non-Primate Lentiviral Vector Administration in the TMJ

[235] Disclosed herein are the effects lentiviral vectors on the temporomandibular joint.
10 Defective feline immunodeficiency virus capable of infecting dividing as well as terminally differentiated cells with the reporter gene lacZ, the expression of which was studied by means of PCR, X-gal histochemistry and β -galactosidase immunocytochemistry were injected into the articular joint space. The results showed successful transduction of hard and soft tissues of the temporomandibular joint. Interestingly, a subset of primary sensory neurons of the ipsilateral
15 trigeminal ganglion also stained positive for the reporter gene, presumably following uptake of the lentiviral vector by peripheral nerve fibers and retrograde transport to the nucleus. These findings indicate that transfer of anti-nociceptive genes, and disclosed herein, genes such as the opioid receptors, can be transferred into nerve cells and relieve pain. For example, lentiviral vectors can serve as the platform for the transfer of anti-nociceptive genes for the management
20 of temporomandibular joint pain.

a) Materials and Methods

(1) Vector Construction and Packaging

[236] The defective, vesicular stomatitis (VSV-G) pseudotyped, feline immunodeficiency virus, FIV(lacZ), capable of transducing dividing, growth-arrested as well as
25 post-mitotic cells (neurons) with the reporter gene lacZ driven by the ubiquitous cytomegalovirus promoter, CMV (Poeschla EM, et al. (1998). *Nature Med* 4: 354-357) was employed. The vectors were kindly donated to us by Dr. Wong-Staal, University of California at San Diego. A schematic description of the vector is depicted in Figure 6A. In addition, a control FIV(Δ' lac) vector carrying an inactive β -galactosidase was constructed by deleting the
30 first 1,000 bp of the lacZ gene (3.75 kb in total), including the transcription initiation site (Fig. 6A). Specifically, the FIV(lacZ) vector was digested *in vitro* with the SstII and Cla I restriction

enzymes overnight at 37°C, followed by agarose gel purification. The ends of the backbone DNA were blunted with the T4 DNA polymerase (Invitrogen, Carlsbad CA) and ligated with T4 ligase (Invitrogen) according to manufacturer's instructions. The FIV(lacZ) and FIV(Δ' lac) vectors were transiently co-transfected along with the packaging and VSV-G vectors into 293H cells (GIBCO/BRL) cultured in DMEM (Invitrogen) plus 10% FBS (Gemini, Woodland CA) using the Lipofectamine 2000 reagent per manufacturer's instructions (Invitrogen), and followed by a fresh media change supplemented by non-essential amino acids (Invitrogen). Sixty hours post-transfection, the supernatant was collected, filtered through .45mm *Surfil®-MF* filter (Corning Separations Division, Acton MA), aliquoted and frozen until further use. Titering was performed on CrfK cells (American Tissue Culture Collection; Manassas, VA) cultured in 24 well tissue culture plates, and assessed at 5×10^7 blue forming units (bfu) / mL by X-gal histochemistry.

(2) Animal Injections

[237] All methods pertinent to animal utilization were approved by the University Committee on Animal Resources. Specifically, 12 male mice, C57BL/6J, under surgical plane of anesthesia (ketamine 60mg/Kg and xylazine 5 mg/Kg administered intraperitoneally) received a single injection of 5×10^6 FIV(lacZ) infectious particles (100 μ l of stock solution) in the joint space of the right TMJ. Four additional mice received a single injection of 5×10^6 FIV(Δ' lac) infectious particles (100 μ l of stock solution) in the joint space of the right TMJ. In brief, the hair of the skin covering the right TMJ was shaved and the skin cleaned with Betadine solution. The joint was approached with an antero-posterior incision between the posterior end of the zygomatic arch and the ear cartilage, followed by a blunt dissection to expose the zygomatic arch and the posterior margin of the articular eminence. The joint space was not exposed during this procedure. The posterior margin of the eminence was identified by palpation and a 1-ml tuberculin syringe with a 271/2 gage needle was employed to inject the experimental solutions in the joint. This surgically assisted intra-articular injection technique was utilized to minimize leakage or spreading of the injectable solution beyond the articular space. (Kyrkanides S, et al. (2002). *J Orofac Pain* 16: 229-235). In addition, 2 mice that received 100 μ l saline injection served as controls. Forty-five days following treatment, the mice were deeply anesthetized by pentobarbital (100mg/Kg IP) and euthanised by transcardial perfusion of 4% paraformaldehyde in phosphate buffered saline (PBS) (Kyrkanides S, et al. (2002a). *J Orofac Pain* 16: 229-235, Kyrkanides S, et al. (2002). *Mol Brain Res* 104: 159-169).

The trigeminal ganglia and brain stem were dissected and sectioned at 20 μ m using a freezing microtome. The TMJ joints were also dissected, decalcified in an EDTA buffered solution, embedded in paraffin and cut at 8 μ m sections. All tissues were stored at -20°C until further processed.

5

(3) X-Gal Histochemistry

[238] Sections of trigeminal ganglia were processed by X-gal histochemistry and evaluated under light microscopy. Specifically, the sections were washed in 0.15M phosphate buffered saline (PBS) pH 7.2 for 60 min, followed by overnight processing in a staining solution containing 5-bromo-4chloro-3-indolyl- β -D-galactopyranoside (1mg/ml), potassium ferricyanide 10 (3mM), potassium ferrocyanide (3mM), NP-40 (0.02%) in 0.1M PBS pH 7.2 (Invitrogen) and $MgCl_2$ (1.3mM). The tissue was then washed in PBS for 30 min, and briefly rinsed with dH_2O . Considerable attention was given so that only the bacterial form of β -galactosidase was detected. The slides were cover slipped with DPX mounting medium (Fluka, Neu-Ulm, Switzerland) and examined under a light microscope (BX51 Olympus; Tokyo, Japan). Color 15 microphotographic images were captured in TIFF 16-bit format using a *SPOT RT Color* CCD digital camera attached onto the microscope and connected to a PC computer.

(4) Cell Counting

[239] The mouse ganglia (1.5mm X 2mm X 3mm) were sectioned sagittally on a freezing cryotome along their long axis into 20 μ m thick sections. A total of 42 sections were 20 approximately produced from each ganglion, which were sequentially collected onto 3 glass slides, whereby each slide contained representative ganglion sections 60 μ m apart of each other. One glass slide of each ganglion was processed by X-gal histochemistry and was employed in cell counting: all X-gal positive (blue) cells were counted on each tissue section on the slides. Since the tissue sections were 60 μ m apart, counting all blue cells on a single slide gave a 25 representative number of infected cells in each ganglion while avoiding overlap between sections and subsequently any "double counting".

(5) Immunocytochemistry

[240] Tissue sections from trigeminal ganglia were analyzed by immunocytochemistry employing a rabbit anti β -galactosidase polyclonal antibody (Chemicon INTL, Temecula CA). 30 In brief, sections were washed in BPS for 60 min followed by a 30 min blocking step in normal goat serum (4% in PBS) and overnight incubation in the primary antibody solution containing

rabbit anti β -galactosidase polyclonal antibody (1:2,500), 0.5% Triton-X, 4% normal goat serum (Invitrogen), 1% bovine serum albumin (Sigma; St Louis, MO) in PBS. The next morning the tissue was washed in PBS for 60 min, followed by a 30 min blocking step and incubated for 90 min in the secondary antibody solution containing a goat anti-rabbit polyclonal antibody
 5 (1:2,000), Triton-X (0.5%) and normal goat serum (0.15%) in PBS. Subsequently, the tissue was washed in PBS for 30 min and incubated in a avidin-biotin complex solution (ABC kit; Vector Laboratories, Burlingame CA), and was then washed in 0.1M sodium acetate buffered solution (pH 7.4) for 30 min. The tissue was then reacted in a DAB (3,3'-diaminobenzidine) – Nickel solution in 0.1M sodium acetate buffered solution (pH 7.4) for 5 min, followed by a 15
 10 min wash in PBS (Kyrkanides S, et al. (2002). *J Orofac Pain* 16: 229-235, Kyrkanides S, et al. (2002). *Mol Brain Res* 104: 159-169). The glass slides were then dehydrated through multiple ethanol solutions, cleared through xylene and cover-slipped using DPX permanent mounting medium. The tissue sections were then studied under a light microscope and microphotographic images were captured as described above.

15 [241] Tissue sections from the temporomandibular joints were first deparaffinized by immersion in a series of xylenes and alcohols, followed by antigen retrieval processing (95°C heating for 15 sec in 0.1 M Tris-HCL buffer pH 8.9) and processing employing the aforementioned immunocytochemical method.

(6) Polymerase Chain Reaction (PCR)

20 [242] The DNA from the left and right trigeminal ganglia of 8 mice (4 control and 4 experimental) was extracted employing the Trizol reagent (Invitrogen) according to manufacturer's instructions. The concentration of the recovered DNA ranged between 17-50 ng/ μ l, and was analyzed for the presence of viral DNA by PCR employing the following primer sets. Detection of FIV viral DNA (Fig.6A): 5'TTT TTC CAG TTC CGT TTA TCC (SEQ ID
 25 NO:35) and TTT ATC GCC AAT CCA CAT CT^{3'} SEQ ID NO. 36 (T_A =58°C; 40 total cycles). Detection of active β -galactosidase gene (Fig. 6A): 5'CCC ATA GTA ACG CCA ATA GG (SEQ ID NO:37) and AAA TGT GAG CGA GTA ACA ACC^{3'} SEQ ID NO. 38 (T_A =59.6°C; 45 total cycles). Detection of genomic DNA was performed utilizing primers designed for the murine G3PDH house keeping gene: ACC ACA GTC CAT GCC ATC AC SEQ ID NO. 39 and
 30 TCC ACC ACC CTG TTG CTG TA SEQ ID NO. 40 (T_A =58°C; 30 cycles). A total of 400 ng was used as DNA template in the PCR reactions. The PCR products were analyzed by agarose

gel (1% w/v) electrophoresis and the images were captured utilizing a KODAK Image Analysis system (Rochester NY).

b) RESULTS

(1) Intra-articular FIV injection resulted in transduction of hard and soft tissues

[243] FIV(lacZ) injection into the TMJ articular space resulted in transfer of the reporter gene lacZ via the lentiviral vector in cells located within the articular capsule. Specifically, cells of the TMJ meniscus, presumably fibroblasts, expressed bacterial β -galactosidase as it was assessed by immunocytochemistry employing appropriate polyclonal antibodies. In addition, cells located in the hypertrophic zone of the condyle, primarily comprised of cartilaginous cells, as well as perivascular cells, including endothelial cells and possibly osteocytes, also stained positive for bacterial β -galactosidase (Fig. 5). There was lack of β -galactosidase in the contralateral joints as well as the saline injected animals. These results indicate FIV successfully infected and stably transferred the reporter gene to cells of hard and soft TMJ tissues.

(2) FIV injection into the TMJ resulted in transduction of trigeminal neurons

[244] Two FIV vectors were employed in our experiment: the wild type FIV(lacZ) and the mutated FIV(Δ' lac) (Fig. 6A). FIV(Δ' lac) is capable of transducing cells with an inactive form of the reporter gene β -galactosidase compared to FIV(lacZ) which carries a full-length lacZ (Fig. 6B & 6C). Injection of either FIV vector in the right TMJ of mice resulted in transduction of neurons located in the ipsilateral trigeminal ganglia as assessed by PCR (Fig. 7A). Full length lacZ gene was detected by PCR only in the FIV(lacZ) treated animals (Fig. 7B), accompanied by neuronal β -galactosidase expression as assessed by X-gal histochemistry. The X-gal staining was localized primarily in the posterolateral part of the ganglion within the cell bodies of cells that appear histologically as neurons (Fig. 8A & 8B). In fact, the cell bodies of the primary sensory neurons that innervate the TMJ are known to localize in this part of the trigeminal ganglion. In contrast, FIV(Δ' lac) injected mice did not display any X-gal positive cells in the ganglia (Fig. 8C). Expression of bacterial β -galactosidase in the trigeminal ganglia was also confirmed by immunocytochemistry in the FIV(lacZ) (Fig. 8D) but not the FIV(Δ' lac) treated mice (Fig. 8E). Moreover, analysis of sections from the brain stem did not reveal any X-

gal positive staining, as it was anticipated since the vectors are defective, do not replicate and cannot infect second order neurons.

(3) Considerable number of trigeminal neurons were infected by FIV(lacZ)

5 [245] The mouse ganglia were on average of the following dimensions: 1.5mm X 2mm X 3mm. As described above, a total of 42 sections (20 μ m thick) were produced approximately from each ganglion, which were sequentially collected onto 3 glass slides, whereby each slide contained representative ganglion sections 60 μ m apart of each other. Consistently, 4 sections were identified containing X-gal (blue) cells on each glass slide, with an average of 93 (+/- 7.64
10 S.D.) blue cells per section. Therefore, we infer that there were approximately 93 cells X 4 sections X 3 glass slides = 1,116 transduced neurons in each right-sided ganglion in the FIV-injected animals. No X-gal positive cells were identified in the saline injected animals. These results suggest that from a total of 5×10^6 infectious particles injected into the articular TMJ space approximately 10^3 nerve fibers were infected resulting in lacZ expression, presumably
15 following uptake of the lentiviral vector by peripheral nerve fibers and retrograde transport to the nucleus.

[246] The results shown herein demonstrated that intra-articular injection of FIV(lacZ) resulted in successful gene transfer to articular TMJ surfaces as well as the joint meniscus. Interestingly, VSV-G does not require interaction between the viral envelope protein and a
20 specific membrane receptor, but instead interacts with a phospholipid component of the cell membrane leading to membrane-fusion mediated entry. This characteristic confers broad host-cell range for VSV-G pseudotyped viruses (Burns JC, et al. (1993). *Proc Natl Acad Sci USA* 90: 8033-8037; Carneiro FA, et al. (2002). *J Virol* 76: 3756-3764). Therefore, it is possible that FIV vectors demonstrate higher infectivity for TMJ tissues than previously described viral vectors
25 (Kuboki T, et al. (1999). *Arc Oral Biol* 44: 701-709), as well as result in prolonged transgene expression secondary to stable transgene integration (Poeschla EM, et al. (1998). *Nature Med* 4: 354-357).

[247] The efficacy of VSV-G pseudotyped FIV vectors to transduce peripheral tissues (Kang Y, et al.(2002). *J Virol* 76: 9378-9388), as well as the brain (Bloemer U, et al. (1997). *J*
30 *Virol* 71: 6641-6649) and cerebellum (Alisky JM, et al. (2002) *Mol Neurosci* 11: 2669-2673) has been previously demonstrated. The present observations of cells staining positively for X-gal in the trigeminal ganglion ipsilateral to the site of injection indicates that FIV virions were

taken up by peripheral nerve projections of trigeminal sensory neurons that lead to infection and expression of the reporter gene lacZ by these neurons. Therefore, VSV-G pseudotyped lentiviruses, such as the defective feline or human immunodeficiency virus, can serve as the platform for the transfer of anti-nociceptive genes to temporomandibular joint tissues as well as the neurons that innervate these structures.

3. Example 3:

a) Vector construction

[248] The ViraPower™ Lentiviral Expression System that can create a replication incompetent HIV-1-based lentivirus (Invitrogen, Carlsbad CA) was employed. This system can deliver and express NSE/Human- μ -opioid receptor in either dividing or non-dividing mammalian cells. First the pLenti6/V5-D-TOPO vector was reconstructed by insert PCR product which was generated base on the multiple cloning site of pIRES vector (Clontech Inc, Palo Alto CA) with the PCR primers: MCS-upper primer 5'-CACCTAATACGACTCACTATAGG-3' SEQ ID NO. 41 and MCS-lower primer 5'-CATTAAACCCTCACTAAAG-3' SEQ ID NO. 42. This 707 bp PCR product was purified and cloned into pLenti6/V5-D-TOPO vector directional (CACC, 4 base pair with overhang sequence will anneal to the GTGG sequence in the pLenti6/V5-D-TOPO vector). Multiple cloning sites were used as the template for PCR amplification to insert NheI site to the 5' end and a multiple enzyme digestion sequence followed by a NotI site to the 3' end of fragment. Then the CMV promoter of pLenti6/V5-D-TOPO was removed with ClaI and SpeI restriction enzyme digestions, followed by isopropanol DNA purification. Both ends of the re-constructed vector were blunted with T4 DNA polymerase (Invitrogen, Carlsbad CA) and ligated with T4 DNA ligase (Invitrogen) according to manufacturer's instruction.

[249] The NSE promoter was originally from pTR-NT3myc-NSE vector (Described in Peel AL. et al., Gene Therapy. 4(1):16-24, 1997). The NSE promoter sequence can be found in SEQ ID NO:52.). The NSE fragment was cut out with BglII and HindIII restriction enzyme digestions. The BglII site of this fragment was blunted. Later on, NSE fragment (2050 bp) was ligated into HindIII and blunted XhoI sites of pBluescript II KS+/- phagemid to form pBluescript II KS-NSE.

[250] Human- μ -opioid receptor (HUMOR) DNA was from pcDNA3-Human- μ -opioid receptor (Wang JB, et al., PNAS USA 90:10230-4) The sequence of this particular Human- μ -opioid receptor is found in SEQ ID NO:53.

[251] This 1.6 Kb fragment was cut out of the vector with EcoRV and XbaI, and
5 inserted into EcoRV/XbaI sites of pBluescript II KS-NSE to form pBluescript II KS-NSE-HUMOR. This structure was digested with KpnI and blunted, followed by the digestion of XbaI. The whole NSE-Human μ -opioid fragment was ligated into pIRES plasmid at XbaI and EcoRI (blunt) sites, and become pIRES- NSE-Human μ -opioid vector. To Insert the NSE-human μ -opioid receptor genes into the constructed pLenti6/V5-D-TOPO without CMV
10 promoter, pIRES-NSE-human μ -opioid was digested with NheI and SalI restriction enzymes and ligated into NheI/SalI sites of pLenti6/V5-d-TOPO at 14°C overnight and pLenti6/V5-D-TOPO-NSE-HUMOR vector was constructed.

[252] In order to increase the efficacy of virus packaging, cPPT sequence was added to the front of NSE of the pLenti6/V5-D-TOPO-NSE-HUMOR. Plasmid pLP1 (SEQ ID NO:49)
15 was used as template to PCR amplifying cPPT fragments with ClaI and NheI sites at both ends (upper primer 5'-atcgcgatcgcctagcttttaaagaaaaggggg-3' SEQ ID NO. 43 and lower primer 5'-taatcgcgatcgaagcaaaattttgaattttgtaattg-3' SEQ ID NO. 44). The PCR products were digested with NheI, and resulting fragment was ligated into the NheI site of pLenti6/V5-D-TOPO-NSE-HUMOR to generate pLenti6/V5-D-TOPO-cPPT-NSE-HUMOR plasmid at 4°C overnight. A
20 schematic description of the vector is disclosed herein.

[253] The pLenti6/v5-D-TOPO-cppt-NSE-HUMOR (SEQ ID NO:48) were transiently co-transfected along with the three packaging plasmids, pLP1(SEQ ID NO:49), pLP2 (SEQ ID NO:50),and pLP/VSVG(SEQ ID NO:51), into 293FT cells (Invitrogen) cultured in DEME (Invitrogen) plus 10% FBS (Gemini, Woodland CA). After 24 hours, the medium was replaced
25 with fresh medium supplemented with non-essential amino acid (Invitrogen). Seventy hours post-transfection, the supernatant was collected and filtered through 0.45 μ m Acrodisc 25 mm syringe filter (Pall Corporation, Gelman Laboratory). Aliquots of virus were frozen at -80°C until further use. Titering of the virus was performed on NIH3T3 cells cultured in 6 well tissue plates and assessed at 3×10^3 colonies (bfu)/ml by blasticidin selection.

b) Infection of albino neurons cells with Lenti6/NSE-Humor virus

[254] Neuro-2 α cell was plated into 6 well plate and cultured in MEM (GIBCO/BRL) with 10% FBS. To infected Lenti6/NSE-HUMOR virus, 1ml viral solution (3×10^3 ip) with 6 μ g polybrene solution was added to N2 α cell culture. After overnight plating, the medium was
5 changed to regular MEM with 10% FBS. The cells were harvested after 96 hours infection.

c) RT Polymerase Chain Reaction (RT-PCR)

[255] The total RNA of the N2 α cells infected with Lenti6/NSE-HUMOR virus was extracted with Trizol reagent (Invitrogen) according to manufacture instruction. 5 μ g total RNA was used to syntheses first strain DNA with SuperScriptTM First-Strand Synthesis System for
10 RT-PCR (Invitrogen). Analysis of the presence of HUMOR gene was done by PCR, employing the following primer sets: 5'-GAATTACCTAATGGGAACATGG-3' (SEQ ID NO:45) and 5'-GCAGACGATGAACACAGC-3' (SEQ ID NO:46) ($T_A = 56^\circ\text{C}$, total 30 cycles). G3PDH house keeping gene was used as quantity PCR control. Detection of genomic G3PDH DNA was
performed with primers 5'-ACCACAGCAATCAC-3' (SEQ ID NO:47) and 5'-
15 TCCACCACCCTGTTGCTGTA-3' (SEQ ID NO:40) ($T_A = 58^\circ\text{C}$, 30 cycles). The PCR products were analyzed by agarose gel (1% w/v) electrophoresis and imaged were captured by a KODAK image analysis system (Rochester NY).

H. Sequences

1. SEQ ID NO:1 Homo sapiens HUMOR, protein Genbank Accession No.
2. SEQ ID NO:2 Homo sapiens HUMOR, cDNA Genbank Accession No
3. SEQ ID NO:3 Murine HUMOR, protein Genbank Accession No
- 5 4. SEQ ID NO:4 Murine HUMOR, cDNA Genbank Accession No
5. SEQ ID NO:5 : human kappa opioid receptor cDNA
6. SEQ ID NO:6 human delta opioid receptor cDNA sequence
7. SEQ ID NO:7 FIV(Opioid receptor construct)
8. SEQ ID NO:8 FIV(LacZ) (a construct can be used for nerve
10 transduction)
9. SEQ ID NO:9 HUMOR degenerate cDNA G to A change at position 94
10. SEQ ID NO:10: HUMOR polypeptide conservative substitution of
Val32 to I32
11. SEQ ID NO:11: Neuron specific enolase promoter
- 15 12. SEQ ID NO:12 FIV backbone
13. SEQ ID NO:13: Packaging vector
14. SEQ ID NO:14 FIV-NSE-HUMOR -pA
15. SEQ ID NO:15: Mu-opioid RECEPTOR Bovine ACCESSION
NP_776833
- 20 16. SEQ ID NO:16: Bos taurus mu opioid receptor mRNA, complete cds.
ACCESSION U89677
17. SEQ ID NO:17:mu opioid receptor - mouse.

18. SEQ ID NO:18: Mus musculus mu opioid receptor cDNA, complete cds
ACCESSION U19380
19. SEQ ID NO:19: mu opioid receptor – rat ACCESSION I56504
20. SEQ ID NO:20: Rat mu opioid receptor mRNA, complete cds.
- 5 21. SEQ ID NO:21: mu opioid receptor [Sus scrofa] porcine ACCESSION
AAB53770
22. SEQ ID NO:22: Sus scrofa porcine mu opioid receptor mRNA, complete
cds ACCESSION AF521309
23. SEQ ID NO:23: DELTA-opioid RECEPTOR ACCESSION AAA18789
- 10 24. SEQ ID NO:24: Human delta opioid receptor mRNA, complete cds
ACCESSION U07882
25. SEQ ID NO:25: delta opioid receptor [Sus scrofa] ACCESSION
AAB39694
- 15 26. SEQ ID NO:26: delta opioid receptor [Rattus norvegicus] ACCESSION
AAA19939
27. SEQ ID NO:27: delta-opioid receptor [Mus musculus] ACCESSION
AAA37522
28. SEQ ID NO:28: Mus musculus delta-opioid receptor mRNA, cpl cds
Acc No. L06322
- 20 29. SEQ ID NO: 29: Homo sapiens (human) kappa opioid receptor
ACCESSION AAA63906
30. SEQ ID NO: 30: Human kappa opioid receptor (hKOR) mRNA,
complete cds Acc No. U17298
- 25 31. SEQ ID NO: 31: kappa opioid receptor [Mus musculus] ACCESSION
AAA39363
32. SEQ ID NO: 32: Mouse kappa opioid receptor mRNA, complete cds
ACCESSION L11065
33. SEQ ID NO: 33: kappa opioid receptor [Rattus norvegicus]
ACCESSION AAA41496

**34. SEQ ID NO: 34: Rattus norvegicus mRNA for kappa opioid receptor,
complete cds ACCESSION D16829**

I. References

- Alisky JM, Hughes SM, Sauter SL, Joly D, Dubensky Jr. TW, Staber, PD, et al. (2002) Transduction of murine cerebellar neurons with recombinant FIV and AAV5 vectors. *Mol Neurosci* 11: 2669-2673.
- 5 Baum BJ, Kok M, Tran SD, Yamano S (2002). The impact of gene therapy on dentistry: a revisiting after six years. *JADA* 133: 35-44.
- Bernick S (1962). The vascular and nerve supply to the temporomandibular joint of the rat. *Oral Surg* 15:488-492.
- 10 Bloemer U, Naldini L, Kafri T, Trono D, Verma IM, Gage FH (1997). Highly efficient and sustained gene transfer in adult neurons with a lentivirus vector. *J Virol* 71: 6641-6649.
- Burns JC, Fiedmann T, Driever W, Burrascanno M, Yee J-K (1993). Vesicular stomatitis virus G glycoprotein pseudotyped retroviral vectors: Concentration to very high titer and efficient gene transfer into mammalian and nonmammalian cells. *Proc Natl Acad Sci USA* 90: 8033-8037.
- 15 Capra NF (1987). Localization and central projections of primary afferent neurons that innervate the temporomandibular joint in cats. *Somatosensory Res* 4: 201-213.
- Carneiro FA, Bianconi L, Weissmueller G, Stauffer F, Da Poian AT (2002). Membrane recognition by vesicular stomatitis virus involves enthalpy-driven protein-lipid interactions. *J Virol* 76: 3756-3764.
- 20 Cox BM, Ginsburg M, Osman OH (1968). Acute tolerance to narcotic drugs in rats. *Br J Pharmacol* 245-256.
- Dreessen D, Halata Z, Strassmann T (1990). Sensory innervation of the temporomandibular joint in the mouse. *Acta Anat* 139:154-160.
- 25 Frommer J, Monroe CW (1966). The morphology and distribution of nerve fibers and endings associated with the mandibular joint of the mouse. *J Dent Res* 45:1762-1766.
- Kang Y, Stein CS, Heth PA, Sinn PL, Penisten AK, Staber PD, et al.(2002). In vivo gene transfer using a nonprimate lentiviral vector pseudotyped with ross river virus glycoproteins. *J Virol* 76: 9378-9388.
- 30 Kido MA, Kiyoshima T, Kondo T, Ayasaka N, Moroi R, Terada Y, Tanaka T (1993). Distribution of substance P and CGRP-like immunoreactive nerve fibers in the rat temporomandibular joint. *J Dent Res* 72:592-598.
- Kido MA, Kondo T, Ayasaka N, Terada Y, Tanaka T (1991). The peripheral distribution of trigeminal nerve fibers in the rat temporomandibular joint studied by an anterograde axonal transport method with wheat germ agglutinin-horseradish peroxidase. *Arch Oral Biol*
- 35 36:397-400.
- Klineberg I (1971). Structure and function of temporomandibular joint innervation. *Ann Royal Coll Surg Engl* 49:268-288

- Kuboki T, Nakanishi T, Kanyama M, Sonoyama W, Fujisawa T, Kobayashi K, et al. (1999). Direct adenovirus-mediated gene delivery to the temporomandibular joint in guinea-pigs. *Arc Oral Biol* 44: 701-709.
- 5 Kyrkanides S, Moore AH, Olschowka JA, Daeschner JC, Williams JP, Hansen JT, O'Banion MK (2002). Cyclooxygenase-2 modulates brain inflammation-related gene expression in central nervous system radiation injury. *Mol Brain Res* 104: 159-169.
- Kyrkanides S, Tallents RH, Macher DJ, Olschowka JO, Stevens SY (2002). Temporomandibular joint nociception: Effects of capsaicin on substance P-like immunoreactivity in the rabbit brain stem. *J Orofac Pain* 16: 229-235.
- 10 Martin WR, Eades CG (1961). Demonstration of tolerance and physical dependence in the dog following a short-term infusion of morphine. *J Pharmacol Exp Ther* 133: 262-270.
- Peel AL, Zolotukhin S, Schrimsher GW, Muzyczka N, Reier PJ. Efficient transduction of green fluorescent protein in spinal cord neurons using adeno-associated virus vectors containing cell type-specific promoters. [Journal Article] *Gene Therapy*. 4(1):16-24, 1997
- 15 Poeschla EM, Wong-Stall F, Looney DI (1998). Efficient transduction of nondividing human cells by feline immunodeficiency virus lentiviral vectors. *Nature Med* 4: 354-357.
- Pohl M, Braz J (2001). Gene therapy of pain: emerging strategies and future directions. *Eur J Pharmacol* 429: 39-48.
- 20 Romfh JH, Capra NF, Gatipon GB (1979). Trigeminal nerve and temporomandibular joint of the cat: A horseradish peroxidase study. *Exp Neurol* 65: 99-106.
- Sessle BJ, Hu JW (1991). Mechanisms of pain arising from articular tissues. *Can J Physiol Pharmacol* 69: 617-626.
- Thilander B (1964). Innervation of the temporomandibular disc in Man. *Acta Odont Scan* 22:151-156.
- 25 Wang JB, Imai Y, Eppler CM, Gregor P, Spivak CE, Uhl GR. (1993). mu opiate receptor: cDNA cloning and expression. *PNAS USA* 90:10230-4
- Wink CS, St'Onge M, Zimny ML (1992). Neural elements in the human temporomandibular articular disc. *J Oral Maxillofac Surg* 50:334-337
- 30 Wu CL, Garry MG, Zollo RA, Yang J (2001). Gene therapy for the management of pain: part II: molecular targets. *Anesthesiology* 95: 216-240.
- Yoshino K, Kawagishi S, Amano N (1998). Morphological characteristics of primary sensory and post-synaptic sympathetic neurons supplying the temporomandibular joint in the cat. *Arc Oral Biol*; 43: 679-686.

V. CLAIMS

What is claimed is:

1. A vector for delivering an opioid receptor to a nerve cell, comprising sequence encoding an opioid receptor and a vector backbone.
2. The vector of claim 1, wherein the opioid receptor is a μ -opioid receptor.
3. The vector of claim 2, wherein the μ -opioid receptor has a sequence with at least 80% identity to the sequence set forth in SEQ ID NO:1.
4. The vector of claim 2, wherein the μ -opioid receptor has a sequence with at least 85% identity to the sequence set forth in SEQ ID NO:1.
5. The vector of claim 2, wherein the μ -opioid receptor has a sequence with at least 90% identity to the sequence set forth in SEQ ID NO:1.
6. The vector of claim 2, wherein the μ -opioid receptor has a sequence with at least 95% identity to the sequence set forth in SEQ ID NO:1.
7. The vector of claims 1-6, wherein the composition further comprises a promoter.
8. The vector of claim 7, wherein the promoter is a nerve cell specific promoter.
9. The vector of claim 8, wherein the promoter is the neuron specific enolase promoter.
The vector of claim 7, wherein the promoter is a NSE promoter.
10. The vector of claim 7, wherein the vector backbone is a lentiviral vector backbone.
11. The vector of claim 10, wherein the lentiviral vector backbone is a feline immunodeficiency vector (FIV).
12. The vector of claim 11, wherein the FIV has a sequence with at least 80% identity to the sequence set forth in SEQ ID NO:7.
13. A cell comprising the vector of claims 1-12.
14. An animal comprising the cell of claim 13.
15. A cell comprising the integrated product of the vector of claims 1-12.
16. An animal comprising the cell of claim 15.
17. A method of reducing pain in a subject, comprising administering the vector of

claims 1-12, to the subject, wherein the vector transduces a nerve cell.

18. The method of claim 17, wherein administering the vector occurs at the point of pain.

19. The method of claim 17, wherein administering the vector occurs at the distal end of the nerve cell.

20. The method of claim 17, wherein administering the vector occurs in the peripheral nervous system.

21. The method of claim 17, wherein administering the vector occurs at the axon or axon terminal of the nerve cell.

22. The method of claim 17, wherein administering the vector occurs at the dendrite of the nerve cell.

23. The method of claim 17, wherein administering the vector occurs at the trigeminal ganglion.

24. The vector of claim 1, wherein the vector comprises the sequence set forth in SEQ ID NO:48.

25. The vector of claim 1, wherein the vector comprises the sequence set forth in SEQ ID NO: 7.

26. A method of producing the vector of claim 1, comprising linking the opiod receptor sequence operably to a promoter.

27. A method of producing the cell of claim 13, comprising transfecting the vector of claim 1 into the cell.

28. An animal produced by the process of administering the vector of claim 1 to the animal.

29. The vector of claim 7, wherein the vector backbone is an HIV vector backbone.





FIG. 1

**FIG. 2**




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8

RT-PCR of N2α Cells Infected with Lenti Viruses



C: plain N2α cells
 L: N2α cells infected with Lenti LacZ virus
 H: N2α cells infected with Lenti NSE-Humor virus
 +: pLenti6/NSE-Humor
 -: primers only

FIG. 9

SEQUENCE LISTING

<110> University of Rochester

Kyrkanides, Stephanos
Tallents, Ross H.<120> Treatment of Pain Through Expression of
Opioid Receptors

<130> 21108.0022P1

<150> 60/448,663

<151> 2003-02-19

<160> 54

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 400

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 1

Met	Asp	Ser	Ser	Ala	Ala	Pro	Thr	Asn	Ala	Ser	Asn	Cys	Thr	Asp	Ala	1	5	10	15
Leu	Ala	Tyr	Ser	Ser	Cys	Ser	Pro	Ala	Pro	Ser	Pro	Gly	Ser	Trp	Val	20	25	30	
Asn	Leu	Ser	His	Leu	Asp	Gly	Asn	Leu	Ser	Asp	Pro	Cys	Gly	Pro	Asn	35	40	45	
Arg	Thr	Asp	Leu	Gly	Gly	Arg	Asp	Ser	Leu	Cys	Pro	Pro	Thr	Gly	Ser	50	55	60	
Pro	Ser	Met	Ile	Thr	Ala	Ile	Thr	Ile	Met	Ala	Leu	Tyr	Ser	Ile	Val	65	70	75	80
Cys	Val	Val	Gly	Leu	Phe	Gly	Asn	Phe	Leu	Val	Met	Tyr	Val	Ile	Val	85	90	95	
Arg	Tyr	Thr	Lys	Met	Lys	Thr	Ala	Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	100	105	110	
Ala	Leu	Ala	Asp	Ala	Leu	Ala	Thr	Ser	Thr	Leu	Pro	Phe	Gln	Ser	Val	115	120	125	
Asn	Tyr	Leu	Met	Gly	Thr	Trp	Pro	Phe	Gly	Thr	Ile	Leu	Cys	Lys	Ile	130	135	140	
Val	Ile	Ser	Ile	Asp	Tyr	Tyr	Asn	Met	Phe	Thr	Ser	Ile	Phe	Thr	Leu	145	150	155	160
Cys	Thr	Met	Ser	Val	Asp	Arg	Tyr	Ile	Ala	Val	Cys	His	Pro	Val	Lys	165	170	175	
Ala	Leu	Asp	Phe	Arg	Thr	Pro	Arg	Asn	Ala	Lys	Ile	Ile	Asn	Val	Cys	180	185	190	
Asn	Trp	Ile	Leu	Ser	Ser	Ala	Ile	Gly	Leu	Pro	Val	Met	Phe	Met	Ala	195	200	205	
Thr	Thr	Lys	Tyr	Arg	Gln	Gly	Ser	Ile	Asp	Cys	Thr	Leu	Thr	Phe	Ser	210	215	220	
His	Pro	Thr	Trp	Tyr	Trp	Glu	Asn	Leu	Leu	Lys	Ile	Cys	Val	Phe	Ile	225	230	235	240

```

Phe Ala Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu
      245                      250                      255
Met Ile Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu
      260                      265                      270
Lys Asp Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Val
      275                      280                      285
Ala Val Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile
      290                      295                      300
Lys Ala Leu Val Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp
305                      310                      315                      320
His Phe Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val
      325                      330                      335
Leu Tyr Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe
      340                      345                      350
Cys Ile Pro Thr Ser Ser Asn Ile Glu Gln Gln Asn Ser Thr Arg Ile
      355                      360                      365
Arg Gln Asn Thr Arg Asp His Pro Ser Thr Ala Asn Thr Val Asp Arg
      370                      375                      380
Thr Asn His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
385                      390                      395                      400

```

<210> 2

<211> 1200

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 2

```

atggacagca ggcgtgcccc cacgaacgcc agcaattgca ctgatgcctt ggcgtactca 60
agttgtctccc cagcaccacag ccccggttcc tgggtcaact tgtcccactt agatggcaac 120
ctgtccgacc catgcggtcc gaaccgcacc gacctggcg ggagagacag cctgtgccct 180
ccgaccggca gtccctccat gatcacggcc atcacgatca tggccctcta ctccatcgtg 240
tgcgtggtgg ggctcttcgg aaacttctctg gtcatgtatg tgattgtcag atacaccaag 300
atgaagactg ccaccaacat ctacattttc aaccttgctc tggcagatgc cttagccacc 360
agtaccctgc ccttcagag tgtgaattac ctaatgggaa catggccatt tggaaccatc 420
ctttgcaaga tagtgatctc catagattac tataacatgt tcaccagcat attcaccctc 480
tgcaccatga gtgttgatcg atacattgca gtctgccacc ctgtcaaggc cttagatttc 540
cgtactcccc gaaatgccaa aattatcaat gtctgcaact ggatcctctc ttcagccatt 600
ggtcttcctg taatgttcat ggctacaaca aaatacaggc aaggttccat agattgtaca 660
ctaacattct ctcacccaac ctggtactgg gaaaacctgc tgaagatctg tgttttcatc 720
ttcgcttca ttatgccagt gctcatcatt accgtgtgct atggactgat gatcttgccg 780
ctcaagagtg tccgcatgct ctctggctcc aaagaaaagg acaggaatct tcgaaggatc 840
accaggatgg tgctgggtgt ggtggctgtg ttcacgtctt gctggactcc cattcacatt 900
tacgtcatca ttaaagcctt gggttacaatc ccagaaacta cgttccagac tgtttcttgg 960
cacttctgca ttgctctagg ttacacaaac agctgcctca acccagtcct ttatgcattt 1020
ctggatgaaa acttcaaacg atgcttcaga gagttctgta tcccaacctc ttccaacatt 1080
gagcaacaaa actccactcg aattcgtcag aacactagag accaccctc caccggccaa 1140
acagtggata gaactaatca tcagctagaa aatctggaag cagaaactgc tccgttgccc 1200

```

<210> 3

<211> 398

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =

synthetic construct

<400> 3

```

Met Asp Ser Ser Ala Gly Pro Gly Asn Ile Ser Asp Cys Ser Asp Pro
 1          5          10          15
Leu Ala Pro Ala Ser Trp Ser Pro Ala Pro Gly Ser Trp Leu Asn Leu
          20          25          30
Ser His Val Asp Gly Asn Gln Ser Asp Pro Cys Gly Pro Asn Arg Thr
          35          40          45
Gly Leu Gly Gly Ser His Ser Leu Cys Pro Gln Thr Gly Ser Pro Ser
          50          55          60
Met Val Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile Val Cys Val
          65          70          75          80
Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile Val Arg Tyr
          85          90          95
Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu
          100          105          110
Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Val Asn Tyr
          115          120          125
Leu Met Gly Thr Trp Pro Phe Gly Asn Ile Leu Cys Lys Ile Val Ile
          130          135          140
Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Cys Thr
          145          150          155          160
Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu
          165          170          175
Asp Phe Arg Thr Pro Arg Asn Ala Lys Ile Val Asn Val Cys Asn Trp
          180          185          190
Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met Ala Thr Thr
          195          200          205
Lys Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe Ser His Pro
          210          215          220
Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe Ile Phe Ala
          225          230          235          240
Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Ile
          245          250          255
Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu Lys Asp
          260          265          270
Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Val Ala Val
          275          280          285
Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile Lys Ala
          290          295          300
Leu Ile Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp His Phe
          305          310          315          320
Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val Leu Tyr
          325          330          335
Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe Cys Ile
          340          345          350
Pro Thr Ser Ser Thr Ile Glu Gln Gln Asn Ser Ala Arg Ile Arg Gln
          355          360          365
Asn Thr Arg Glu His Pro Ser Thr Ala Asn Thr Val Asp Arg Thr Asn
          370          375          380
His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
          385          390          395

```

<210> 4

<211> 1194

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 4

```

atggacagca ggcgcggccc agggaaacatc agcgactgct ctgacccctt agctcctgca 60
agttggtccc cagcacctgg ctccctggctc aacttgctcc acgttgatgg caaccagtcc 120
gacccatgcg gtcctaaccg caccggggctt ggccgggagcc acagcctgtg ccctcagacc 180
ggcagccctt ccatggtcac agccatcacc atcatggccc tctattctat cgtgtgtgta 240
gtgggcctct ttggaaactt cctgggtcatg tatgtgattg taagatatac caaaatgaag 300
actgccacca acatctacat tttcaacctt gctctggcag atgccttagc cactagcacg 360
ctgccctttc agagtgttaa ctacctgatg ggaacgtggc cctttggaaa catcctctgc 420
aagatcgtga tctcaataga ctactacaac atgttcacca gtatcttcac cctctgcacc 480
atgagtgtag accgctacat tgccgtctgc caccgggtca aggccttggg tttccgtacc 540
ccccgaaatg ccaaaattgt caatgtctgc aactggatcc tctcttctgc cattgggtctg 600
cccgtaatgt tcatggcaac cacaaaatac aggcaggggt ccatagattg caccctcact 660
ttctctcctc ccacatggta ctgggagaac ctgctcaaaa tctgtgtctt catcttcgcc 720
ttcatcatgc cggctcctcat catcactgtg tgttatggac tgatgatctt acgactcaag 780
agtgtccgca tgctgtcggg ctccaaagaa aaggacagga acctgcgcag gatcaccgag 840
atggtgctgg tggctcgtggc tgtattttatt gtctgctgga ccccatcca catctatgtc 900
atcatcaaag cactgatcac gattccagaa accactttcc agactgtttc ctggcacttc 960
tgcatgtgct tgggttacac aaacagctgc ctgaacccag ttctttatgc gttcctggat 1020
gaaaacttca aacgatgttt tagagagttc tgcatcccaa ctctctccac aatcgaacag 1080
caaaactctg ctcgaaatccg tcaaaacact agggaaacacc cctccacggc taatacagt 1140
gatcgaacta accaccagct agaaaatctg gaagcagaaa ctgctccatt gccc 1194

```

<210> 5

<211> 2423

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 5

```

accgcgctc gtacgtgccc ctccgcgggc agctcctgac tcatcggggg ctccgggtca 60
catgcgcccg cgcggcccta taggcgcctc ctccgcgggc cgccggggag ccgcagccgc 120
cgccgccact gccactcccg ctctctcagc gccgcgctcg ccaccgccac cgccaccgcc 180
actaccaccg tctgagctcg cagtcgccag atcccagcca tcatgtccat agagaagatc 240
tgggcccggg agatcctgga ctccgcgggc aacccacagc tggagggtgga tctctatact 300
gccaaaggtc ttttccgggc tgcagtgcgc agtggagcct ctacgggcat ctatgaggcc 360
ctggagctga gggatggaga caaacagcgt tacttaggca aagggtgcct gaaggcagt 420
gaccacatca actccaccat cgcgccagcc ctcatcagct caggtctctc tgtggtggag 480
caagagaaac tggacaacct gatgctggag ttggatggga ctgagaacaa atccaagttt 540
ggggccaatg ccactcctggg tgtgtctctg gccgtgtgta aggcaggggc agctgagcgg 600
gaactgcccc tgtatcgcca cattgctcag ctggccggga actcagacct catcctgcct 660
gtgccggcct tcaacgtgat caatggtggc tctcatgctg gcaacaagct ggccatgcag 720
gagttcatga tccctccagt gggagctgag agctttcggg atgccatgcg actaggtgca 780
gaggtctacc atacactcaa gggagtcac aaggacaaat acggcaagga tgccaccaat 840
gtgggggatg aagggtggctt tgcctccaat atcctggaga acagtgaagc cttggagctg 900
gtgaaggaag ccacgcacaa ggctggctac acggaaaaga tcgttattgg catggatgtt 960
gctgcctcag agttttatcg tgatggcaaa tatgacttgg acttcaagtc tccactgat 1020
ccttcccgat acatcactgg ggaccagctg ggggcaactc accaggactt tgtcagggac 1080
tatcctgtgg tctccattga ggaccattt gaccaggatg attgggctgc ctggtccaag 1140
ttcacagcca atgtaggat ccagattgtg ggtgatgacc tgacagtgac caacccaaaa 1200
cgtattgagc gggcagtggg agaaaaggcc tgcaactgtc tgctgctcaa ggtcaaccag 1260
atcggtcctg tcaactgaag catccaagcg tgcaagctgg cccaggagaa tggctggggg 1320
gtcatggtga gtcacgctc aggagagact gaggacacat tcattgctga cctggtggtg 1380
gggctgtgca caggccagat caagactggt gccccgtgcc gttctgaacg tctggctaaa 1440
tacaaccagc tcatgagaat tgaggaagag ctgggggatg aagctcgctt tgccggacat 1500
aacttccgta atcccagtgt gctgtgattc ctctgcttgc ctggagacgt ggaacctctg 1560

```

```

tctcatcctc ctggaacctt gctgtcctga tctgtgatag ttcacccctt gagatccctt 1620
gagccccagg gtgcccagaa cttccctgat tgacctgctc cgctgctcct tggettacct 1680
gacctcttgc tgtctctgct cgccctcctt tctgtgccct actcattggg gttccgcact 1740
ttccacttct tcctttctct ttctctcttc cctcagaaac tagaaatgtg aatgaggatt 1800
attataaaaag ggggtccgtg gaagaatgat cagcatctgt gatgggagcg tcagggttgg 1860
tgtgtcgagg tgtagagag ggaccatgtg tcaattgtgc tttgctcttg tcccacgtgt 1920
cttccacttt gcatatgagc cgtgaactgt gcatagtgtc gggatggagg ggagtgttgg 1980
gcatgtgata acgcctggct aataaggctt tagtgtattt atttatttat ttattttatt 2040
tgtttttcat tcatccatt aatcatttcc ccataactca atggcctaaa actggcctga 2100
cttgggggaa cgatgtgtct gtatttcatg tggctgtaga tccaagatg actggggtgg 2160
gaggtcttgc tagaatggga agggtcatag aaagggcctt gacatcagtt ctttgtgtg 2220
tactcactga agcctgcgtt ggtccagagc ggaggctgtg tgcctggggg agttttcctc 2280
tatacatctc tccccaaccc taggttccct gttcttctc cagctgcacc agagcaacct 2340
ctcactcccc atgccacgtt ccacagttgc caccacctct gtggcattga aatgagcacc 2400
tccattaaag tctgaatcag tgc                                     2423

```

<210> 6

<211> 1773

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 6

```

cdgaggagcc tgcgctgctc ctggctcaca gcgctccggg cgaggagagc gggcggaaccg 60
gggggctggg ccggtgcggg cggcgaggca ggcggacgag gcgagagac agcgggggcgg 120
ccggggcgcg gcacgcggcg ggtcggggcc ggctctgcc ttgccgctcc cctcgcgctcg 180
gatccccgcg ccaggcagc oggtggagag ggacgcggcg gacgccgga gccatggaac 240
cggccccctc cgccggcgcc gagctgcacc ccccgctctt cgccaacgcc tcggagcgcct 300
accctagcgc cttcccagc gctggcgcca atgcgctggg gccgccaggc gcgcggagcg 360
cctcgctcct cgccctggca atcgccatca ccgcgctcta ctggcccggtg tgcgcgctgg 420
ggctgctggg caacgtgctt gtcattgtcg gcatcgctcg gtacactaag atgaagacgg 480
ccaccaacat ctacatcttc aacctggcct tagccgatgc gctggccacc agcacgctgc 540
ctttccagag tgccaagtac ctgatggaga cgtggccctt cggcgagctg ctctgcaagg 600
ctgtgctctc catcgactac tacaatatgt tcaccagcat cttcacgctc accatgatga 660
gtgttgaccg ctacatcgct gtctgaccgc cctgtcaagg cctggacttc cgcacgcct 720
ccaaggccaa gctgatcaac atctgtatct ggttccctggc ctcaggcggtt ggcgtgcccc 780
tcatggatcat ggctgtgacc cgtccccggg acggggcagt ggtgtgcatg ctccagttcc 840
ccagccccag ctggtactgg gacacgggtg ccaagatctg cgtgttctc ttgccttcg 900
tggtgcccac cctcatcatc accgtgtgct atggcctcat gctgctgcgc ctgcgcagt 960
tgcgcctgct gtcgggctcc aaggagaagg accgcagcct gcggcgcatc acgcgcagt 1020
tgetggtggt tgtgggcgcc ttctgtgtgt gttgggcgcc catccacatc ttctcatcg 1080
tctggacgct ggtggacatc gaccggcgcg acccgctggt ggtggctgcg ctgcacctgt 1140
gcatcgcgct gggctacgcc aatagcagcc tcaaccccggt gctctacgct ttctcgacg 1200
agaacttcaa gcgctgcttc cgccagctct gccgcaagcc ctgcggccgc ccagacccca 1260
gcagcttcag ccgcgcccgc gaagccacgg ccgcgagcg tgtaccgcgc tgcaccccg 1320
ccgatggtcc cggcggtggc gctgccgcct gaccaggcca tccggcccc agacgcccct 1380
ccctagttgt acccgaggc cacatgagtc ccagtgggag gcgcgagcca tgatgtggag 1440
tggggcccagt agataggtcg gagggtttt ggaccgccag atggggcctc tgtttcggag 1500
acgggaccgg gccgctagat gggcatgggg tgggcctctg gtttggggcg aggcagagga 1560
cagatcaatg gcgcagtgc tctggtctg gtgccccgt ccacggctct aggtggggcg 1620
ggaaagccag tgaactcagg agaggagcg gacctgtggc tctacaactg agtccttaa 1680
cagggcatct ccaggaaggc ggggcttcaa ccttgagaca gcttcgggtt ctaacttgg 1740
gccggacttt cggagttggg ggggtccggg ccc                                     1773

```

<210> 7

<211> 9426

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 7

```

atgCGgtttt ggcagtacat caatgggCGt ggatagCGgt ttgactcacg gggatttcca 60
agtctccacc ccattgacgt caatgggagT ttgttttggc accaaaatca acgggacttt 120
ccaaaatgtc gtaacaactc cgccccattg acgcaaattg gcggtaggcg tgtacgggtg 180
gaggtctata taagcagagc tctgtgaaac ttcgaggagt ctctttgttg aggacttttg 240
agttctccct tgaggctccc acagatacaa taaatatttg agattgaacc ctgtcgagta 300
tctgtgtaat cttttttacc tgtgagggtc cggaatccgg gccgagaact tcgcagttgg 360
cgcccgaaca gggacttgat tgagagtgat tgaggaagtg aagctagagc aatagaaagc 420
tgtaaagcag aactcctgct gacctaaata gggaagcagt agcagacgct gctaacagtG 480
agtatctcta gtgaagcgga ctcgagctca taatcaagtc attgtttaaa ggccagata 540
aattacatct ggtgactctt cgcggaacct caagccagga gattcgccga gggacagtca 600
acaaggtagg agagattcta cagcaacatg gggaatggac aggggCGaga ttggaaaatg 660
gccattaaga gatgtagtaa tgttgctgta ggagtagggg ggaagagtaa aaaaatttgg 720
gaagggaatt tcagatgggc cattagaatg gctaattgat ctacaggacg agaacctggT 780
gatataccag agacttttaga tcaactaagg ttggttattt gcgatttaca agaaagaaga 840
gaaaatttgg gatctagcaa agaaattgat atggcaattg tgacattaaa agtctttgCG 900
gtagcaggac ttttaaatat gacgggtgtc tactgtctgt gcagctgaaa atatgtattc 960
tcaaatggga ttagacacta ggccatctat gaaagaagca ggtggaaaag aggaaggccc 1020
tccacaggca tatcctattc aaacagtaaa tggagtacca caatatgtag cacttgaccc 1080
aaaaatgggt tccattttta tggaaaaggc aagagaagga ctaggagggt aggaagttca 1140
actatggttt actgccttct ctgcaaatTT aacacctact gacatggcca cattaataat 1200
ggccgcacca ggggtgcgctg cagataaaga aatattggat gaaagcttaa agcaactgac 1260
agcagaatat gatcgcacac atccccctga tgctccaga ccattacctt attttactgc 1320
agcagaaatt atgggtatag gattaactca agaacaacaa gcagaagcaa gatttgcacc 1380
agctaggatg cagtgtagag catggtatct cgaggcatta ggaaaatttg ctgccataaa 1440
agctaagtct cctcgagctg tgcagttaag acaaggagct aaggaagatt attcatcctt 1500
tatagacaga ttgtttgccc aaatagatca agaacaaaat acagctgaag ttaagttata 1560
tttaaaacag tcattgagca tagctaattg taatgcagac tgtaaaaagg caatgagcca 1620
ccttaagcca gaaagtaccc tagaagaaaa gttgagagct tgtcaagaaa taggctcacc 1680
aggatataaa atgcaactct tggcagaagc tcttacaanaa gttcaagtag tgcaatcaa 1740
aggatcagga ccagtgtgtt ttaattgtaa aaaccaggga catctagcaa gacaatgtag 1800
agaagtgaan aaatgtaata aatgtggaaa acctggtcat gtagctgcca aatgttggca 1860
aggaaataga aagaattgta caagggaaga aagggataca acaattacaa aagtgggaag 1920
attgggtagg atggatagga aatattccac aatattttaa gggactattg ggaggtatct 1980
tggaatagg attaggagtg ttattattga ttttatgttt acctacattg gttgattgta 2040
taagaaattg tatccacaag atactaggat acacagtaat tgcaatgcct gaagtagaag 2100
gagaagaat acaaccacaa atggaattga ggagaaatgg taggcaatgt ggcattgtct 2160
aaaaagagga ggaatgatga agtatctcag acttatttta taaggagatg actgtgtctg 2220
gttcttccct ttgaggaagg tatgtcatat gaaatccatt cgaaatcaaa ttcctcctct 2280
gtctgccccaa tccttccaac cccctatggt ggtatggctg acacagaaaa tgtctgtctc 2340
tgtatgggac atttgccctt cttctccaaa tataagacag gatgaggcct agcttttgtc 2400
gctccaaagt tttaaaagaa cacattgcac ggcatttagg gactctaaag ggtggaggag 2460
gaatgaggga attgcatcat gccaaaggctg gtcctcatcc atcactgctt ccaggGCCca 2520
gagtggcttc caggagggtat tcttacaagg gaagcccgat ctgtagctaa cactcagagc 2580
ccattttcct gcgttaaccc ctcggacact caatacagg agtaacatga tcagtGacct 2640
gggggagctg gccaaactgc gggacctgcc caagctgagg gccttgggtg tgctggacaa 2700
ccctctgccc gatgagactg actaccgcca ggaggccctg gtgcagatgg cacacctaga 2760
gcgcctagac aaagagtact atgaggacga ggaccgggca gaagctgagg agatccgaca 2820
gaggctgaag gaggaacagg agcaagaact cgaccgggac caagacatgg aaccgtacct 2880
cccGCCaact tagtggtctc tctagcctgc agggacagta aaggtgatgg caggaaggca 2940
gcccccgag gtcaaaggct gggcacgCGg gaggagaggc cagagtcaga ggctcggggt 3000
atctcagata tgaaggaaaag atgagagagg ctcaggaaga ggtaagaaaa gacacaagag 3060
accagagaag ggagaagaat tagagaggga ggcagaggac cgctgtctct acagacatag 3120
ctggtataga ctgggaggaa gggatgaacc ctgagcgcat gaagggaagg aggtggctgg 3180
tggtatatgg aggatgtagc tgggCCaggg aaaagatcct gcactaaaaa tctgaagcta 3240
aaaataacag gacacggggT ggagggcgca aaggagggca gattgaggca gagagactga 3300

```


gaggcctggg gatgtgggca ttccggtagg gcacacagtt cacttgtcct ctctttttcc 3360
aggaggccar agatgctgac ctcaagaact cataataccc cagtggggac caccgcattc 3420
atagccctgt tacaagaagt gggagatggt cctttttgtc ccagactgga aatccattac 3480
atcccagggc tcaggttctg tgggtggcat ctctgtgtgg cttgttctgt gggcctacct 3540
aaagtccaa gcacagctct caagcagatc cgaggcgact aagatgctag taggggtgt 3600
ctggagagaa gagccgagga ggtgggctgt gatggatcag ttcagctttc aaataaaaag 3660
gcgtttttat attctgtgtc gagttcgtga acccctgtgg tgggcttctc catctgtctg 3720
ggtagtagcc tgccactata ctggaataag gggacgcctg cttccctcga gttggctgga 3780
caaggttatg agcatccgtg tacttatggg gttgccagct tggctcctgga tcgcccgggc 3840
ccttccccca cccgttcggg tccccaccac caccgcgct cgtaçgtgcg tctccgcctg 3900
cagctcttga ctcatcgggg ccccggggtc acatgcgctc gctcggctct ataggcgccg 3960
ccccctgcc acccccgcc cgcgctggga ccgcagccg ccgccaactc tgctctctct 4020
gcgcgcgcgc cgtcaccacc gccaccgcca ccgctgagt ctgcagtcct cgaggaactg 4080
aaaaaccaga aagttaactg gtaagtttag tctttttgtc ttttatttca ggtcccggt 4140
ccggtgggtg tgcaaatcaa agaactgtc ctgagtgat gttgccttta cttctaggcc 4200
tgtacggaag tgttacttct gctctaaaag ctgcggaatt gtacccgcgg ccaagctaag 4260
cttgatatcg aattccggat gagcctctgt gaactactaa ggtgggaggg ggctatacgc 4320
agaggagaat gtcagatgct cagctcggtc ccctccgct gacgctcctc tctgtctcag 4380
ccaggactgg tttctgtaa aaacagcagg agctgtggca gcggcgaaag gaagcggctg 4440
aggcgcttgg aaccgaaaa gtctcggtgc tctcgctac ctgcacagc cgtgcccgc 4500
cgccgctcag taccatggac agcagcgtg cccccacgaa cgccagcaat tgcactgatg 4560
ccttggcgta ctcaagttgc tccccagcac ccagccccg ttcctgggtc aacttgtccc 4620
acttagatgg caacctgtcc gaöccatgcg gtccgaaccg caccgacctg ggccggagag 4680
acagcctgtg ccctccgacc ggcagtcctt ccagatcac ggccatcacg atcatggccc 4740
tctactccat cgtgtgcgtg gtggggctct tcggaaactt cctggctcatg tatgtgattg 4800
tcagatacac caagatgaag actgccacca acatctacat tttcaacctt gctctggcag 4860
atgccttagc caccagtacc ctgcccctcc agagtgtgaa ttacctaaty ggaacatggc 4920
catttggaaac catcctttgc aagatagtga tctccataga ttactataac atgttcacca 4980
gcataatcac cctctgcacc atgagtgttg atcgatacat tgcagtctgc caccctgtca 5040
aggccttaga tttccgtact ccccgaaatg ccaaaattat caatgtctgc aactggatcc 5100
tctcttcagc cattggtctt cctgtaatgt tcatggctac aacaaaatac aggcaagggt 5160
ccatagattg tacactaaca ttctctcatc caacctggta ctgggaaaaac ctgctgaaga 5220
tctgtgtttt catcttcgcc ttcattatgc cagtgtcat cattaccgtg tgctatggac 5280
tgatgatctt gcgcctcaag agtgtccgca tgctctctgg ctccaaagaa aaggacagga 5340
atcttogaag gatcaccagg atggtgctga tgggtgggac tgtgtctatc gtctgtctgc 5400
ctcccattca catctacgtc atcattaaag ccttgggtac aatcccagaa actacgttcc 5460
agactgtttc ttggcacttc tgcattgtct taggttacac aaacagctgc ctcaaccag 5520
tcctttatgc atttctggat gaaaacttca aacgatgctt cagagagttc tgtatcccaa 5580
cctcttccaa cattagcaaa caaaactcca ctogaattcg tcagaacatc tagagaccac 5640
ccctccacgg ccaatacagt ggatagaact aatcatcagc tagaaaatct ggaagcagaa 5700
actgctccgt tgccctagag ctgctgact agcctcgact gtgccttcta gttgccagcc 5760
atctgttgtt tgcccctccc ccgtgccttc cttgacctg gaaggtgcca ctcccactgt 5820
cctttcctaa taaaatgagg aaattgcac ctagtgtctg agtaggtgtc attctattct 5880
gggggggtgg gtggggcagg acagcaaggg ggaggattgg gaagacaata gcaggcatgc 5940
tggggtaaaa aagaaaaaag ggtggactgg gatgagtatt ggaaccctga agaaatagaa 6000
agaatgctta tggactaggg actgtttacg aacaaatgat aaaaggaaat agctgagcat 6060
gactcatagt taaagcgcta gcagctgcct aaccgcaaaa ccacatccta tggaaagctt 6120
gctaatacag tataagttgt tccattgtaa gagtatataa ccagtgtctt gtgaaacttc 6180
gaggagtctc tttgttgagg acttttgagt tctcccttga ggctcccaca gatacaataa 6240
atatttgaga ttgaaccctg tcgagtatct gtgtaatctt ttttacctgt gaggtctcgg 6300
aatccgggccc gagaacttcg cagcgccgcg tcatgaccga ccaagcgacg cccaacctgc 6360
catcacgaga tttcgattcc accgcccgcct tctatgaaag gttgggcttc ggaatcgctt 6420
tccgggacgc cggctggatg atcctccagc gcggggatct catgctggag ttcttcgccc 6480
accccaactt gtttattgca gcttataatg gttacaaata aagcaatagc atcaciaatt 6540
tcacaaataa agcatttttt tcaactgcatt ctagtgtgtg ttgttccaaa ctcatcaatg 6600
tatcttatca tgtctggatc ccgtcgacct cgagagcttg gcgtaatcat ggtcatagct 6660
gtttcctgtg tgaattgtt atccgctcac aattccacac aacatacagc ccggaagcat 6720
aaagtgtaaa gcctgggggt cctaattgag gagctaaact acattaattg cgttgcgctc 6780
actgcccgtt ttccagtcg gaaacctgtc gtgccagctg cattaatgaa tcggccaacg 6840
cgcggggaga ggcggtttgc gtattgggag ctcttccgct tcctcgctca ctgactcgct 6900
gcgctcggtc gttcggtgc ggcgagcggt atcagctcac tcaaaggcgg taatacggtt 6960

```

atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc 7020
caggaaccgt aaaaaggccg cgttgctggc gtttttccat aggtccgcc cccctgacga 7080
gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac tataaagata 7140
ccaggcggtt cccctggaa gctccctcgt gcgctctcct gttccgaccc tgccgcttac 7200
cggatacctg tccgccttcc tcccttcggg aagcgtggcg ctttctcaat gctcacgctg 7260
taggtatctc agttcggtgt aggtcgttcg ctccaagctg ggctgtgtgc acgaaccccc 7320
cgttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca acccggttaag 7380
acacgactta tcgcactgg cagcagccac tggtaacagg attagcagag cgaggatagt 7440
aggcgggtgt acagagttct tgaagtgggt gcctaactac ggctacacta gaaggacagt 7500
atgttggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg 7560
atccggcaaaa caaacaccg ctggtagcgg tgggtttttt gtttgcaagc agcagattac 7620
gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacggggg ctgacgctca 7680
gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac 7740
ctagatcctt ttaaatataa aatgaagttt taaatcaatc taaagtatat atgagtaaac 7800
ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt 7860
tcgttcatcc atagtgtcct gactccccgt cgtgtagata actacgatac gggagggctt 7920
accatctggc cccagtgtg caatgatacc gcgagacca cgctcacccg ctccagattt 7980
atcagcaata aaccagccag ccggaaggcg cgagcgaga agtggctctg caactttatc 8040
cgctccatc cagtctatta attgttgccg ggaagctaga gtaagttagt cgccagttaa 8100
tagtttgccg aacgttggtt ccattgtctac aggcacgtg gtgtcacgct cgtcgttttg 8160
tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat ccccatgtt 8220
gtgcaaaaaa gcggttagct ccttcgggtc tccgatcgtt gtcagaagta agttggccgc 8280
agtgttatca ctcattggtt tggcagcact gcataattct cttactgtca tgccatccgt 8340
aagatgcttt tctgtgactg gtgagtactc acccaagtca ttctgagaat agtgtatgct 8400
gcgaccgagt tgctcttgcc cggcgtcaat acgggataat accgcgccac atagcagaac 8460
tttaaaagtg ctcattcttg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc 8520
gctgttgaga tccagttcga tgaacccac tcgtgcaccc aactgatctt cagcatcttt 8580
tactttcacc agcgtttctg ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg 8640
aataagggcg acacggaaat gttgaatact catactcttc ctttttcaat attattgaag 8700
catttatcag ggttattgtc tcatgagcgg atacatattt gaatgtattt agaaaaataa 8760
acaaaatagg gttccgcgca catttcccg aaaagtgcc cctgacgtcg acggatcggg 8820
agatctcccg atccctatg gtcgactctc agtacaatct gctctgatgc cgcatagtta 8880
agccagtatc tgctccctgc ttgtgtgttg gaggtcgtg agtagtgccg gagcaaaatt 8940
taagctacaa caaggcaagg cttgaccgac aattgcatga agaactctgt taggggttagg 9000
cgttttgccg tgcttcgcga tgtacggggc agatatacgc gttgacattg attattgact 9060
agttattaat agtaatacat tacgggggtc ttagttcata gcccatatat ggagttccgc 9120
gttacataac ttacggtaaa tggcccgcct ggctgaccgc ccaacgaccc ccgcccattg 9180
acgtcaataa tgacgtatgt tcccatagta acgccaatag ggactttcca ttgacgtcaa 9240
tgggtggact atttacggta aactgccac ttggcagtac atcaagtgt tcatatgcc 9300
agtacgcccc ctattgacgt caatgacggg aaatggcccg cctggcatta tgcccagtac 9360
atgaccttat gggactttcc tacttggcag tacatctacg tattagtcac cgctattacc 9420
atggtg

```

<210> 8
 <211> 12745
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:/note =
 synthetic construct

```

<400> 8
atgcggtttt ggcagtacat caatgggcgt ggatagcggg ttgactcacg gggatttcca 60
agtctccacc ccattgacgt caatgggagt ttgttttggc accaaaatca acgggacttt 120
ccaaaatgtc gtaacaactc cgccccattg acgcaaatgg gcggtaggcg tgtacgggtg 180
gaggtctata taagcagagc tctgtgaaac ttcgaggagt ctctttgttg aggacttttg 240
agttctccct tgaggctccc acagatacaa taaatatttg agattgaacc ctgtcagata 300
tctgtgtaat cttttttacc tgtgaggtct cggaatccgg gccgagaact tcgcagttgg 360

```

cgccgaaca	gggacttgat	tgagagtgat	tgaggaagtg	aagctagagc	aatagaaagc	420
tgtaagcag	aactcctgct	gacctaaata	gggaagcagt	agcagacgct	gctaacagt	480
agtatctcta	gtgaagcggg	ctcgagctca	taatcaagtc	attgtttaa	ggccagata	540
aattacatct	ggtgactcct	cgcggacctt	caagccagga	gattcgccga	gggacagtca	600
acaaggtagg	agagattcta	cagcaacatg	gggaatggac	agggcgaga	ttggaaaatg	660
gccattaaga	gatgtagtaa	tggtgctgta	ggagtagggg	ggaagagtaa	aaaatttgga	720
gaaggggaatt	tcagatgggc	cattagaatg	gctaattgat	ctacaggacg	agaacctggt	780
gatataccag	agactttaga	tcaactaagg	ttggttattt	gcgatttaca	agaaagaaga	840
gaaaaatttg	gatctagcaa	agaaattgat	atggcaattg	tgacattaaa	agtctttgcg	900
gtagcaggac	ttttaaatat	gacgggtgtc	tactgctgct	gcagctgaaa	atatgtattc	960
tcaaatggga	ttagacacta	ggccatctat	gaaagaagca	ggtggaaaag	aggaaggccc	1020
tccacaggca	tatcctattc	aaacagtaaa	tggagtacca	caatatgtag	cacttgaccc	1080
aaaaatggtg	tccattttta	tggaaaaggc	aagagaagga	ctaggagggtg	aggaagttca	1140
actatggttt	actgccttct	ctgcaaattt	aacacctact	gacatggcca	cattaataat	1200
ggccgcacca	gggtgcgctg	cgataaaaga	aatattggat	gaaagcttaa	agcaactgac	1260
agcagaatat	gatcgcacac	atccccctga	tgctcccaga	ccattaccct	attttactgc	1320
agcagaaatt	atgggtatag	gattaactca	agaacaacaa	gcagaagcaa	gatttgcacc	1380
agctaggatg	cagtgtagag	catggtatct	cgaggcatta	ggaaaattgg	ctgccataaa	1440
agctaagtct	cctcgagctg	tgagtttaag	acaaggagct	aagggaagatt	attcatcctt	1500
tatagacaga	ttgtttgccc	aaatagatca	agaacaaaaat	acagctgaag	ttaagttata	1560
tttaaaacag	tcattgagca	tagctaattgc	taatgcagac	tgtaaaaagg	caatgagcca	1620
ccttaagcca	gaaagtaacc	tagaagaaaa	gttgagagct	tgtcaagaaa	taggctcacc	1680
aggatataaa	atgcaactct	tggcagaagc	tcttacaana	gttcaagtag	tgcaatcaaa	1740
aggatcagga	ccagtgtgtt	tttaattgtaa	aaaaccagga	catctagcaa	gacaatgtag	1800
agaagtgaag	aaatgtaata	aatgtggaaa	acctgggtcat	gtagctgcca	aatgttggca	1860
aggaaataga	aagaattgta	caaggggaaga	aagggatata	acaattacaa	aagtgggaag	1920
attgggtagg	atggatagga	aatattccac	aatattttaa	gggactattg	ggaggtatct	1980
tgggaatagg	attaggagt	ttattattga	ttttatgttt	acctacattg	gttgattgta	2040
taagaaattg	taccacaag	atactaggat	acacagtaat	tgcaatgcct	gaagtgaag	2100
gagaagaaat	acaaccacaa	atggaattga	ggagaaatgg	taggcaatgt	ggcatgtctg	2160
aaaaagagga	ggaatgatga	agtatctcag	acttatttta	taaggagat	actgtgctga	2220
gttcttccct	ttgaggaagg	tatgtcatat	gaatccattt	cgaatcaaat	caaactaata	2280
aagtatgtat	tgtaaggtaa	aaagaaaaga	caaagaagaa	gaagaaagaa	gaaagccttc	2340
agtacattta	tattggctca	tgtccaatat	gaccgccatg	ttgacattga	ttattgacta	2400
gttattaata	gtaatcaatt	acgggggtcat	tagttcatag	cccatatatg	gagttccgcg	2460
ttacataact	tacggttaatt	ggccgcctg	ctgaccgccc	aacgaccccc	gcccattgac	2520
gtcaataatg	acgtatgttc	ccatagtaac	actttccatt	gacgtcaatg	2580	
ggtggagtat	ttacggtaaa	ctgcccactt	ggcagtagat	caagtgtatc	atatgccaa	2640
tccggccccc	tattgacgtc	aatgacggta	aatggccgcg	ctggcattat	gcccagtaca	2700
tgaccttacg	ggactttggt	acttggcagt	acatctacgt	attagtcac	gctattacca	2760
tggtgatgag	gttttggcag	tacaccaatg	ggcgtggata	gcggtttgac	tcacggggat	2820
ttccaagtct	ccacccattt	gacgtcaatg	ggagtttggt	ttggcaccaa	aatcaacggg	2880
actttccaaa	atgtcgtaat	aacccgcgcc	cgttgacgca	aatgggcggg	aggcgtgtac	2940
ggtgggaggt	ctatataagc	agagctcggt	tactgaaccg	tcagatcgcc	tggagacgcc	3000
atccacgctg	ttttgacctc	catagaagac	accgggaccg	atccagcctc	cgcgccggg	3060
aacggtgcat	tggaaacgcg	attccccgtg	ccaagagtga	cgtaagtacc	gcctatagac	3120
tctataggca	cacccttttg	gctcttatgc	atgctatact	gtttttggct	tggggcctat	3180
acacccccgc	tccttatgct	ataggtgatg	gtatagctta	gcctatagg	gtgggttatt	3240
gaccattatt	gaccactccc	ctattgggtga	cgatactttc	cattactaat	ccataacatg	3300
gctctttgce	acaactatct	ctattggcta	tatgccaata	ctctgtcctt	cagagactga	3360
cacggactct	gtatttttac	aggatgggg	cccattttat	atttacaana	tcacatatac	3420
aacaacgcg	tccccgtg	cgcagttttt	tattaaacat	agcgtgggat	ctccacgcga	3480
atctcggtta	cgtgttccg	acatgggctc	ttctccggta	gcggcgagag	ttccacatcc	3540
gagccctggt	cccatgcctc	cagcggtcca	tggtcgctcg	gcagctcctt	gctcctaaca	3600
gtggaggcca	gacttaggca	cagcacaatg	cccaccacca	ccagtgtgcc	gcacaaggcc	3660
gtggcggtag	ggtatgtgtc	tgaaaatgag	ctcgagagatt	gggctcgac	cgtgacgcag	3720
atggaagact	taaggcagcg	gcagaagaag	atgcaggcag	ctgagttggt	gtattctgat	3780
aagagtcaga	ggttaactccc	gttgcggttc	tgtaaacgg	ggagggcagt	gtagtctgag	3840
cagtactcgt	tgctgccgcg	cgcgccacca	gacataatag	ctgacagact	aacagactgt	3900
tcctttccat	gggtcttttc	tgcagtcacc	gtcgtcgaag	cttatgacca	tgattacgga	3960
ttcactggcc	gtcgttttac	aacgtcgtga	ctgggaaaac	cctggcggtta	cccaacttaa	4020

tcgccttgca	gcacatcccc	ctttcgccag	ctggcgtaat	agcgaagagg	cccgaccga	4080
tcgcctttcc	caacagttgc	gcagcctgaa	tggcgaatgg	cgctttgcct	ggtttccggc	4140
accagaagcg	gtgccggaaa	gctggctgga	gtgcgatctt	cctgaggccg	atactgtcgt	4200
cgtccctca	aactggcaga	tgcacggtta	cgatgcgccc	atctacacca	acgtaaccta	4260
tcccattacg	gtcaatccgc	cgtttggtcc	cacggagaat	ccgacgggtt	gttactcgct	4320
cacatttaat	gttgatgaaa	gctggctaca	ggaaggccag	acgcgaatta	tttttgatgg	4380
cgtaactcg	gcgtttcatc	tgtggtgcaa	cgggcgctgg	gtcggttacg	gccaggacag	4440
tcgtttgccc	tctgaatttg	acctgagcgc	atttttacgc	gccggagaaa	accgcctcgc	4500
ggtgatggtg	ctgcgttgga	gtgacggcag	ttatctggaa	gatcaggata	tgtggcggat	4560
gagcggcatt	ttccgtgacg	tctcgttgct	gcataaaccg	actacacaaa	tcagcgattt	4620
ccatgttgcc	actcgtttta	atgatgattt	cagccgcgct	gtactggagg	ctgaagtcca	4680
gatgtgcggc	gagttgcgtg	actacctacg	ggtaacagtt	tctttatggc	agggtgaaac	4740
gcaggtcgcc	agcggcaccg	cgcctttcgg	cggtgaaatt	atcgatgagc	gtggtggtta	4800
tgccgatcgc	gtcacactac	gtctgaacgt	cgaaaaccgc	aaactgtgga	gcgccgaaat	4860
ccggaatctc	tatcgtgcgg	tggttgaact	gcacaccgcc	gacggcacgc	tgattgaagc	4920
agaagcctgc	gatgtcgggt	tccgcgaggt	gcggattgaa	aatggtctgc	tgctgctgaa	4980
cggcaagccg	ttgctgattc	gaggcgtaa	ccgtcacgag	catcatcttc	tgcatggtca	5040
ggtcatggat	gagcagacga	tgggtgcagga	tatcctgctg	atgaagcaga	acaactttaa	5100
cgcctgctgc	tgttcgcatt	atccgaacca	tccgctgtgg	tacacgctgt	gcgaccgcta	5160
cggcctgtat	gtggtggatg	aagccaatat	tgaaaccac	ggcatgggtc	caatgaatcg	5220
tctgaccgat	gatccgcgct	ggctaccggc	gatgagcgaa	cgcgtaacgc	gaatggtgca	5280
gcgcgatcgt	aatcacccga	gtgtgatcat	ctggctcgtg	gggaatgaat	caggccacgg	5340
cgctaatacc	gacgcgctgt	atcgctggat	caaatctgtc	gatccttccc	gcccggctga	5400
gtatgaagcg	ggcggagccg	acaccacggc	caccgatatt	atttgcgccg	tgtacgcgcg	5460
cgtggatgaa	gaccagccct	tcccggctgt	gccgaaatgg	tccatcaaaa	aatggctttc	5520
gctacctgga	gagacgcgcc	cgtgatcct	ttgcgaatac	gcccacgcga	tgggtaacag	5580
tcttgccggt	ttcgttaaat	actggcaggc	gtttcgtcag	tatccccgtt	tacagggcgg	5640
cttcgtctgg	gactgggtgg	atcagtcgct	gattaaatat	gatgaaaacg	gcaaccctgt	5700
gtcggcttac	ggcggtgatt	ttggcgatac	gccgaacgat	cgccagttct	gtatgaacgg	5760
tctggtcttt	gccgacgcga	cgcgcaccc	agcgtgacg	gaagcaaaac	accagcagca	5820
gtttttccag	ttccgtttat	cggggcaaac	catcgaaagt	accagcgaat	acctgttccg	5880
tcatagcgat	aacgagctcc	tgcaactggat	ggtggcgctg	gatggtaagc	cgctggcaag	5940
cggatgaagt	cctctggatg	tcgctccaca	aggtaaacag	ttgattgaac	tgctgaact	6000
accgcagccg	gagagcgccg	ggcaactctg	gctcacagta	cgcgtagtgc	aaccgaacgc	6060
gaccgcgatg	tcagaagccg	ggcacatcag	cgcctggcag	cagtggcgct	tggcggaaaa	6120
cctcagtggt	acgctccccg	cgcgtcccca	cgccatcccc	catctgacca	ccagcgaaat	6180
ggatttttgc	atcgagctgg	gtaataagcg	ttggcaattt	aaccgccagt	caggctttct	6240
ttcacagatg	tggtattggc	ataaaaaaca	ctcgtgacg	ccgctgcgcg	atcagttccg	6300
ccgtgcaccg	ctggataacg	acattggcgt	aagtgaagcg	accgcatttg	accctaacgc	6360
ctgggtcgaa	cgttggaagg	cggcgggcca	ttaccaggcc	gaagcagcgt	tgttgcatg	6420
cacggcagat	acacttgctg	atgcggtgct	gattacgacc	gctcacgcgt	ggcagcatca	6480
ggggaaaacc	ttattttatca	gccggaaaac	ctaccggatt	gatggtagt	gtcaaatggc	6540
gattaccggt	gatgttgaag	tggcgagcga	tacaccgcat	ccggcgcgga	ttggcctgaa	6600
ctgccagctg	gcgcaggtag	cagagcgggt	aaactggctc	ggattagggc	cgcaagaaaa	6660
ctatccccag	cgccttactg	cgcctgttt	tgaccgctgg	gatctgccat	tgtcagacat	6720
gtataccccg	tacgtcttcc	cgagcgaaaa	ccgtctgcgc	tgccggagcg	gcgaattgaa	6780
ttatggccca	caccagtggc	gcggcgactt	ccagttcaac	atcagccgct	acagtcaaca	6840
gcaactgatg	gaaaccagcc	atcgccatct	gctgcacgcg	gaagaaggca	catggctgaa	6900
tatcgacggt	ttccatatgg	ggattggtgg	cgacgactcc	tggagcccgt	cagtatcggc	6960
ggaattccag	ctgagcgccg	gtcgctacca	ttaccagttg	gtctggtgtc	aaaaataact	7020
cgatcgacca	gagctgagat	cctacaggag	tccagggtcg	gagagaaaa	ctctgaagag	7080
gatgatgaca	gagttagaag	atcgcttcag	gaagctattt	ggcacgactt	ctacaacggg	7140
agacagcaca	tagattctg	aagatgaacc	tcctaaaaaa	gaaaaaagg	tggactggga	7200
tgagtattgg	aaccttgaag	aaatagaaag	aatgcttatg	gactagggac	tgtttacgaa	7260
caaatgataa	aaggaaatag	ctgagcatga	ctcatagtta	aagcgctagc	agctgcctaa	7320
ccgcaaaacc	acatcctatg	gaaagcttgc	taatgacgta	taagttgttc	cattgtaaga	7380
gtatataacc	agtgtcttgt	gaaacttcga	ggagtctctt	tgttgaggac	ttttgagttc	7440
tcccttgagg	ctcccacaga	tacaataaat	atttgagatt	gaacctgtc	gagtatctgt	7500
gtaatctttt	ttacctgtga	ggtctcggaa	tccgggcccga	gaacttcgca	gcggccgctc	7560
gagcatgcat	ctagagggcc	ctattctata	gtgtcaccta	aatgctagag	ctcgtgatc	7620
agcctcgact	gtgccttcta	gttgccagcc	atctgttgtt	tgccctccc	ccgtgccttc	7680

cttgaccctg	gaaggtgcc	ctcccactgt	cctttcctaa	taaaatgagg	aaattgcac	7740
gcattgtctg	agtaggtgtc	attctattct	ggggggtggg	gtggggcagg	acagcaagg	7800
ggaggattgg	gaagacaata	gcaggcatgc	tggggatgog	gtgggctcta	tggcttctga	7860
ggcggaaga	accagctggg	gctcgagggg	ggatccccac	gcgccctgta	gcggcgcat	7920
aagcgcggg	ggtgtggtgg	ttacgcgcag	cgtgaccgct	acacttgcca	gcgccctagc	7980
gcccgtcct	ttcgctttct	tcccttcctt	tctcgccacg	ttcgccggct	ttccccgtca	8040
agctctaaat	cggggcatcc	ctttagggtt	ccgatttagt	gctttacggc	acctcgaccc	8100
caaaaaactt	gattagggtg	atggttcacg	tagtgggcca	tcgccctgat	agacgggttt	8160
tcgccctttg	acgttggagt	ccacgttctt	taatagtgg	ctcttgttcc	aaactggaac	8220
aacactcaac	cctatctcgg	tctattcttt	tgatttataa	gggatttttg	ggatttcggc	8280
ctattgggta	aaaaatgagc	tgatttaaca	aaaatttaac	gcgaatttta	acaaaatatt	8340
aacgtttaca	atttaaatat	ttgcttatac	aactctcctg	tttttggggc	ttttctgatt	8400
atcaaccggg	gtgggtaccg	agctcgaatt	ctgtggaatg	tgtgtcagtt	aggggtgtgga	8460
aagtccccag	gctccccagg	caggcagaag	tatgcaaagc	atgcatctca	attagtccagc	8520
aaccagggtg	ggaaagtccc	caggctcccc	agcaggcaga	agtatgcaaa	gcacgcacct	8580
caattagtc	gcaaccatag	tcccggccct	aactccggcc	atcccggccc	taactccggc	8640
cagtcccgcc	cattctccgc	cccatggctg	actaattttt	tttatttatg	cagaggccga	8700
ggccgcctcg	gcctctgagc	tattccagaa	gtagtggaga	ggcttttttg	gaggccctagg	8760
cttttgcaaa	aagctcccg	gagcttggat	atccattttc	ggatctgatc	aagagacagg	8820
atgaggatcg	tttcgcata	ttgaacaaga	tggattgcac	gcagggttctc	cggccgcttg	8880
ggtggagagg	ctattcggct	atgactgggc	acaacagaca	atcggctgct	ctgatgccgc	8940
cgtgttccgg	ctgtcagcgc	aggggcgccc	ggttcttttt	gtcaagaccg	acctgtccgg	9000
tgccctgaat	gaactgcagg	acgaggcagc	gcggctatcg	tggctggcca	cgacgggctg	9060
tccttgccga	gctgtgctcg	acgttgtcac	tgaagcggga	agggactggc	tgctattggg	9120
cgaagtgccg	gggcaggatc	tctgtcatc	tcaccttget	cctgccgaga	aagtatccat	9180
catggctgat	gcaatgcggc	ggctgcatac	tcttgatccg	gctacctgcc	cattcgacca	9240
ccaagcgaaa	catcgcatcg	agcagcacg	tactcgatg	gaagccggct	ttgtcgatca	9300
ggatgatctg	gacgaagagc	atcaggggct	cgcgccagcc	gaactgttgc	ccaggctcaa	9360
ggcgcgcatg	ccgacggcgc	aggatctcgt	cgtgaccat	ggcgatgcct	gcttgccgaa	9420
tatcatgggtg	gaaaatggcc	gcttttctgg	attcatcgac	tgtggccggc	tgggtgtggc	9480
ggaccgctat	caggacatag	cggttggtac	ccgtgatatt	gctgaagagc	ttggcgccga	9540
atgggctgac	cgcttctcgc	tgttttacgg	tatcgccgct	cccgatctgc	agcgcatcgc	9600
cttctatcgc	cttcttgacg	agttcttctg	agcgggactc	tggggttcga	aatgaccgac	9660
caagcgacgc	ccaactgcc	atcacgagat	tctgattcca	ccgccgcctt	ctatgaaagg	9720
ttgggtctcg	gaatcgtttt	ccgggacgcc	ggctggatga	tcttcagcgc	cggggatctc	9780
atgctggagt	tcttcgcccc	ccccacttg	tatttgtag	cttataatgg	ttacaaataa	9840
agcaatagca	tcacaaat	cacaaataaa	gcattttttt	cactgcattc	tagttgtggg	9900
ttgtccaaac	tcatcaatgt	atcttatcat	gtctggatcc	cgctgacctc	gagagcttgg	9960
cgtaatcatg	gtcatagctg	tttctgtgt	gaaattgtta	tccgctcaca	attccacaca	10020
acatacggagc	cggaagcata	aagtgtaaag	cctgggggtgc	ctaagtgtg	agctaactca	10080
cattaattgc	gttgcgctca	ctgcccgtt	tccagtcggg	aaacctgtcg	tgccagctgc	10140
attaatgaat	cgcccaacgc	gcggggagag	gcgggttgcg	tattggcgcg	tcttcgcgtt	10200
cctcgctcac	tgactcgctg	cgctcggtcg	tctggctgcg	gcgagcggtg	tcagctcact	10260
caaaaggcgg	aatacgggta	tccacagaat	caggggataa	cgcaggaaag	aacatgtgag	10320
caaaaggcca	gcaaaaggcc	aggaaccgta	aaaaggccgc	ggtgtggtgg	tttttccata	10380
ggctccgccc	ccctgacgag	catcacaaaa	atcgacgctc	aagtcagagg	tggcgaaacc	10440
cgacaggact	ataaagatac	caggcggttc	cccctggaag	ctccctcggt	cgctctcctg	10500
ttccgaccct	gccgcttacc	ggatacctgt	ccgcctttct	cccttcggga	agcgtggcgc	10560
tttctcaatg	ctcacgctgt	aggtatctca	gttcgggtgta	ggtcgttcgc	tccaagctgg	10620
gctgtgtgca	cgaaccccc	gttcagcccc	accgctgcgc	cttatccggt	aactatcgtc	10680
ttgagtccaa	cccggttaaga	cacgacttat	cgccactggc	agcagccact	ggtaacagga	10740
ttagcagagc	gaggtatgta	ggcggtgcta	cagagtctct	gaagtgggtg	cctaactacg	10800
gctacactag	aaggacagta	tttggtatct	gcgctctgct	gaagccagtt	accttcggaa	10860
aaagagttgg	tagctcttga	tccggcaaac	aaaccaccgc	tggtagcggt	ggtttttttg	10920
tttgcaagca	gcagattacg	cgcagaaaaa	aaggatctca	agaagatcct	ttgatctttt	10980
ctacgggggtc	tgacgctcag	tggaaacgaa	actcacgtta	agggattttg	gtcatgagat	11040
tatcaaaaag	gatcttcacc	tagatccttt	taaatataaa	atgaagtttt	aatcaatct	11100
aaagtataata	tgagtaaaat	tggcttgaca	gttaccatg	cttaatcagt	gaggcaccta	11160
tctcagcgat	ctgtctat	cggtcatcca	tagttgcctg	actccccgtc	gtgtagataa	11220
ctacgatagc	ggagggctta	ccatctggcc	ccagtgtgct	aatgataacc	cgagaccac	11280
gctcaccggc	tccagattta	tcagcaataa	accagccagc	cgaaggggcc	gagcgagaa	11340

```

gtggctctgc aactttatcc gctccatcc agtctattaa ttgttgccgg gaagctagag 11400
taagtagttc gccagttaat agtttgcgca acgttggtgc cattgctaca ggcatcgagg 11460
tgtcacgctc gtcgttttgt atggcttcat tcagctccgg ttcccaacga tcaaggcgag 11520
ttacatgatc ccccatgttg tgcaaaaaag cggtagctc cttcggtcct ccgatcggtg 11580
tcagaagtaa gttggccgca gtgttatcac tcatggttat ggcagcactg cataattctc 11640
ttactgtcat gccatccgta agatgctttt ctgtgactgg tgagtactca accaagtcat 11700
tctgagaata gtgtatgcgg cgaccgagtt gctcttgccc ggcgtcaata cgggataata 11760
ccgcgccaca tagcagaact ttaaaagtgc tcatcattgg aaaacgttct tcggggcgaa 11820
aactctcaag gatcttaccg ctgttgagat ccagttcgat gtaaccact cgtgcacca 11880
actgatcttc agcatctttt actttcacca gcgtttctgg gtgagcaaaa acaggaaggc 11940
aaaatgccgc aaaaaaggga ataaggcgca cacggaaatg ttgaatactc atactcttcc 12000
tttttcaata ttattgaagc atttatcagg gttattgtct catgagcggg tacatatttg 12060
aatgtattta gaaaaataaa caaatagggg ttccgcgcac atttccccga aaagtgccac 12120
ctgacgtcga cggatcggga gatctcccca tccccatgg tcgactctca gtacaatctg 12180
ctctgatgcc gcatagttaa gccagtatct gctccctgct tgtgtgttgg aggtcgctga 12240
gtagtgcgcy agcaaaattt aagctacaac aaggcaaggc ttgaccgaca attgcatgaa 12300
gaatctgctt aggggttagg gttttgcgct gcttcgcgat gtacgggcca gatatacgcg 12360
ttgacattga ttattgacta gttattaata gtaatcaatt acgggggtcat tagttcatag 12420
cccatatatg gagttccgcy ttacataact tacggtaaat ggcccgcctg gctgaccgcc 12480
caacgacccc cgccattga cgtcaataat gacgtatgtt cccatagtaa cgccaatagg 12540
gactttccat tgacgtcaat ggggtggacta tttacggtaa actgcccact tggcagtaca 12600
tcaagtgtat catatgccaa gtacgcccc tattgacgct aatgacggta aatggccgc 12660
ctggcattat gccagtaga tgaccttat ggactttcct acttggcagt acatctacgt 12720
attagtcacg gctattacca tgggtg

```

<210> 9

<211> 1200

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 9

```

atggacagca ggcgtgcccc cacgaacgcc agcaattgca ctgatgcctt ggcgtactca 60
agttgctccc cagcaccag ccccggttcc tggatcaact tgtcccactt agatggcaac 120
ctgtccgacc catgcggtcc gaaccgcacc gacctgggcy ggagagacag cctgtgccct 180
ccgaccggca gtccctccat gatcacggcc atcacgatca tggccctcta ctccatcgtg 240
tgctggtgg ggctcttcgg aaacttcctg gtcattgtatg tgattgtcag atacaccaag 300
atgaagactg ccaccaacat ctacattttc aaccttgctc tggcagatgc cttagccacc 360
agtaccctgc ccttcagag tgtgaattac ctaatgggaa catggccatt tggaaaccatc 420
ctttgcaaga tagtgatctc catagattac tataacatgt tcaccagcat attcaccctc 480
tgcaccatga gtgttgatcg atacattgca gtctgccacc ctgtcaaggc cttagatttc 540
cgtactcccc gaaatgccaa aattatcaat gtctgcaact ggatcctctc ttcagccatt 600
ggctcttcctg taatgttcat ggctacaaca aaatacaggc aagggttccat agattgtaca 660
ctaacattct ctcatccaac ctggtactgg gaaaacctgc tgaagatctg tgttttcatc 720
ttcgcttca ttatgccagt gctcatcatt accgtgtgct atggactgat gatcttgcyg 780
ctcaagagtg tccgcatgct ctctggctcc aaagaaaagg acaggaatct tcgaaggatc 840
accaggatgg tgctggtggg ggtggctgtg ttcatcgtct gctggactcc cattcacatt 900
tacgtcatca ttaaagcctt gggtacaatc ccagaaacta cgttcagac tgtttcttgg 960
cactctgca ttgctctagg ttacacaac agctgcctca accagtcct ttatgcattt 1020
ctggatgaaa acttcaaacg atgcttcaga gagtctgtga tcccaacctc ttccaacatt 1080
gagcaacaaa actccactcg aattcgtcag aacactagag accaccctc caggccaat 1140
acagtggata gaactaatca tcagctagaa aatctggaag cagaaactgc tccgttgccc 1200

```

<210> 10

<211> 400

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 10

```

Met Asp Ser Ser Ala Ala Pro Thr Asn Ala Ser Asn Cys Thr Asp Ala
 1          5          10          15
Leu Ala Tyr Ser Ser Cys Ser Pro Ala Pro Ser Pro Gly Ser Trp Ile
          20          25          30
Asn Leu Ser His Leu Asp Gly Asn Leu Ser Asp Pro Cys Gly Pro Asn
          35          40          45
Arg Thr Asp Leu Gly Gly Arg Asp Ser Leu Cys Pro Pro Thr Gly Ser
          50          55          60
Pro Ser Met Ile Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile Val
          65          70          75          80
Cys Val Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile Val
          85          90          95
Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu
          100          105          110
Ala Leu Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Val
          115          120          125
Asn Tyr Leu Met Gly Thr Trp Pro Phe Gly Thr Ile Leu Cys Lys Ile
          130          135          140
Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu
          145          150          155          160
Cys Thr Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys
          165          170          175
Ala Leu Asp Phe Arg Thr Pro Arg Asn Ala Lys Ile Ile Asn Val Cys
          180          185          190
Asn Trp Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met Ala
          195          200          205
Thr Thr Lys Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe Ser
          210          215          220
His Pro Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe Ile
          225          230          235          240
Phe Ala Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu
          245          250          255
Met Ile Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu
          260          265          270
Lys Asp Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Val
          275          280          285
Ala Val Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile
          290          295          300
Lys Ala Leu Val Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp
          305          310          315          320
His Phe Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val
          325          330          335
Leu Tyr Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe
          340          345          350
Cys Ile Pro Thr Ser Ser Asn Ile Glu Gln Gln Asn Ser Thr Arg Ile
          355          360          365
Arg Gln Asn Thr Arg Asp His Pro Ser Thr Ala Asn Thr Val Asp Arg
          370          375          380
Thr Asn His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
          385          390          395          400

```

<210> 11

<211> 1986

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 11

```

tcctcctctg ctcgcccaat ccttccaacc ccctatggtg gtatggctga cacagaaaat 60
gtctgctcct gtatgggaca tttgcccctc ttctccaaat ataagacagg atgaggccta 120
gcttttctg ctccaaagtt ttaaaagaac acattgcacg gcatttaggg actctaaagg 180
gtggaggagg aatgaggga atgcatcatg ccaaggctgg tcctcatcca tcaactgttc 240
cagggccag agtggcttcc aggaggtatt cttacaaagg aagcccgatc tgtagctaac 300
actcagagcc cattttcctg cggttaacccc tcccgcacct atatacagga gtaacatgat 360
cagtgcacct ggggagctgg ccaaactgcg ggacctgcc aagctgaggg ccttgggtgct 420
gtcggacaac ccctgtgccg atgagactga ctaccgccag gaggccctgg tgcagatggc 480
acacctagag cgcctagaca aagagtacta tgaggacgag gaccgggcag aagctgagga 540
gatccgacag aggctgaagg aggaacagga gcaagaactc gaccgggacc aagacatgga 600
accgtacctc ccgccaactt agtggctcct ctgacctgca gggacagtaa aggtgatggc 660
aggaaggcag ccccggagg tcaaaggctg ggcacgcggg aggagaggcc agagtcaagg 720
gtcgcgggta cttcagatat gaaggaaaga tgagagaggc tcaggaagag gtaagaaaag 780
acacaagaga ccagagaagg gagaagaatt agagaggagg gcagaggacc gctgtctcta 840
cagacatagc tggtagagac tgggaggaag ggatgaacc tgagcgcatg aagggaagga 900
ggtggctggt ggtatatgga ggatgtagct gggccaggga aaagatcctg cactaaaaat 960
ctgaagctaa aaataacagg acacgggggtg gagaggcgaa aggagggcag attgaggcag 1020
agagactgag aggcctgggg atgtgggcat tccggtaggg cacacagttc acttgtcttc 1080
tctttttcca ggaggccara gatgctgacc tcaagaactc ataatacccc agtggggacc 1140
accgcattca tagccctgtt acaagaagtg ggagatgttc ctttttgtcc cagactggaa 1200
atccattaca tcccagggct caggttctgt ggtggtcacc tctgtgtggc ttgttctgtg 1260
ggcctacct aagtcctaag cacagctctc aagcagatcc gaggcgacta agatgctagt 1320
aggggttgc tggagagaag agccgaggag gtgggctgtg atggatcagt tcagctttca 1380
aataaaaagg cgtttttata ttctgtgtcg agttcgtgaa cccctgtggt gggcttctcc 1440
atctgtctgg gttagtacct gccactatac tggaaataagg ggacgcctgc ttccctcgag 1500
ttgctggac aaggttatga gcatccgtgt acttatgggg ttgccagctt ggtcctggat 1560
cgcccgggcc ctccccccac ccgttcgggt cccaccacc acccgcgctc gtacgtgcgt 1620
ctccgcctgc tccctctgac tcatcggggc cccgggtca catgcgctcg ctgggtcta 1680
taggcgcgc cccctgcca cccccgcgc gcgctgggag ccgcagccgc cgccactcct 1740
gctctctctg cgccgcccgc gtcaccaccg ccaccgccac cggctgagtc tgcagtcctc 1800
gaggaactga aaaaccagaa agttaactgg taagtttagt ctttttgtct tttatttcag 1860
gtcccgatc cggtggtggt gcaaatcaaa gaactgctcc tcagtggatg ttgcctttac 1920
ttctaggcct gtacggaagt gttacttctg ctctaaaagc tgcggaattg taccgcggc 1980
caagct 1986

```

<210> 12

<211> 5982

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 12

```

atgcggtttt ggcagtacat caatgggctg ggatagcggg ttgactcacg gggatttcca 60
agtctccacc ccattgacgt caatgggagt ttgttttggc accaaaatca acgggacttt 120
ccaaaatgtc gtaacaactc cgccccattg acgcaaatgg gcggtaggcg tgtacgggtg 180
gaggtctata taagcagagc tctgtgaaac ttcgaggagt ctctttgttg aggacttttg 240
agttctccct tgaggctccc acagatacaa taaatatttg agattgaacc ctgtcgagta 300
tctgtgtaat cttttttacc tgtgaggtct cggaatccgg gccgagaact tcgcagtttg 360
cgcccgaaca gggacttgat tgagagtgat tgaggaaagt aagctagagc aatagaaagc 420

```


tgtaagcag	aactcctgct	gacctaaata	gggaagcagt	agcagacgct	gctaacagt	480
agtatctcta	gtgaagcggg	ctcgagctca	taatcaagtc	attgtttaaa	ggcccagata	540
aattacatct	ggtgactctt	cgcgacacct	caagccagga	gattcgccga	gggacagtca	600
acaaggtagg	agagattcta	cagcaacatg	gggaatggac	aggggcgaga	ttggaaaatg	660
gccattaaga	gatgtagtaa	tggtgctgta	ggagtagggg	ggaagagtaa	aaaatttggg	720
gaaggggaatt	tcagatgggc	cattagaatg	gctaattgtat	ctacaggacg	agaacctggg	780
gatataccag	agactttaga	tcaactaagg	ttggttattt	gcgattttaca	agaaagaaga	840
gaaaaatttg	gatctagcaa	agaaattgat	atggcaattg	tgacattaaa	agtctttgcg	900
gtagcaggac	ttttaaatat	gacgggtgtc	tactgctgct	gcagctgaaa	atatgtattc	960
tcaaatggga	ttagacacta	ggccatctat	gaaagaagca	ggtggaaaag	aggaaggccc	1020
tccacaggca	tatcctattc	aaacagtaaa	tggagtacca	caatatgtag	cacttgacct	1080
aaaaatgggtg	tccattttta	tggaaaaggc	aagagaagga	ctaggagggtg	aggaagttca	1140
actatggttt	actgccttct	ctgcaaattt	aacacctact	gacatggcca	cattaataat	1200
ggcgcaccca	gggtgcgctg	cagataaaga	aattatggat	gaaagcttaa	agcaactgac	1260
agcagaatat	gatcgacac	atccccctga	tgctcccaga	ccattaccct	attttactgc	1320
agcagaaatt	atgggtatag	gattaactca	agaacaacaa	gcagaagcaa	gatttgcacc	1380
agctaggatg	cagtgtagag	catggtatct	cgaggcatta	ggaaaattgg	ctgccataaa	1440
agctaagtct	cctcgagctg	tgagtttaag	acaaggagct	aaggaagatt	attcatcctt	1500
tatagacaga	ttgtttgccc	aaatagatca	agaacaaaaat	acagctgaag	ttaagttata	1560
tttaaaacag	tcattgagca	tagctaattg	taatgcagac	tgtaaaaagg	caatgagcca	1620
ccttaagcca	gaaagtaccc	tagaagaaaa	gttgagagct	tgtaagaaa	taggctcacc	1680
aggataataa	atgcgaactct	tggcagaagc	gtttacaaaa	gttcaagtag	tgcaatcaaa	1740
aggatcagga	ccagtggtgtt	ttaattgttaa	aaaaccagga	catctagcaa	gacaatgtag	1800
agaagtgaag	aatgttaata	aatgtggaaa	acctggtcat	gtagctgcca	aatgttggca	1860
aggaaataga	aagaattgta	caagggaaga	aagggatata	acaattacaa	aagtgggaag	1920
attgggtagg	atggatagga	aatattccac	aatattttaa	gggactattg	ggaggtatct	1980
tgggaatagg	attaggagtg	ttattattga	ttttatgttt	acctacattg	gttgattgta	2040
taagaaattg	tatccacaag	atactaggat	acacagtaat	tgcaatgcct	gaagtagaag	2100
gagaagaata	acaaccacaa	atggaattga	ggagaaatgg	taggcaatgt	ggcatgtctg	2160
aaaaagagga	ggaatgatga	agtatctcag	acttatttta	taaggagagat	actgtgctga	2220
gttcttccct	ttgaggaagg	tatgtcatat	gaatccattt	cgaaatcaaa	ttagagctcg	2280
ctgatcagcc	tcgactgtgc	cttctagtgt	ccagccatct	gttgtttgcc	cctccccctg	2340
gccttccttg	accctggaag	gtgccactcc	cactgtcctt	tcctaataaa	atgaggaaat	2400
tgcatcgcat	tgtctgagta	ggtgtcattc	tattctgggg	ggtgggggtg	ggcaggacag	2460
caagggggag	gattgggaag	acaatagcag	gcatgctggg	gtaaaaaga	aaaaaggggtg	2520
gactgggatg	agtattggaa	ccctgaagaa	atagaaagaa	tgcttatgga	ctagggactg	2580
ttacagataa	atgataaaa	ggaaatagct	catagttaaa	gcgctagcga	agttgttcca	2640
ctgcctaacc	gcaaaaaccac	atcctatgga	aagcttgcta	atgacgtata	agttgttcca	2700
ttgtaagagt	atataaccag	tgctttgtga	aacttcgagg	agtctctttg	ttgaggactt	2760
ttgagttctc	ccttgagggt	cccacagata	caataaatat	ttgagattga	accctgtcga	2820
gtatctgtgt	aatctttttt	acctgtgagg	tctcggaatc	cgggccgaga	acttcgcagc	2880
ggcgcgtcat	gaccgaccaa	gcgacgcca	acctgccatc	acgagatttc	gattccaccg	2940
ccgccttcta	tgaagggttg	ggcttcggaa	tcgttttccg	ggacgcgggc	tgatgatccc	3000
tccagcgcgg	ggtatctcatg	ctggagttct	tcgcccaccc	caacttggtt	attgcagctt	3060
ataatggtta	caaataaagc	aatagcatca	caaatttcac	aaataaagca	tttttttcac	3120
tgcatcttag	ttgtggtttg	tccaaactca	tcaatgtatc	ttatcatgtc	tggtatcccg	3180
cgacctcgag	agcttgccgt	aatcatgggtc	atagctgttt	cctgtgtgaa	attgttatcc	3240
gctcacaatt	ccacacaaca	tacgagccgg	aagcataaag	tgtaaaagcct	ggggtgccta	3300
atgagtgaag	taactcacat	taattgcgtt	gcgctcactg	cccgttttcc	agtcgggaaa	3360
cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcgtat	3420
tgggcgtctc	tccgtcttct	cgctcactga	ctcgctgcgc	tcggtcgttc	ggctgcggcg	3480
agcggatata	gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataacgc	3540
aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgtt	3600
gctggcggtt	ttccatagga	tccgcccccc	tgacgagcat	cacaaaaaatc	gacgctcaag	3660
tcagaggtgg	cgaaacccga	caggactata	aagataccag	gcgtttcccc	ctggaagctc	3720
cctcgtgcgc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	3780
ttcgggaagc	gtggcgcttt	ctcaatgctc	acgctgtagg	tatctcagtt	cggtgtaggt	3840
cgttcgcctc	aagctgggct	gtgtgcacga	acccccctgt	cagccccgacc	gctgcgctt	3900
atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	3960
agccactggg	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	4020
gtggtggcct	aactacgggt	acactagaag	gacagtattt	ggtatctgcg	ctctgctgaa	4080

```

gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa ccaccgctgg 4140
tagcgggtggt ttttttggtt gcaagcagca gattacgcgc agaaaaaaag gatctcaaga 4200
agatcctttg atcttttcta cggggtctga cgctcagtg aacgaaaact cacgttaagg 4260
gatttttggtc atgagattat caaaaaggat cttcacctag atccttttaa attaaaaatg 4320
aagtttttaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt accaatgctt 4380
aatcagtgag gcacctatct cagcgatctg tctatttctg tcatccatag ttgcctgact 4440
ccccgtcgtg tagataacta cgatacggga gggcttacca tctggcccca gtgctgcaat 4500
gataccgcga gaccacgct caccggctcc agatttatca gcaataaacc agccagccgg 4560
aagggccgag cgcagaagt gtcctgcaac tttatccgcc tccatccagt ctattaattg 4620
ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg ttgttgccat 4680
tgctacaggc atcgtgggtg cacgctcgtc gtttggtatg gcttcattca gctccggttc 4740
ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg ttagctcctt 4800
cggtcctccg atcgttgtca gaagtaagt ggcgcagtg ttatcactca tggttatggc 4860
agcactgcat aattctotta ctgtcatgcc atccgtaaga tgcttttctg tgactggtga 4920
gtactcaacc aagtcattct gagaatagt tatgcggcga ccgagttgct cttgcccggc 4980
gtcaatacgg gataataccg cgccacatag cagaacttta aaagtgtca tcattggaag 5040
acgttcttcc gggcgaaaac tctcaaggat cttaccgctg ttgagatcca gttcgatgta 5100
accactcgt gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg 5160
agcaaaaaaca ggaaggcaaa atgcccga aaaggaata agggcgacac ggaaatgttg 5220
aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat 5280
gagcggatac atatttggaat gtatttagaa aaataaaca atagggttc cgcgcacatt 5340
tccccgaaaa gtgccacctg acgtcgacgg atcgggagat ctcccgatcc cctatggtcg 5400
actctcagta caatctgctc tgatgccgca tagttaagcc agtatctgct ccctgcttgt 5460
gtgttgagg tcgctgagta gtgcgcgagc aaaatttaag ctacaacaag gcaaggcttg 5520
accgacaatt gcatgaagaa tctgcttagg gttaggcgtt ttgcgctgct tcgcgatgta 5580
cgggccagat atacgcgttg acattgatta ttgactagtt ataatagta atcaattacg 5640
gggtcattag ttcatagccc atatatggag ttccgcgtta cataacttac ggtaaattggc 5700
ccgcctggct gaccgcccac cgaccccgc ccattgacgt caataatgac gtatgttccc 5760
atagtaacgc caataggac tttccattga gctcaatggg tggactattt acggtaaaact 5820
gccacttggt cagtacatca agtgtatcat atgccaagta cgccccctat tgacgtcaat 5880
gacggtaaat ggccgcctg gcattatgcc cagtacatga ccttatggga ctttctact 5940
tggcagtaca tctacgtatt agtcacgct attaccatg tg 5982

```

<210> 13

<211> 13361

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 13

```

tggcagtaca tcaagtgtat catatgccaa gtacgcccc tattgacgtc aatgacggta 60
aatggccgc ctggcattat gcccagtaca tgaccttat ggactttcct acttggcagt 120
acatctacgt attagtcac gctattacca tgggtgatgc gttttggcag tacatcaatg 180
ggcgtggata gcggtttgac tcacggggat ttccaagtct ccacccatt gacgtcaatg 240
ggagtttggt ttggcaccaa aatcaacggg actttccaaa atgtcgtaac aactccgcc 300
cattgacgca aatgggcggg aggcgtgtac ggtgggaggt ctatataagc agagctctct 360
ggctaactag agaaccact gcttaactgg cttatcgaaa ttaatacgac tcaactatag 420
gagacccaag cttggtaccg agctcggatc cactagtaac ggccgccagt gtgctggaat 480
tctgcagata tccatcacac tggcggccca taatcaagtc attgtttaaa ggccagata 540
aattacatct ggtgactctt cgcggacctt caagccagga gattcgccga gggacagtca 600
acaaggtagg agagattcta cagcaacatg gggaaatggac aggggcgaga ttggaaaatg 660
gccattaaga gatgtagtaa tgttgctgta ggagtagggg ggaagagtaa aaaatttga 720
gaaggaacgt tcagattggc cattagaatg gctaagtgt ctacaggacg agaacctgg 780
gataataacc agactttaga tcaactaagg ttggttattt gcgatttaca agaaagaaga 840
gaaaaatttg gatctagcaa agaaattgat atggcaattg tgacattaaa agtctttgcg 900
gtagcaggac ttttaaatat gacggtgtct actgctgctg cagctgaaaa tatgtattct 960

```

caa atgggat	tagacactag	gccatctatg	aaagaagcag	gtggaaaaga	ggaaggccct	1020
ccacagggcat	atcctatttca	aacagtaaat	ggagtaccac	aatatgtagc	acttgaccca	1080
aaa atgggtgt	ccattttttat	ggaaaaggca	agagaaggac	taggaggtga	ggaagttaa	1140
ctatggtttta	ctgcctttctc	tgcaaattta	acacctactg	acatggccac	attaataatg	1200
gccgcaccag	ggtgcgctgc	agataaagaa	atattggatg	aaagcttaaa	gcaactgaca	1260
gcagaatatg	atcgcacaca	tccccctgat	gctcccagac	cattacccta	ttttactgca	1320
gcagaaatta	tgggtatagg	attaactcaa	gaacaacaag	cagaagcaag	atttgcacca	1380
gctaggatgc	agtgtagagc	atgggtatctc	gaggcattag	gaaaattggc	tgccataaaa	1440
gctaagtctc	ctcgagctgt	gcagttaaga	caaggagcta	aggaagatta	ttcatccttt	1500
atagacagat	tgtttgcccc	aatagatcaa	gaacaaaata	cagctgaagt	taagttatat	1560
ttaaaacagt	cattgagcat	agctaagtct	aatgcagact	gtaaaaaggc	aatgagccac	1620
cttaagccag	aaagtaccct	agaagaaaag	ttgagagctt	gtcaagaaat	aggctcaacca	1680
ggatataaaa	tgcaactcct	ggcagaagct	cttacaaaag	ttcaagtagt	gcaatcaaaa	1740
ggatcaggac	cagtgtgttt	taattgtaaa	aaaccaggac	atctagcaag	acaatgtaga	1800
gaagtgaaaa	aatgtaataa	atgtgaaaaa	cctgggtcatg	tagctgccaa	atgttggcaa	1860
ggaaatagaa	agaattcggg	aaactgggaag	gcggggcgag	ctgcagcccc	agtgaataaa	1920
atgcagcaag	cagtaatgcc	atctgcacct	ccaatggagg	agaaactatt	ggatttataa	1980
attataataa	agtaggtact	actacaacat	tagaaaagag	gccagaaata	ctcatatttg	2040
taaatggata	tcctataaaa	tttttattag	acacaggagc	agatataaca	attttaaata	2100
ggagagattt	tcaagtaaaa	aattctatag	aaaatgggaag	gcaaaatatg	attggagtag	2160
gaggaggaaa	gagaggaaca	aattatatta	atgtacattt	agagattaga	gatgaaaatt	2220
ataagacaca	atgtatatatt	ggtaatgttt	gtgtccttaga	agataactca	ttaatacaac	2280
cattattagg	gagagataat	atgattaaat	tcaattattag	gttagtaatg	gctcaaattt	2340
ctgataagat	tcagtagta	aaagtaaaaa	tgaaggatcc	taataaagga	cctcaataaa	2400
aacaatggcc	attaacaaat	gaaaaaattg	aagccttaac	agaaatagta	gaaagactag	2460
aaagagaagg	gaaagtaaaa	agagcagatc	caaataatcc	atggaataca	ccagtatttg	2520
ctataaaaaa	gaaaagtggg	aaatggagaa	tgctcataga	ttttagagaa	ttaaacaaac	2580
taactgagaa	aggagcagag	gtccagtggg	gactacctca	tcctgctggg	ttacaaataa	2640
aaaaacaagt	aacagtatta	gatatagggg	atgcatattt	caccattcct	cttgatccag	2700
attatgctcc	ttatacagca	tttactttac	ctagaaaaaa	taatgcggga	ccaggaagga	2760
gatttgtgtg	gtgtagtcta	ccacaaggct	ggattttaag	tccattgata	tatcaaagta	2820
cattagataa	tataatacaa	ccttttatta	gacaaaatcc	tcaattagat	atttaccat	2880
atatggatga	catttatata	ggatcaaatt	taagtaaaaa	ggagcataaa	gaaaaggtag	2940
agaattaag	aaaattacta	ttatgggtggg	gatttgaaac	tccagaagat	aaattacagg	3000
aagaaccccc	atatacatgg	atgggttatg	aattacatcc	attaacatgg	acaatacaac	3060
agaaacagtt	agacattcca	gaacagccca	ctctaaatga	gttgcaaaaa	ttagcaggaa	3120
aaattaattg	ggctagccaa	gctattccag	acttgagtat	aaaagcatta	actaacatga	3180
tgagaggaaa	tcaaaaccta	aattcaacaa	gacaaaggac	taaagaagct	cgactggaa	3240
tacaaaaggc	taaaaaggct	atagaagaac	aggtacaact	aggatactat	gaccccagta	3300
aggagttata	tgctaaatta	agtttgggtg	gaccacatca	aataagttat	caagtataat	3360
agaaggatcc	agaaaagata	ctatgggtatg	gaaaaatgag	tagacaaaag	aaaaaggcag	3420
aaaatacatg	tgatatagcc	ttaagagcat	gctataagat	aagagaagag	tctattataa	3480
gaataggaaa	agaaccaaga	tatgaaatac	ctacttctag	agaagcctgg	gaatcaaatt	3540
taattaattc	accatatcct	aaggccccac	ctcctgaggt	agaatatatc	catgctgctt	3600
tgaatataaa	gagagcgtaa	agtatgataa	aagatgctcc	aataccagga	gcagaaacat	3660
ggtatataga	tggaggtaga	aagctaggaa	agcagcctat	tggaacagata	3720	
caggaaagtg	caaagtgatg	gaattagaag	gcagtaatca	gaaggcagaa	atacaagcat	3780
tattattggc	attaaaagca	ggatcagagg	agatgaatat	tataacagat	tcacaatatg	3840
ttataaatat	tattcttcaa	caaccagata	tgatggaggg	aatctggcaa	gaagttttag	3900
aagaattgga	gaagaaaaca	gcaatattta	tagattgggt	cccaggacat	aaaggatttc	3960
caggaaatga	ggaagtagat	aagctttgtc	aaacaatgat	gataatagaa	ggggatggga	4020
tattagataa	aaggctcagaa	gatgcaggat	atgattttatt	agctgcaaaa	gaaatacatt	4080
tattgccagg	agaggtaaaa	gtaataccaa	caggggtaaa	gctaattgtg	cctaaaggat	4140
attggggatt	aataatagga	aaaagctcga	tagggagtaa	aggattggat	gtattaggag	4200
gggtaataga	cgaaggatat	cgaggtgaaa	ttggagtaat	aatgattaat	gtatcaagaa	4260
aatcaatcac	cttaattggaa	cgacaaaaga	tagcacaatt	aataatattg	ccttgtaaac	4320
atgaagtatt	agaacaagga	aaagtagtaa	tggattcaga	gagaggagac	aatgggttatg	4380
ggtcaacagg	agtattctcc	tcttgggttg	acagaattga	ggaagcagaa	ataaatcatg	4440
aaaaatttca	ctcagatcca	cagtacttaa	ggactgaatt	taattttacct	aaaatggtag	4500
cagaagagat	aagacgaaaa	tgcccagtat	gcagaatcag	aggagaacaa	gtgggaggac	4560
aattgaaaat	agggcctggt	atctggcaaa	tggattgcac	acactttgat	ggcaaaaataa	4620

ttcttgtggg	tatacatgtg	gaatcaggat	atatatgggc	acaaataatt	tctcaagaaa	4680
ctgctgactg	tacagttaaa	gctgtcttac	aattgttgag	tgctcataat	gttactgaat	4740
tacaaacaga	taatggacca	aattttaaaa	atcaaaagat	ggaaggagta	ctcaattaca	4800
tgggtgtgaa	acataagttt	ggatccccag	ggaacccaca	gtcacaagca	ttagttgaaa	4860
atgtaaatca	tacattaaaa	gtttggatto	ggaaatTTTT	gcctgaaaca	acctccttgg	4920
ataatgcctt	atctctcgct	gtacatagtc	tcaattttaa	aagaagaggt	aggataggag	4980
ggatggcccc	ttatgaatta	ttagcacaac	aagaatcctt	aagaatacaa	gattatTTTT	5040
ctgcaatacc	acaaaaattg	caagcacagt	ggatttatta	taaagatcaa	aaagataaga	5100
aatggaaagg	accaatgaga	gtagaatact	ggggacaggg	atcagtatta	ttaaaggatg	5160
aagagaaggg	atattttcct	atacctagga	gacacataag	gagagttcca	gaaccctgcg	5220
ctctccttga	aggggtagag	tgaagaagat	tggcaggtaa	gtagaagact	ctttgcagtg	5280
ctccaaggag	gagtaaatag	cgctatgcta	tacatatcta	ggctacctcc	ggatgaaaga	5340
gaaaagtata	aaaaagactt	caagaaaaga	ctttttgaca	cagaaacagg	atttataaag	5400
agactacgga	aagctgaagg	aataaaatgg	agctttcata	ctagagatta	ttacatagga	5460
tatgtcagag	aaatggtggc	aggatccact	acatcattaa	gtctaaggat	gtatatatat	5520
ataagtaacc	cactatggca	ttctcagtat	cgtccaggtt	tgaaaaattt	caataaggaa	5580
tggccttttg	taaatatgtg	gataaaaaca	ggatttatgt	gggatgatat	tgaaaaacaa	5640
aatatttgta	taggaggaga	agtttcacca	ggatggggac	cagggatggg	aggatatagca	5700
ataaaagctt	ttagtgttgg	cgaaagaaag	attgaggcta	ctcctgtaat	gattataaga	5760
ggagaaatag	atccaaaaaa	atggtgcgga	gattgttggg	atttaattgtg	tcttagaaac	5820
tcacctccaa	agactttaca	aagactcgct	atggtggcgt	gtggcgtgcc	ggctaagaag	5880
tggcgaggat	gctgtaatca	acgctttgtt	tctccttaca	gaacgcctgc	tgatttagag	5940
gtcattcaat	ccaagcccag	ctggaacctg	ttatggtcgg	gagaattatg	aatggaagac	6000
ataatagtat	tattcaatag	ggtcactgag	aaactagaaa	aagaattagc	tatcagaata	6060
tttgtattag	cacatcaatt	agaaagggac	aaagctatta	gattactaca	aggattatTT	6120
tggagatata	gattttaagaa	accccgagta	gattattggt	tatggttggtg	gtgttgcaaa	6180
ttctattatt	ggcagttgca	atctacatta	tcaataacta	ctgcttagaa	atatttagat	6240
taatatTTTca	tttgcaacaa	taagaatggc	agaaggattt	gcagccaata	gacaatggat	6300
aggactagaa	gaagctgaag	agttattaga	ttttgatata	gcaacacaaa	tgagtgaaga	6360
aggaccacta	aatccaggag	taaaccatt	tagggtagct	ggaataacag	aaaaagaaaa	6420
gcaaaactac	tgtaacatat	tacaacctaa	gttacaagat	ctaaggaacg	aaattcaaga	6480
ggtaaaactg	gaagaaggaa	atgcaggtaa	gtttagaaga	gcaagatttt	taaggatttc	6540
tgatgaaagt	gtattgtccc	tgggtcatgc	gttcatagga	tattgtatat	atttaggtaa	6600
tcgaaataag	ttaggatcct	taagacatga	ctttagatata	gaagcaccac	aagaagagtg	6660
ttataataat	agagagaagg	gtacaactga	caatataaaa	tatggtagac	gatgtgcct	6720
aggaacgggt	actttgtacc	tgattttatt	tataggaata	ataatatatt	cacagacaac	6780
caacgctcag	gtagtatgga	gacttccacc	attagtagtc	ccagtagaag	aatcagaaat	6840
aatttttttg	gattgttggg	caccagaaga	acccgcctgt	caggactttc	ttggggcaat	6900
gatacatcta	aaagctaaga	caaataatag	tatacgagag	ggacctacct	tggggaattg	6960
ggctagagaa	atatgggcaa	cattattcaa	aaaggctact	agacaatgta	gaagaggcag	7020
aatatggaaa	agatggaatg	agactataac	aggaccatca	ggatgtgcta	ataacacatg	7080
ttataatggt	tcagtaatat	tacctgatta	tcagtggtat	ttagatagag	tagatacttg	7140
gttacaaggg	aaaataaata	tatcattatg	tctaacagga	ggaaaaatgt	tgtacaataa	7200
agttacaaaa	caattaagct	attgtacaga	cccattacaa	atccactga	tcaattatac	7260
atttggaact	aatcaaacat	gtatgtggaa	tacttcacaa	attcaggacc	ctgaaatacc	7320
aaggccgcgg	ccctggcaac	ccatcaagaa	gctgtagaaa	aggtgactga	agccttaaag	7380
ataaacaact	taagattagt	tacattagag	catcaagtac	tagtaatagg	attaaaagta	7440
gaagctatgg	aaaaatTTTT	gtatacagct	ttcgctatgc	aagaattagg	atgtaatcaa	7500
aatcaatTTT	tctgcaaaat	ccctcctgag	ttgtggacaa	ggtataatat	gactataaat	7560
caaacaatat	ggaatcatgg	aaatataact	ttgggggaat	ggtataacca	aacaaaagat	7620
ttacaacaaa	agttttatga	aataataatg	gacatagaac	aaaataatgt	acaagggaag	7680
aaagggatac	aacaattaca	aaagtgggaa	gattgggtag	gatggatagg	aaatattcca	7740
caatatttta	agggactatt	gggaggatc	ttgggaatag	gattaggagt	gttattattg	7800
attttatggt	tacctacatt	ggttgattgt	ataagaaatt	gtatccacaa	gatactagga	7860
tacacagtaa	ttgcaatgcc	tgaagtagaa	ggagaagaaa	tacaaccaca	aatggaattg	7920
aggagaaatg	gtaggcaatg	tggcatgtct	gaaaaagagg	aggaatgatg	aagtatctca	7980
gacttatTTT	ataagggaga	tactgtgctg	agttcttccc	tttgagggaag	gtatgtcata	8040
tgaatccatt	tcgaatcaaa	tcaaactaat	aaagtatgta	ttgtaaggta	aaaggaaaag	8100
acaaagaaga	agaagaaaga	agaaagcctt	caagaggatg	atgacagagt	tagaagatcg	8160
cttcaggaag	ctattttggca	cgacttctac	aacgggagac	agcacagtag	attctgaaga	8220
tgaacctcct	aaaaaagaaa	aaagggtgga	ctgggatgag	tattggaacc	ctgaagaaat	8280

agaaagaatg	cttatggact	agggactgtt	tacgaacaaa	tgataaaagg	aaatagctga	8340
ctagagggcc	ctattctata	gtgtcaccta	aatgctagag	ctcgctgata	agcctcgact	8400
gtgccttcta	gttgccagcc	atctgttgtt	tgcctctccc	ccgtgccttc	cttgaccctg	8460
gaaggtgcca	ctcccactgt	cctttcctaa	taaaatgagg	aaattgcata	gcattgtctg	8520
agtaggtgtc	attctattct	ggggggtggg	gtggggcagg	acagcaagg	ggaggattgg	8580
gaagacaata	gcaggcatgc	tggggatgcg	gtgggctcta	tggcttctga	ggcggaaga	8640
accagctggg	gctcgagggg	ggatccccac	gcgccttgta	gcggcgcat	aagcgcgcg	8700
gggtgtggtg	ttacgcgcag	cgtgaccgct	acacttgcca	gcgccttagc	gcccgtcct	8760
ttcgctttct	tccttctcct	tctcgccacg	ttcgccggct	ttccccgtca	agctctaaat	8820
cggggcatcc	ctttagggtt	cggatttagt	gctttacggc	acctcgaccc	caaaaaactt	8880
gatttagggg	atgggttcacg	tagtgggcca	tcgccttgat	agacggtttt	tcgccttttg	8940
acgttggagt	ccacgttctt	taatagtggg	ctcttgttcc	aaactggaac	aacactcaac	9000
cctatctcgg	tctattcttt	tgatttataa	ggatttttgg	ggatttcggc	ctattggtta	9060
aaaaatgagc	tgatttaaca	aaaatttaac	gcgaatttta	acaaaatatt	aacgtttaca	9120
atttaaatat	ttgcttatac	aatcttcctg	tttttggggc	ttttctgatt	atcaaccggg	9180
gtgggtaccg	agctcgaatt	ctgtggaatg	tgtgtcagtt	aggggtgtgga	aagtccccag	9240
gctccccagg	caggcagaag	tatgcaaaag	atgcatctca	attagtcagc	aaccagggtg	9300
ggaaagtccc	caggctcccc	agcaggcaga	agtatgcaaa	gcatgcatct	caattagtca	9360
gcaaccatag	tcccgccctt	aactccgccc	atcccgcctc	taactccgcc	cagttccgcc	9420
cctatctcgg	ccctaggctg	actaattttt	ttattttatg	cagaggccga	ggccgcctcg	9480
gcctctgagc	tattccagaa	gtagtggagg	ggcttttttg	gaggccctagg	cttttgcaaa	9540
aagctcccg	gagcttggat	atccattttc	ggatctgata	aagagacagg	atgaggatcg	9600
tttcgcatga	ttgaacaaga	tggattgcac	gcaggttctc	cggccgcttg	ggtggagagg	9660
ctattcggct	atgactgggc	acaacagaca	atcggtgct	ctgatgccgc	cgtgttccgg	9720
ctgtcagcgc	aggggcgccc	ggttcttttt	gtcaagaccg	acctgtccgg	tgccctgaat	9780
gaactgcagg	acgaggcagc	gcggtatctg	tggctggcca	cgacgggctg	tccttgcgca	9840
gctgtgctcg	acgttgtcac	tgaagcggga	agggactggc	tgctattggg	cgaagtgcg	9900
gggcaggatc	tctgtgcatc	tcaccttgct	cctgccgaga	aagtatccat	catggctgat	9960
gcaatgcggc	ggctgcatac	gcttgatccg	gctacctgcc	cattcgacca	ccaagcgaaa	10020
catcgcatcg	agcgagcacg	tactcgatg	gaagccggtc	ttgtcgatca	ggatgatctg	10080
gacgaagagc	atcaggggct	cgcgccagcc	gaactgttcg	ccaggctcaa	ggcgcgcatg	10140
cccgcggcgc	aggatctcgt	cgtgaccat	ggcgatgcct	gcttgccgaa	tatcatggtg	10200
gaaaatggcc	gcttttcttg	attcatcgac	tgtggccggc	tgggtgtggc	ggaccgctat	10260
caggacatag	cgttggctac	ccgtgatatt	gctgaagagc	ttggcgcgca	atgggctgac	10320
cgttctcctg	tgtctgcatc	cccgatccgc	agcgcatcgc	cttctatcgc	cttctatcgc	10380
cttcttgacg	agttcttctg	agcgggactc	tgggggttca	aatgaccgac	caagcgacgc	10440
ccaacctgcc	atcacgagat	ttcgattcca	ccgcgcctt	ctatgaaagg	ttgggcttcg	10500
gaatcgtttt	ccgggacgcc	ggctggatga	tcctccagcg	cggggatctc	atgctggagt	10560
tcttcgcccc	cccgaacttg	tttattgcag	cttataatgg	ttacaaataa	agcaatagca	10620
tcacaaat	cacaaataaa	gcattttttt	cactgcattc	tagttgtggg	ttgtccaaac	10680
tcataaatgt	atcttatcat	gtctggatcc	cgtcgacctc	gagagcttgg	cgtaatcatg	10740
gtcatagtg	tttctgtgtg	gaaattgtta	tcctgtcaca	attccacaca	acatacagc	10800
cgcgaagcata	aatgtgaaag	cctgggggtg	cctaagtgatg	agctaactca	cattaattgc	10860
gttgcgctca	ctgcccgcct	tccagtcggg	aaacctgtcg	tgccagctgc	attaatgaat	10920
cggccaacgc	gcggggagag	gcgggttgcg	tattgggcgc	tcttcgcgtt	cctcgctcac	10980
tgactcgctg	cgtcgggtcg	ttcggtgcg	gcgagcggta	tcagctcact	caaaggcggt	11040
aatacggtta	tccacagaat	caggggataa	cgcaggaaag	aacatgtgag	caaaaggcca	11100
gcaaaaggcc	aggaaccgta	aaaaggccgc	ggtgtgtggc	tttttccata	ggctccgccc	11160
cctgacgag	catcacaaaa	atcgacgctc	aagtacagag	tggcgaaacc	cgacaggact	11220
ataaagatac	caggcgtttc	cccctggaag	ctccctcgtg	cgtctccttg	ttccgacct	11280
gccgcttacc	ggataacctg	cgcctttct	cccttcggga	agcgtggcgc	tttctcaatg	11340
ctcacgctgt	aggatatctc	gttcggtgta	ggtcggtcgc	tccaagctgg	gctgtgtgca	11400
cgaaccccc	gttcagcccg	accgctgcgc	cttatccggt	aactatcgtc	ttgagtccaa	11460
cccggtaaga	cacgacttat	cgcactggc	agcagccact	ggtaacagga	ttagcagagc	11520
gaggatgtga	ggcggtgcta	cagagtctt	gaagtgggtg	cctaactacg	gctacactag	11580
aaggacagta	tttgggtatc	gcgctctgct	gaagccagtt	accttcggaa	aaagagttgg	11640
tagctcttga	tccggcaaac	aaaccaccgc	tggtagcggt	ggtttttttg	tttgcaagca	11700
gcagatctacg	cgcagaaaaa	aaggatctca	agaagatcct	tgatctttt	ctacggggtc	11760
tgcagctcag	tgggaacgaaa	actcacgtaa	agggatcttg	gtcatgagat	tatcaaaaag	11820
gatcttcacc	tagatccttt	taaattaaaa	atgaagtgtt	aaatcaatct	aaagtatata	11880
tgagtaaact	tggtctgaca	gttaccaatg	cttaatcagt	gaggcaccta	tctcagcgat	11940

```

ctgtctatatt cggtcatcca tagttgctg actccccgtc gtgtagataa ctacgatacg 12000
ggagggttta ccattctggcc ccagtgtgc aatgataccg cgagaccac gctcaccggc 12060
tccagattta tcagcaataa accagccagc cggaggggcc gagcgagaa gtgggtcctgc 12120
aactttatcc gcctccatcc agtctattaa ttgttgccgg gaagctagag taagtagttc 12180
gccagttaat agtttgcgca acgttggtgc cattgtctaca ggcacgtgg tgtcacgctc 12240
gtcgtttggt atggcttcat tcagctccgg tccccaacga tcaaggcgag ttacatgatc 12300
ccccatgttg tgcaaaaaaag cggttagctc cttcggctct ccgatcgttg tcagaagtaa 12360
gttgccgca gtgttatcac tcatggttat ggcagcactg cataattctc ttactgtcat 12420
gccatccgta agatgctttt ctgtgactgg tgagtactca accaagtcac tctgagaata 12480
gtgtatgcgg cgaccgagtt gctcttgccc ggcgtcaata cgggataata ccgcgccaca 12540
tagcagaact ttaaaagtgc tcatcattgg aaaaacgttct tcggggcgaa aactctcaag 12600
gatcttaccg ctggtgagat ccagttcgat gtaaccact cgtgcacca actgatcttc 12660
agcatctttt actttcacca gcgtttctgg gtgagcaaaa acaggaaggc aaaatgcccgc 12720
aaaaaaggga ataaggcgca cacggaaatg ttgaatactc atactcttcc tttttcaata 12780
ttattgaagc atttatcagg gttattgtct catgagcgga tacatatttg aatgtattta 12840
gaaaaataaa caaatagggg ttccgcgcac atttccccga aaagtgcac ctgacgtcga 12900
cggatcggga gatctccgca tccccatgg tcgactctca gtacaatctg ctctgatgcc 12960
gcatagttaa gccagtatct gctccctgct tgtgtgttgg aggtcgtga gtagtgccgc 13020
agcaaaattt aagctacaac aaggcaaggc ttgaccgaca attgcatgaa gaatctgctt 13080
agggttaggc gttttgcgct gcttcgcgat gtacgggcca gatatacgcg ttgacattga 13140
ttattgacta gttattaata gtaatcaatt acggggcat tagttcatag cccatatatg 13200
gagttccgcy ttacataact tacggtaaat gggccgctg gctgaccgcc caacgacccc 13260
cgccattga cgtcaataat gacgtatgtt cccatagtaa cgccaatagg gactttccat 13320
tgacgtcaat ggggtgacta tttacggtaa actgcccact c 13361

```

<210> 14

<211> 9569

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 14

```

atgcggtttt ggcagtagat caatgggagc ggatagcggc ttgactcacg gggatttcca 60
agtctccacc ccattgacgt caatgggagc ttgttttggc accaaaatca acgggacttt 120
ccaaaatgtc gtaacaactc cgccccattg acgcaaatgg gcggtaggcg tgtacggttg 180
gaggtctata taagcagagc tctgtgaaac ttcgaggagt ctctttgttg aggacttttg 240
agttctccct tgaggtctcc acagatacaa taaatatttg agattgaacc ctgtcgagta 300
tctgtgtaat cttttttacc tgtgaggtct cggaaatccg gccgagaact tcgcagtttg 360
cgcccgaaaca gggacttgat tgagagtgat tgaggaagtg aagctagagc aatagaaaagc 420
tgtaagcag aactcctgct gacctaaata gggaagcagt agcagacgct gctaaccagt 480
agtatctcta gtgaagcgga ctcgagctca taatcaagtc attgtttaaa ggcccagata 540
aattacatct ggtgactctt cgccgacatt caagccagga gattcgccga gggacagtca 600
acaaggtagg agagattcta cagcaacatg gggaatggac aggggagaga ttggaaaatg 660
gccattaaga gatgtagtaa tgttgctgta ggagtagggg ggaagagtaa aaaatttgga 720
gaagggaatt tcagatgggc cattagaatg gctaattgtat ctacaggagc agaacctggt 780
gatataccag agactttaga tcaactaagg ttggttattt gcgatttaca agaaagaaga 840
gaaaaatttg gatctagcaa agaaattgat atggcaattg tgacattaaa agtctttgcy 900
tagcaggac ttttaaata gacgggtgtc taactgtgct gcagctgaaa atatgtattc 960
tcaaatggga tttagacacta ggccatctat taatgaagca ggtggaaaag aggaaggccc 1020
tccacaggca taccctattc aaacagtaaa tggagtacca caatatgtag cacttgaccc 1080
aaaaatggty tccattttta tggaaaaggc aagagaagga ctaggagggt aggaagttca 1140
actatggttt actgccttct ctgcaaattt aacacctact gacatggcca cattaataat 1200
ggccgcacca ggggtgcgctg cagataaaga aatattggat gaaagcttaa agcaactgac 1260
agcagaatat gatcgacac atccccctga tgctcccgca ccattaccct attttactgc 1320
agcagaatatt atgggtatag gattaactca agaacaacaa gcagaagcaa gatttgcacc 1380
agctaggatg cagtgtagag catggtatct cgaggcatta ggaaaattgg ctgccataaa 1440

```

agctaagtct	cctcgagctg	tgcagttaag	acaaggagct	aaggaagatt	attcatcctt	1500
tatagacaga	ttgtttgccc	aaatagatca	agaacaaaat	acagctgaag	ttaagttata	1560
tttaaaacag	tcattgagca	tagctaattgc	taatgcagac	tgtaaaaagg	caatgagcca	1620
ccttaagcca	gaaagtaccc	tagaagaaaa	gttgagagct	tgtcaagaaa	taggctcacc	1680
aggatataaa	atgcaactct	tggcagaagc	tcttacaaaa	gttcaagtag	tgcaatcaaa	1740
aggatcagga	ccagtggtgt	ttaattgttaa	aaaaccagga	catctagcaa	gacaatgtag	1800
agaagtgaaa	aaatgtaata	aatgtggaaa	acctgggtcat	gtagctgcca	aatgttggca	1860
aggaaataga	aagaattgta	caagggaaga	aaggggataca	acaattacaa	aagtgggaag	1920
attgggtagg	atggatagga	aatattccac	aatattttaa	gggactattg	ggaggtatct	1980
tgggaatagg	attaggagtg	ttattattga	ttttatgttt	acctacattg	gttgattgta	2040
taagaaattg	tatccacaag	atactaggat	acacagtaat	tgcaatgcct	gaagtagaag	2100
gagaagaaat	acaaccacaa	atggaattga	ggagaaatgg	taggcaatgt	ggcatgtctg	2160
aaaaagagga	ggaatgatga	agtatctcag	acttatttta	taagggagat	actgtgctga	2220
gttcttccct	ttgaggaagg	tatgtcatat	gaatccattt	cgaatcaaat	tccccagcat	2280
gctgtctatt	gtcttcccaa	tcctccccct	tgtgttctctg	ccccaccca	ccccccagaa	2340
tagaatgaca	cctactcaga	caatgcgatg	caatttcctc	attttattag	gaaaggacag	2400
tgggagtggc	accttccagg	gtcaaggaag	gcacggggga	ggggcaaaaa	acagatggct	2460
ggcaactaga	aggcacagtc	gaggctgatc	agcgagctct	gggcaacgga	gcagtttctg	2520
cttccagatt	ttctagctga	tgattagttc	tatccactgt	attggccgtg	gaggggtggg	2580
ctctagatgt	tctgacgaat	tgcagtgagg	ttttgttgct	caatgttggg	agaggttggg	2640
atacagaact	ctctgaagca	tcggttgaag	ttttcatcca	gaaatgcata	aaggactggg	2700
ttgaggcagc	tggttgtgta	acctagagca	atgcagaagt	gccaagaaac	agtctggaac	2760
gtagtttctg	ggattgtaac	caaggcttta	atgatgacgt	aaatgtgaat	gggagtcocag	2820
cagacgatga	acacagccac	caccaccagc	accatcctgg	tgatccttcg	aagattcctg	2880
tccttttctt	tggagccaga	gagcatgcgg	acactcttga	ggcgcaagat	catcagtcga	2940
tagcacacgg	taatgatgag	cactggcata	atgaaggcga	agatgaaaac	acagatcttc	3000
agcaggtttt	cccagtacca	gggttgatga	gagaatgtta	gtgtacaatc	tatggaacct	3060
tgctgttatt	ttgtttagtc	catgaacatt	acaggaagac	caatggctga	agagaggatc	3120
cagttgcaga	cattgataat	tttggcattt	cggggagtag	ggaaatctaa	ggccttgaca	3180
gggtggcaga	ctgcaatgta	tcgatcaaca	ctcattgggtg	agagggtgaa	tatgctgggtg	3240
aaatggttat	agtaattctat	ggagatcact	atccttgcaaa	ggatgggttc	aaatggccat	3300
gttcccatga	ggtaattcac	actctggaag	ggcagggtag	tggtggctaa	ggcatctgcc	3360
agagcaagg	tgaaaatgta	gatgttgggtg	gcagtcttca	tcttgggtga	tctgacaatc	3420
acatacatga	ccaggaagtt	tccgaagagc	cccaccacgc	acacgatgga	gtagaggggc	3480
atgatcgtga	tggccctgat	catggaggga	ctgccggctg	gagggcacag	gctgtctctc	3540
ccgccaggt	cgggtgcggt	cggaccgcat	gggtcggaca	ggttgccatc	taagtgggac	3600
aagttgaccc	aggaaccggg	gctgggtgct	ggggagcaac	ttgagtacgc	caaggcatca	3660
gtgcaattgc	tggcgttcgt	gggggcagcg	atgctgtcca	tggtactgac	ggccgggcgg	3720
gcaccgctgt	gcgaggtagc	caggagcacc	gagacttttc	gggttccaag	cgcctcagcc	3780
gcttcccttc	gccgctgcca	cagctcctgc	tgtttcttac	agaaaccagt	cctggctgag	3840
acagagagga	gcgtcaggcg	gaggggaccg	agctgagcat	ctgacattct	cctctgcgta	3900
tagccccctc	ccaccttagt	agttcacaga	ggctcatccg	gaattcgata	tcaagcttag	3960
cttggccgcg	ggtacaattc	cgcagctttt	agagcagaag	taacacttcc	gtacaggcct	4020
agaagtaaa	gcaacatcca	ctgaggagca	gttctttgat	ttgcaccacc	accggatccg	4080
ggacctgaaa	taaaagacaa	aaagactaaa	cttaccagtt	aactttctgg	tttttcagtt	4140
cctcgaggac	tgcagactca	cctgggtggc	gtggcggtgg	tgacggcggc	ggcgagaga	4200
gagcaggagt	ggcgggcggt	gcggtctcca	gcgcggggcg	gggttgggca	ggggggcgcg	4260
cctatagagc	cgagcgagcg	catgtgaccc	gggggccccg	atgagtcaag	agctgcaggc	4320
ggagacgcac	gtacgagcgc	gggtgggtgg	ggggaaccga	acgggtgggg	gaagggcccg	4380
ggcgatccag	gaccaagctg	gcaaccccat	aagtacacgg	atgctcataa	ccttgtccag	4440
ccaactcgag	ggaagcaggc	gtccccttat	tccagtatag	tggcaggtag	taaccagac	4500
agatggagaa	gcccaccaca	ggggttcacg	aactcgacac	agaatataaa	aacgcctttt	4560
tatttgaaag	ctgaactgat	ccatcacagc	ccacctcttc	ggctcttctc	tccagacaac	4620
ccctactagc	atcttagtcg	cctcgatctc	ccttgagagc	tgtgcttagg	acttttaggt	4680
ggcccacaga	acaagccaca	cagagatgac	caccacagaa	cctgagcctc	gggatgtaat	4740
ggatttccag	tctgggacaa	aaaggaacat	ctcccacttc	ttgtaacagg	gctatgaatg	4800
cggtgggtccc	cactggggta	ttatgagttc	ttgaggtcag	catctytggc	ctcctggaaa	4860
aagagaagac	aagtgaactg	tgtgccctac	cggaatgccc	acatccccag	gcctctcagt	4920
ctctctgcct	caatctgccc	tcctttcgcc	tctccacccc	gtgtcctgtt	attttttagct	4980
tcagattttt	agtgcaggat	cttttccttg	gcccagctac	atcctccata	taccaccagc	5040
cacctccttc	ccttcatgcg	ctcagggttc	atcccttctc	cccagtcctc	accagctatg	5100

tctgtagaga	cagcggtcct	ctgcctccct	ctctaattct	tctcccttct	ctggtctctt	5160
gtgtcttttc	ttacctcttc	ctgagcctct	ctcatctttc	cttcatactt	gagatacccg	5220
cagcctctga	ctctggcctc	tcctcccgcg	tgcccagcct	ttgacctccg	ggggctgcct	5280
tcctgccatc	acctttactg	tccttgccagg	ctagaggagc	cactaagttg	gcgggaggta	5340
cggttccatg	tcttggtccg	ggtcgagttc	ttgctcctgt	tcctccttca	gcctctgtcg	5400
gatctcctca	gcttctgccc	ggtcctcgte	ctcatagtac	tctttgtcta	ggcgctctag	5460
gtgtgccatc	tgcaccaggg	cctcctggcg	gtagtcagtc	tcacggcac	aggggttgte	5520
cagcagcacc	aaggccctca	gcttgggcag	gtcccgagc	ttggccagct	ccccagggtc	5580
actgatcatg	ttactcctgt	atatgagggtc	gggaggggtt	aacgcaggaa	aatgggctct	5640
gagtgttagc	tacagatcgg	gcttcctttg	taagaatacc	tcctggaagc	cactctgggc	5700
cctggaagca	gtgatggatg	aggaccagcc	ttggcatgat	gcaattccct	cattcctcct	5760
ccacccttta	gagtccctaa	atgcctgtca	atgtgttctt	ttaaaacttt	ggagcagcaa	5820
aagctaggcc	tcacctgtct	ttatatattg	agaagagggg	caaagtctcc	atacaggagc	5880
agacattttc	tgtgtcagcc	ataccaccat	agggggttgg	aaggattggg	cgagcagagg	5940
aggaaatgga	attgaggaga	aatggtaggc	aatgtggcat	gtctgaaaaa	gaggaggaa	6000
gatgaagtat	ctcagactta	ttttataagg	gagatactgt	gctgagttct	tccttttgag	6060
gaaggatagt	catatgaatc	catttcgaat	caaattccta	aaaaagaaaa	aagggtggac	6120
tgggatgagt	attggaaccc	tgaagaaata	gaaagaatgc	ttatggacta	gggactgttt	6180
acgaacaaat	gataaaagga	aatagctgag	catgactcat	agttaaagcg	ctagcagctg	6240
cctaaccgca	aaaccacatc	ctatggaaag	cttgctaattg	acgtataagt	tgttccattg	6300
taagagtata	taaccagtgc	tttgtgaaac	ctcgaggagt	ctctttgttg	aggacttttg	6360
agttctccct	tgaggctccc	acagatacaa	taaataattg	agattgaacc	ctgtcgagta	6420
tctgtgtaat	cttttttacc	tgtgaggtct	cggaaatccg	gccgagaact	tcgcagtgac	6480
cgaccaagcg	acgccaacc	tgccatcacg	agatttcgat	tcaccgccg	ccttctatga	6540
aagggtgggc	ttcggaatcg	ttttccggga	cgccggctgg	atgatcctcc	agcgcgggga	6600
tctcatgctg	gagttcctcg	cccaccccaa	ctgttttatt	gcagcttata	atgggtacaa	6660
ataaagcaat	agcatcacaa	atttcacaaa	taaagcattt	ttttcactgc	attctagttg	6720
tggtttgtcc	aaactcatca	atgtatctta	tcagtgtctg	atcccgctga	cctcgagagc	6780
ttggcgtaat	catggtcata	gctgtttcct	gtgtgaaatt	gttatccgct	cacaattcca	6840
cacaacatac	gagccggaag	cataaagtgt	aaagcctggg	gtgcctaatg	agtgaagtaa	6900
ctcacattaa	ttgcgttgcg	ctcaactgcc	gctttccagt	cgggaaacct	gtcgtgccag	6960
ctgcattaat	gaatcgccca	acgcgcgggg	agaggcggtt	tgcgtattgg	gcgctcttcc	7020
gcttctctcg	tcactgactc	gctgcgctcg	gtcgttcggc	tgcggcgagc	ggtatcagct	7080
cactcaaaag	cggtaatcac	gttatccaca	gaatcagggg	ataacgcagg	aaagaacatg	7140
tgagcaaaaag	gccagcaaaa	ggccaggaac	cgtaaaaagg	ccgcgttgct	ggcgtttttc	7200
cataggctcc	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	gagggtggcg	7260
aaccgcagac	gactataaag	ataccaggcg	tttccccctg	gaagctccct	cgtgcgctct	7320
cctgttccga	ccctgcgcgt	taccggatac	ctgtccgcct	ttctcccttc	gggaagcggt	7380
gcgctttctc	aatgctcacg	ctgtaggtat	ctcagttcgg	tgtaggtcgt	tcgctccaag	7440
ctgggctgtg	tgcacgaacc	ccccgttcag	cccgaccgct	gcgccttata	cggtactat	7500
cgtcttgagt	ccaaccgggt	aagacacgac	ttatcgccac	tggcagcagc	cactggtaac	7560
aggattagca	gagcgaggta	tgtaggcggt	gctacagagt	tcttgaagtg	gtggcctaac	7620
tacggctaca	ctagaaggac	agtatttggt	atctgcgctc	tgtgaagcc	agttaccttc	7680
ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cgggtggttt	7740
tttgtttgca	agcagcagat	tacgcgcaga	aaaaaaaggat	ctcaagaaga	tcctttgatc	7800
ttttctacgg	ggtctgacgc	tcagtggaa	gaaaactcac	gttaagggat	tttgggtcatg	7860
agattatcaa	aaaggatctt	cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca	7920
atctaaagta	tatatgagta	aacttggtct	gacagttacc	aatgcttaat	cagtgaggca	7980
cctatctcag	cgatctgtct	atctcggtca	tccatagttg	cctgactccc	cgtcgtgtag	8040
ataactacga	tacgggaggg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	8100
ccacgctcac	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	8160
agaagtgggc	ctgcaacttt	atccgcctcc	atccagctca	tttaattgtg	ccgggaagct	8220
agagtaagta	gttcgcaggt	taatagtttg	cgcaacggtg	ttgccattgc	tacaggcatc	8280
gtggtgtcac	gctcgtcggt	tggtagggct	tcattcagct	ccggttccca	acgatcaagg	8340
cgagttacat	gatcccccat	gttgtgcaaa	aaagcggtta	gctccttcgg	tcctccgatc	8400
gttgtcagaa	gtaagtggc	cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat	8460
tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	ctggtgagta	ctcaaccaag	8520
tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gccccgcgtc	aatacgggat	8580
aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	8640
cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	8700
cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	8760


```

aggcaaaatg cgcacaaaaa gggaataagg gcgacacgga aatgttgaat actcatactc 8820
ttcctttttc aatattattg aagcatttat caggggttatt gtctcatgag cggatacata 8880
tttgaatgta tttagaaaaa taaacaaata ggggttcgcg gcacatttcc ccgaaaagtg 8940
ccacctgacg tcgacggatc gggagatctc ccgatcccct atggtcgact ctcagtacaa 9000
tctgctctga tgccgcatag ttaagccagt atctgctccc tgcttggtg ttggaggtcg 9060
ctgagtagtg cgcgagcaaa atttaagcta caacaaggca aggcttgacc gacaattgca 9120
tgaagaatct gcttaggggtt aggcgttttg cgctgcttcg cgatgtacgg gccagatata 9180
cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 9240
atagcccata tatggagttc cgcgttacat aacttacggg aaatggcccg cctggctgac 9300
cgccccacga cccccgcccc ttgacgtcaa taatgacgta tgttcccata gtaacgcaa 9360
tagggacttt ccattgacgt caatgggtgg actatttacg gtaaactgcc cacttggcag 9420
tacatcaagt gtatcatatg ccaagtacgc ccctattga cgtcaatgac ggtaaatggc 9480
ccgcctggca ttatgccagc tacatgacct tatgggactt tcctacttgg cagtacatct 9540
acgtattagt catcgctatt accatgggtg                                     9569

```

<210> 15

<211> 401

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 15

```

Met Asp Ser Gly Ala Val Pro Thr Asn Ala Ser Asn Cys Thr Asp Pro
 1          5          10          15
Phe Thr His Pro Ser Ser Cys Ser Pro Ala Pro Ser Pro Ser Ser Trp
          20          25          30
Val Asn Phe Ser His Leu Glu Gly Asn Leu Ser Asp Pro Cys Gly Pro
          35          40          45
Asn Arg Thr Glu Leu Gly Gly Ser Asp Arg Leu Cys Pro Ser Ala Gly
          50          55          60
Ser Pro Ser Met Ile Thr Ala Ile Ile Ile Met Ala Leu Tyr Ser Ile
          65          70          75          80
Val Cys Val Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile
          85          90          95
Val Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn
          100          105          110
Leu Ala Leu Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser
          115          120          125
Val Asn Tyr Leu Met Gly Thr Trp Pro Phe Gly Thr Ile Leu Cys Lys
          130          135          140
Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr
          145          150          155          160
Leu Cys Thr Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val
          165          170          175
Lys Ala Leu Asp Leu Arg Thr Pro Arg Asn Ala Lys Ile Ile Asn Ile
          180          185          190
Cys Asn Trp Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met
          195          200          205
Ala Thr Thr Lys Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe
          210          215          220
Ser His Pro Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe
          225          230          235          240
Ile Phe Ala Phe Ile Met Pro Ile Leu Ile Ile Thr Val Cys Tyr Gly
          245          250          255

```

```

Leu Met Ile Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys
                260                265                270
Glu Lys Asp Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val
                275                280                285
Val Ala Val Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile
                290                295                300
Ile Lys Ala Leu Ile Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser
305                310                315                320
Trp His Phe Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro
                325                330                335
Val Leu Tyr Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu
                340                345                350
Phe Cys Ile Pro Thr Ser Ser Thr Ile Glu Gln Gln Asn Ser Thr Arg
                355                360                365
Ile Arg Gln Asn Thr Arg Asp His Pro Ser Thr Ala Asn Thr Val Asp
370                375                380
Arg Thr Asn His Gln Leu Glu Asn Leu Glu Ala Glu Thr Thr Pro Leu
385                390                395                400
Pro

```

<210> 16
 <211> 1415
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:/note =
 synthetic construct

```

<400> 16
gcctgacgct cctctctggc tccgcccggg ttggtcgctg taagaaataa caggagctgt 60
ggcagcggcg aagacgaagc ggctcgcgcg tggaaaccga aaagtcaggg tgctcgcggt 120
tactcccaac gtggtcccag ccggcggtca gcaccatgga cagcggcgcc gtccccacga 180
acgccagcaa ctgcaactgat cccttcacac acccttcaag ttgctcccca gcacctagtc 240
ccagctcctg ggtcaacttc tcccacttag aaggcaacct gtccgaccca tgcggtccga 300
accgcaccga gctgggaggg agcgacagac tgtgcccttc ggccggcagc ccttccatga 360
tcacggccat catcatcatg gccctctact ccatcgctgt cgtgggtggg ctcttcggaa 420
acttcctggt catgtatgtg attgtcaggt acacaaaaat gaagactgcc accaacatct 480
atattttcaa cctcgccctg gcagatgccc tggcaaccag taccctgcct ttccagagt 540
tcaattacct gatgggaaca tggccgtttg gaaccatcct gtgcaagatt gtgatctcca 600
tagattacta caatatgttc accagcatat tcaccctctg caccatgagt gtggatcgct 660
acattgcagt ctgccatcct gtcaaggccc tggatttaag cactccccgt aatgccaa 720
tcatcaacat ctgcaactgg atcctctctt cagccattgg tctgcctgtg atgttcatgg 780
caacgacaaa gtaccggcaa ggttccatag attgtacact aacattctct caccacaagt 840
ggtactggga aaacctgctg aaaatctgtg ttttcatctt tgccttcac atgcctatcc 900
tcatcattac agtgtgttat gggctgatga tcttacgcct caagagtgtc cgcagtctct 960
ctggctccaa agaaaaggac aggaacctgc gaagaatcac caggatgggt ctggtgggtg 1020
tggctgtgtt cattgtctgc tggacgccc ttcacatcta cgtcatcatt aaagccttga 1080
tcacaatccc ggaaactact ttccagaccg tttcctggca cttctgcatt gctctaggtt 1140
ataccaacag ttgcctcaac cccgtccttt atgcatttct ggatgaaaac ttcaaacgat 1200
gcttcagaga gttctgtatc ccaacttctt ccaccattga gcagcaaac tccactcgaa 1260
ttcgtcagaa caccagagac caccctcca cagccaatac ggtggatagg actaaccatc 1320
agctagaaaa tctggaagca gaaaccactc cgttacccta actgggtctc ataccattca 1380
gaccctcact gagcttagac gccacatcta tatga 1415

```

<210> 17
 <211> 398
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 17

```

Met Asp Ser Ser Ala Gly Pro Gly Asn Ile Ser Asp Cys Ser Asp Pro
 1           5           10           15
Leu Ala Pro Ala Ser Cys Ser Pro Ala Pro Gly Ser Trp Leu Asn Leu
          20           25           30
Ser His Val Asp Gly Asn Gln Ser Asp Pro Cys Gly Pro Asn Arg Thr
          35           40           45
Gly Leu Gly Gly Ser His Ser Leu Cys Pro Gln Thr Gly Ser Pro Ser
          50           55           60
Met Val Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile Val Cys Val
65           70           75           80
Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile Val Arg Tyr
          85           90           95
Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu
          100          105          110
Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Val Asn Tyr
          115          120          125
Leu Met Gly Thr Trp Pro Phe Gly Asn Ile Leu Cys Lys Ile Val Ile
          130          135          140
Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Cys Thr
145          150          155          160
Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu
          165          170          175
Asp Phe Arg Thr Pro Arg Asn Ala Lys Ile Val Asn Val Cys Asn Trp
          180          185          190
Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met Ala Thr Thr
          195          200          205
Lys Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe Ser His Pro
210          215          220
Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe Ile Phe Ala
225          230          235          240
Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Ile
          245          250          255
Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu Lys Asp
          260          265          270
Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Ala Val
          275          280          285
Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile Lys Ala
290          295          300
Leu Ile Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp His Phe
305          310          315          320
Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val Leu Tyr
          325          330          335
Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe Cys Ile
          340          345          350
Pro Thr Ser Ser Thr Ile Glu Gln Asn Ser Ala Arg Ile Arg Gln
          355          360          365
Asn Thr Arg Glu His Pro Ser Thr Ala Asn Thr Val Asp Arg Thr Asn
          370          375          380
His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
385          390          395

```

<210> 18

<211> 2229

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 18

```

cggatcctta gcatcccca agcgctccg tgtacttcta aggtgggagg gggatacaag 60
cagaggagaa tatcggacgc tcagacgttc cattctgcct gccgctcttc tctggttcca 120
ctagggttg tccttgtaag aaactgacgg agcctagggc agctgtgaga ggaagaggct 180
ggggcgccctg gaacccgaac actcttgagt gctctcagtt acagcctacc gagtccgcag 240
caagcattca gaaccatgga cagcagcgcc gggccaggga acatcagcga ctgctctgac 300
cccttagctc ctgcaagttg gtccccagca cctggctcct ggctcaactt gtcccacgtt 360
gatggcaacc agtccgaccc atgcggtcct aaccgcacgg ggcttgccgg gagccacagc 420
ctgtgccctc agaccggcag cccttccatg gtcacagcca tcaccatcat ggccctctat 480
tctatcgtgt gtgtagtggg cctctttgga aacttcctgg tcatgtatgt gattgtaaga 540
tataccaaaa tgaagactgc caccacatc tacattttca accttgcctt ggcagatgcc 600
ttagccacta gcacgtgccc ctttcagagt gttaactacc tgatgggaac gtggcccttt 660
ggaaacatcc tctgcaagat cgtgatctca atagactact acaacatgtt caccagtatc 720
ttcacctctc gcaccatgag tgtagaccgc tacattggcg tctgccaccg ggtcaaggcc 780
ctggatttcc gtacccccg aaatgccaaa attgtcaatg tctgcaactg gatcctctct 840
tctgccattg gtctgcccgt aatgttcatg gcaaccacaa aatacaggca ggggtccata 900
gattgcaccc tcactttctc tcatcccaca tgggtactggg agaacctgct caaaatctgt 960
gtcttcatct tcgccttcat catgccggtc ctcacatca ctgtgtgtta tggactgatg 1020
atcttacgac tcaagagtgt ccgcatgctg tcgggctcca aagaaaagga caggaacctg 1080
cgcaggatca cccggatggt gctggtggtc gtggctgtat ttattgtctg ctggaccccc 1140
atccacatct atgtcatcat caaagcactg atcacgattc cagaaaccac tttccagact 1200
gtttcctggc acttctgcat tgccttgggt tacacaaaca gctgcctgaa cccagttctt 1260
tatgcgttcc tggatgaaaa cttcaaacga tgttttagag agttctgcat cccaacttcc 1320
tccacaatcg aacagcaaaa ctctgctcga atccgtcaaa aactaggga acaccctcc 1380
acggctaata cagtggatcg aactaaccac cagctagaaa atctggaagc agaaactgct 1440
ccattgccct aactgggtcc caccgcatcc agaccctcgc taaacttaga ggctgccatc 1500
tacttggaat caggttgctg tcagggtttg tgggaggctc tggtttctct gaaaagcatc 1560
tgatcctgca tcattcaaa gtcattcctc ctggctattc acgtacacg tcagagacac 1620
tcagactgtg tcaagcactc agaaggaaga gactgcaggc cactactgaa tccagctcat 1680
gtacagaaac atccaatgga ccacaatact ctgtggtatg tgatttgtga tcaacataga 1740
aggtgacctt tccctatgtg gaatttttaa tttcaaggaa atacttatga tctcatcaag 1800
ggaaaaatag atgtcacttg ttaaattcac tgtagtgtat cataaaggaa aagctacctc 1860
tgacctctag cccagtcacc ctctatggaa agttccatag ggaatatgtg agggaaaatg 1920
ttgcttccaa attaaatttt cacctttatg ttatagtcta gttaagacat cagggggcatc 1980
tctgtttctt ggttttgtat tgtttgaaag aagacatctt cctccctagc tgcgtgttga 2040
aaatgaaagg gatttaaaac acagtgtcaa ctgcagaata gttgattctc gactgaagg 2100
gggggggcta atcttcccaa ttctttccat gtctccaag tggtcacaag gtcaaactca 2160
gagtcaccca gtaagctcat catgccacca ttctgagcaa aatccttga ttctgtctca 2220
gaatggtgg

```

<210> 19

<211> 398

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 19

```

Met Asp Ser Ser Thr Gly Pro Gly Asn Thr Ser Asp Cys Ser Asp Pro
 1           5           10          15
Leu Ala Gln Ala Ser Cys Ser Pro Ala Pro Gly Ser Trp Leu Asn Leu
          20          25          30

```

Ser His Val Asp Gly Asn Gln Ser Asp Pro Cys Gly Leu Asn Arg Thr
 35 40 45
 Gly Leu Gly Gly Asn Asp Ser Leu Cys Pro Gln Thr Gly Ser Pro Ser
 50 55 60
 Met Val Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile Val Cys Val
 65 70 75 80
 Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile Val Arg Tyr
 85 90 95
 Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu
 100 105 110
 Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Val Asn Tyr
 115 120 125
 Leu Met Gly Thr Trp Pro Phe Gly Thr Ile Leu Cys Lys Ile Val Ile
 130 135 140
 Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Cys Thr
 145 150 155 160
 Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu
 165 170 175
 Asp Phe Arg Thr Pro Arg Asn Ala Lys Ile Val Asn Val Cys Asn Trp
 180 185 190
 Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met Ala Thr Thr
 195 200 205
 Lys Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe Ser His Pro
 210 215 220
 Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Gly Ile Phe Ala
 225 230 235 240
 Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Ile
 245 250 255
 Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu Lys Asp
 260 265 270
 Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Val Ala Val
 275 280 285
 Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile Lys Ala
 290 295 300
 Leu Ile Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp His Phe
 305 310 315 320
 Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val Leu Tyr
 325 330 335
 Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe Cys Ile
 340 345 350
 Pro Thr Ser Ser Thr Ile Glu Gln Gln Asn Ser Thr Arg Val Arg Gln
 355 360 365
 Asn Thr Arg Glu His Pro Ser Thr Ala Asn Thr Val Asp Arg Thr Asn
 370 375 380
 His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
 385 390 395

<210> 20

<211> 1401

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 20

cactgagcttc tgectgccgc tcttctctgg ttccactagg gctgggtccat gtaagaatct 60
 gacggagcct agggcagctg tgagaggaag aggctggggc gcgtggaacc cgaaaagtct 120
 gagtgtcttc agttacagcc tacctagtcc gcagcaggcc ttcagcacca tggacagcag 180

```

caccggccca gggaacacca gcgactgctc agaccctta gctcaggcaa gttgctcccc 240
agcacctggc tcctgggtca acttgctcca cgttgatggc aaccagtcgc atccatgcgg 300
tctgaaccgc accgggcttg gcggaacga cagcctgtgc cctcagaccg gcagcccttc 360
catggtcaca gccattacca tcatggccct ctactctatc gtgtgtgtag tgggcctctt 420
cggaacttc ctggcatgt atgtgattgt aagatacacc aaaatgaaga ctgccaccaa 480
catctacatt ttcaaccttg ctctggcaga cgccttagcg accagtacac tgccctttca 540
gagtgtcaac tacctgatgg gaacatggcc cttcggaacc atcctctgca agatcgtgat 600
ctcaatagat tactacaaca tgttcaccag catattcacc ctctgcacca tgagcgtgga 660
ccgctacatt gctgtctgcc acccagtcaa agccctggat ttcggtaccg cccgaaatgc 720
caaaatcgct aacgtctgca actggatcct ctctctgcc atcgggtctgc ctgtaatgtt 780
catggcaacc acaaaatata ggcaggggtc catagattgc accctcacgt tctcccacc 840
aacctggtac tgggagaacc tgctcaaat ctgtgtcttt atcttcgctt tcatcatgcc 900
ggtcctcatc atcactgtgt gttacggcct gatgatctta cgactcaaga gcgttcgcat 960
gctatcgggc tccaaagaaa aggacaggaa tctgcgcagg atcaccgga tgggtgctggt 1020
ggtcgtggct gtatttatcg tctgctggac ccccatccac atctacgtca tcatcaaagc 1080
gctgatcacg attccagaaa ccacatttca gacggtttcc tggcacttct gcattgcttt 1140
gggttacacg aacagctgcc tgaatccagt tctttacgcc ttcctggatg aaaacttcaa 1200
gcgatgcttc agagagttct gcacccaac ctcgtccacg atcgaacagc aaaactccac 1260
tcgagtcctg cagaacacta gggaacatcc ctccacggct aatacagtgg atcgaactaa 1320
ccaccagcta gaaaatctgg aggcagaaac tgctccattg ccctaactgg gtctcacacc 1380
atccagaccg tcgctaagct t                                     1401

```

<210> 21

<211> 401

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 21

```

Met Asp Ser Ser Ala Asp Pro Arg Asn Ala Ser Asn Cys Thr Asp Pro
1      5      10      15
Phe Ser Pro Ser Met Cys Ser Pro Val Pro Ser Pro Ser Ser Trp
20      25      30
Val Asn Phe Ser His Leu Glu Gly Asn Leu Ser Asp Pro Cys Ile Arg
35      40      45
Asn Arg Thr Glu Leu Gly Gly Ser Asp Ser Leu Cys Pro Pro Thr Gly
50      55      60
Ser Pro Ser Met Val Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile
65      70      75      80
Val Cys Val Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile
85      90      95
Val Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn
100     105     110
Leu Ala Leu Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser
115     120     125
Val Asn Tyr Leu Met Gly Thr Trp Pro Phe Gly Thr Ile Leu Cys Lys
130     135     140
Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr
145     150     155     160
Leu Cys Thr Met Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val
165     170     175
Lys Ala Leu Asp Phe Arg Thr Pro Arg Asn Ala Lys Ile Ile Asn Val
180     185     190
Cys Asn Trp Ile Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met
195     200     205
Ala Thr Thr Lys Tyr Arg Asn Gly Ser Ile Asp Cys Ala Leu Thr Phe
210     215     220

```

Ser His Pro Thr Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe
 225 230 235 240
 Ile Phe Ala Phe Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly
 245 250 255
 Leu Met Ile Leu Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys
 260 265 270
 Glu Lys Asp Arg Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val
 275 280 285
 Val Ala Val Phe Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile
 290 295 300
 Ile Lys Ala Leu Ile Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser
 305 310 315 320
 Trp His Phe Cys Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro
 325 330 335
 Val Leu Tyr Ala Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu
 340 345 350
 Phe Cys Ile Pro Thr Ser Ser Thr Ile Glu Gln Gln Asn Ser Ala Arg
 355 360 365
 Ile Arg Gln Asn Thr Arg Asp His Pro Ser Thr Ala Asn Thr Val Asp
 370 375 380
 Arg Thr Asn His Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu
 385 390 395 400
 Pro

<210> 22

<211> 1881

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 22

atgagctgtg gtgtacttct aagatgggag ggggcaacaa gcagaagata atgtcagaag 60
 cttagctctc cttctgcctg acgctcctct ctgggtccgc ctgggttggc ttctgtaaga 120
 agtagcagga gccgtggcgg gggctggagg aagcggctga ggcgcgtgga acccgaaaag 180
 cccgggtgat cgcggttacc tctactgcgtg gtcccagccg cccagccgtc agcaccatgg 240
 acagcagcgc tgacccccga aacgccagca attgcaactga tcccttctcg ccctcttcaa 300
 tgtgtctccc agtacctagc cccagctcct ggggtcaactt ctcccaactta gaaggcaacc 360
 tgtccgaccc atgcattcgg aaccgcaccg agctgggagg gagcgacagc ctgtgccctc 420
 cgaccggcag tcttcccatg gtcacggcca tcaccatcat ggccctctac tccatcgtgt 480
 gcgtgggtggg tctcttcgga aacttcctgg tcatgtatgt gattgtcaga tacacaaaaa 540
 tgaagactgc caccaacatc tatattttca acctgtctct ggcggatgcc ttagccacca 600
 gtaccctacc cttccagagt gtcaattacc taatgggaac gtggccggtt ggaaccatcc 660
 tctgcaagat cgtgatctcc atagattact acaatatgtt caccagcata ttcaccctct 720
 gtaccatgag cgtggatcgc tacatcgccg tctgccatcc cgtcaaggcc ctggacttcc 780
 gcactccccg caacgccaaa atcatcaacg tctgcaactg gatcctctct tcagccattg 840
 gtctgcctgt gatgttcatg gcaacaacaa agtaccggaa tggttccata gattgtgcac 900
 taacattctc tcacccaacc tggtagctgg aaaacctgct gaaaatctgt gttttcatct 960
 ttgccttcat catgcctgtc ctcattcatta cgggtgtgta tgggctgatg atcttacgcc 1020
 tcaagagtgt tcgcatgtct tctggctcca aagaaaagga taggaacctg cgaagaatca 1080
 ccaggatggg gctgggtggt gtggctgtgt tcattgtctg ctggactccc attcacattt 1140
 acgtcatcat taaagccttg attacaattc cagaaactac tttccagact gtgtcctggc 1200
 acttctgcat tgctctaggt tatacaaaca gctgcctgaa cccagtcctt tatgcatttc 1260
 tggatgaaaa cttcaaacga tgcttcagag agttctgtat cccaacctcc tccaccattg 1320
 agcagcaaaa ctccgctcga atccgtcaaa acaccagaga ccacctctcc acggccaaca 1380
 cgggtggacag gaccaaccat cagctagaaa atctggaagc agaaactgct ccattgccct 1440
 aaccagggtg catgccattc agatcctcaa tgagctaaga cagccaccat ctacgtggaa 1500

```

gcagggttgcc atgagaatgt gtgggaggca ctattttcct aggaaagtgc ctgctctgag 1560
tcatcaaatc tgtttcctct ctggccgctc tgctctgcac atgagaggga catccaaact 1620
aaatcaagca ctaggaagga aagaactaat ccacatggag tttgcctgtg cacataatct 1680
caaggaagat gacccatggg accgaaacat gctgtggtat gtgcgttgag gtcacctca 1740
aagatggccc ttctgtatgt aatgtgctgt tttcaagcaa atgtttacgt cctcatcaaa 1800
gaaaaaatgt cagttgttaa attcaccata gtaacttgta aaggctacct ctgatcgaag 1860
catcttatgt ggaaatccaa g                                     1881

```

<210> 23

<211> 372

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 23

```

Met Glu Pro Ala Pro Ser Ala Gly Ala Glu Leu Gln Pro Pro Leu Phe
1           5           10           15
Ala Asn Ala Ser Asp Ala Tyr Pro Ser Ala Phe Pro Ser Ala Gly Ala
20           25           30
Asn Ala Ser Gly Pro Pro Gly Ala Arg Ser Ala Ser Ser Leu Ala Leu
35           40           45
Ala Ile Ala Ile Thr Ala Leu Tyr Ser Ala Val Cys Ala Val Gly Leu
50           55           60
Leu Gly Asn Val Leu Val Met Phe Gly Ile Val Arg Tyr Thr Lys Met
65           70           75           80
Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala
85           90           95
Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Ala Lys Tyr Leu Met Glu
100          105          110
Thr Trp Pro Phe Gly Glu Leu Leu Cys Lys Ala Val Leu Ser Ile Asp
115          120          125
Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Thr Met Met Ser Val
130          135          140
Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu Asp Phe Arg
145          150          155          160
Thr Pro Ala Lys Ala Lys Leu Ile Asn Ile Cys Ile Trp Val Leu Ala
165          170          175
Ser Gly Val Gly Val Pro Ile Met Val Met Ala Val Thr Arg Pro Arg
180          185          190
Asp Gly Ala Val Val Cys Met Leu Gln Phe Pro Ser Pro Ser Trp Tyr
195          200          205
Trp Asp Thr Val Thr Lys Ile Cys Val Phe Leu Phe Ala Phe Val Val
210          215          220
Pro Ile Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Leu Leu Arg Leu
225          230          235          240
Arg Ser Val Arg Leu Leu Ser Gly Ser Lys Glu Lys Asp Arg Ser Leu
245          250          255
Arg Arg Ile Thr Arg Met Val Leu Val Val Val Gly Ala Phe Val Val
260          265          270
Cys Trp Ala Pro Ile His Ile Phe Val Ile Val Trp Thr Leu Val Asp
275          280          285
Ile Asp Arg Arg Asp Pro Leu Val Val Ala Ala Leu His Leu Cys Ile
290          295          300
Ala Leu Gly Tyr Ala Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe
305          310          315          320
Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Gln Leu Cys Arg Lys Pro
325          330          335

```


Cys Gly Arg Pro Asp Pro Ser Ser Phe Ser Arg Ala Arg Glu Ala Thr
 340 345 350
 Ala Arg Glu Arg Val Thr Ala Cys Thr Pro Ser Asp Gly Pro Gly Gly
 355 360 365
 Gly Ala Ala Ala
 370

<210> 24

<211> 1773

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
 synthetic construct

<400> 24

```

ccgaggagacc tgcgctgctc ctggctcaca gcgctccggg cgaggagagc gggcgggaccg 60
gggggctggg ccggtgcggg cggcgaggca ggcggacgag ggcagagac agcggggcg 120
ccggggcgcg gcacgcggcg ggtcggggcc ggctctgccc ttgcgctcc cctcgcgtcg 180
gatccccgcg cccaggcagc cgggtggagag ggacgcggcg gacgccggca gccatggaac 240
cggccccctc cgccggcgcc gagctgcagc ccccgctctt cgccaacgcc tcggacgcct 300
accctagcgc cttccccagc gctggcgcca atgcgtcggg gccgccaggc gcgcggagcg 360
cctcgtccct cgccctggca atcgccatca ccgcgtcta ctgcggcgtg tgcgccgtgg 420
ggctgctggg caacgtgctt gtcattgttc gcatcgtccg gtacactaag atgaagacgg 480
ccaccaacat ctacatcttc aacctggcct tagccgatgc gctggccacc agcacgctgc 540
ctttccagag tgccaagtac ctgatggaga cgtggccctt cggcgagctg ctctgcaagg 600
ctgtgctctc catcgactac tacaatatgt tcaccagcat cttcacgctc accatgatga 660
gtgttgaccg ctacatcgct gtctgccacc ctgtcaaggc cctggacttc cgcacgcctg 720
ccaaggccaa gctgatcaac atctgtatct gggctcctggc ctcaggcggt ggcggtgcca 780
tcatggctcat ggctgtgacc cgtccccggg acggggcagt ggtgtgcatg ctccagttcc 840
ccagccccag ctggtactgg gacacggtga ccaagatctg cgtgttcctc ttgccttcg 900
tggtgcccac cctcatcatc accgtgtgct atggcctcat gctgctgcgc ctgcgcagtg 960
tgcgccctgct gtccgggctcc aaggagaagg accgcagcct gggcgcatc acgcgcagtg 1020
tgctgggtgt tgtgggccc ttctgtgtgt gttgggccc catccacatc ttctcatcg 1080
tctggacgct ggtggacatc gaccggcgcg acccgctggt ggtggctgcg ctgcacctgt 1140
gcagcgcgct gggctacgcc aatagcagcc tcaaccccg tctctacgct ttctctgacg 1200
agaacttcaa gcgctgcttc cgccagctct gccgcaagcc ctgcggccgc ccagacccca 1260
gcagcttcag ccgcgcccgc gaagccacgg ccgcgagcg tgtaccgcgc tgcaccccg 1320
ccgatggtcc cggcggtggc gctgccgct gaccaggcca tccggccccc agacgcccct 1380
ccctagtgtg acccgaggc cacatgagtc ccagtgggag gcgcgagcca tgatgtggag 1440
tgggggcagt agataggtcg gagggctttg ggaccgccag atggggcctc tgtttcggag 1500
acgggaccgg gccgctagat gggcatgggg tgggcctctg gtttggggcg aggcagagga 1560
cagatcaatg gcgcagtgcc tctggtctgg gtgccccgt ccacggctct aggtggggcg 1620
ggaaagccag tgactccagg agaggagcgg gacctgtggc tctacaactg agtccttaa 1680
cagggcatct ccaggaaggc ggggcttcaa ccttgagaca gcttcgggtt ctaacttga 1740
gccggacttt cggagttggg gggccgggg ccc 1773

```

<210> 25

<211> 228

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
 synthetic construct

<400> 25

Gly Ile Val Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile
 1 5 10 15

```

Phe Asn Leu Ala Leu Ala Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe
      20      25      30
Gln Ser Ala Lys Tyr Leu Met Glu Thr Trp Pro Phe Gly Glu Leu Leu
      35      40      45
Cys Lys Ala Val Leu Ser Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile
      50      55      60
Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val Cys His
      65      70      75      80
Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Ala Lys Ala Lys Leu Ile
      85      90      95
Asn Ile Cys Ile Trp Val Leu Ala Ser Gly Val Gly Val Pro Ile Met
      100      105      110
Val Met Ala Val Thr Arg Pro Arg Asp Gly Ala Val Val Cys Met Leu
      115      120      125
Gln Phe Pro Ser Pro Ser Trp Tyr Trp Asp Thr Val Thr Lys Ile Cys
      130      135      140
Val Phe Leu Phe Ala Phe Val Val Pro Ile Leu Val Ile Thr Val Cys
      145      150      155      160
Tyr Gly Leu Met Leu Leu Arg Leu Arg Ser Val Arg Leu Leu Ser Gly
      165      170      175
Ser Lys Glu Lys Asp Arg Ser Leu Arg Arg Ile Thr Arg Met Val Leu
      180      185      190
Val Val Val Gly Ala Phe Val Val Cys Trp Ala Pro Ile His Ile Phe
      195      200      205
Val Ile Val Trp Thr Leu Val Asp Ile Asp Arg Arg Asp Pro Leu Val
      210      215      220
Val Ala Ala Leu
      225

```

<210> 26

<211> 372

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 26

```

Met Glu Pro Val Pro Ser Ala Arg Ala Glu Leu Gln Phe Ser Leu Leu
  1      5      10      15
Ala Asn Val Ser Asp Thr Phe Pro Ser Ala Phe Pro Ser Ala Ser Ala
      20      25      30
Asn Ala Ser Gly Ser Pro Gly Ala Arg Ser Ala Ser Ser Leu Ala Leu
      35      40      45
Ala Ile Ala Ile Thr Ala Leu Tyr Ser Ala Val Cys Ala Val Gly Leu
      50      55      60
Leu Gly Asn Val Leu Val Met Phe Gly Ile Val Arg Tyr Thr Lys Leu
      65      70      75      80
Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala
      85      90      95
Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Ala Lys Tyr Leu Met Glu
      100      105      110
Thr Trp Pro Phe Gly Glu Leu Leu Cys Lys Ala Val Leu Ser Ile Asp
      115      120      125
Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Thr Met Met Ser Val
      130      135      140
Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu Asp Phe Arg
      145      150      155      160

```

```

Thr Pro Ala Lys Ala Lys Leu Ile Asn Ile Cys Ile Trp Val Leu Ala
      165      170      175
Ser Gly Val Gly Val Pro Ile Met Val Met Ala Val Thr Gln Pro Arg
      180      185      190
Asp Gly Ala Val Val Cys Thr Leu Gln Phe Pro Ser Pro Ser Trp Tyr
      195      200      205
Trp Asp Thr Val Thr Lys Ile Cys Val Phe Leu Phe Ala Phe Val Val
      210      215      220
Pro Ile Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Leu Leu Arg Leu
      225      230      235      240
Arg Ser Val Arg Leu Leu Ser Gly Ser Lys Glu Lys Asp Arg Ser Leu
      245      250      255
Arg Arg Ile Thr Arg Met Val Leu Val Val Val Gly Ala Phe Val Val
      260      265      270
Cys Trp Ala Pro Ile His Ile Phe Val Ile Val Trp Thr Leu Val Asp
      275      280      285
Ile Asn Arg Arg Asp Pro Leu Val Val Ala Ala Leu His Leu Cys Ile
      290      295      300
Ala Leu Gly Tyr Ala Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe
      305      310      315      320
Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Gln Leu Cys Arg Ala Pro
      325      330      335
Cys Gly Gly Gln Glu Pro Gly Ser Leu Arg Arg Pro Arg Gln Ala Thr
      340      345      350
Ala Arg Glu Arg Val Thr Ala Cys Thr Pro Ser Asp Gly Pro Gly Gly
      355      360      365
Gly Ala Ala Ala
      370

```

<210> 27

<211> 372

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 27

```

Met Glu Leu Val Pro Ser Ala Arg Ala Glu Leu Gln Ser Ser Pro Leu
  1      5      10      15
Val Asn Leu Ser Asp Ala Phe Pro Ser Ala Phe Pro Ser Ala Gly Ala
      20      25      30
Asn Ala Ser Gly Ser Pro Gly Ala Arg Ser Ala Ser Ser Leu Ala Leu
      35      40      45
Ala Ile Ala Ile Thr Ala Leu Tyr Ser Ala Val Cys Ala Val Gly Leu
      50      55      60
Leu Gly Asn Val Leu Val Met Phe Gly Ile Val Arg Tyr Thr Lys Leu
      65      70      75      80
Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala
      85      90      95
Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Ala Lys Tyr Leu Met Glu
      100      105      110
Thr Trp Pro Phe Gly Glu Leu Leu Cys Lys Ala Val Leu Ser Ile Asp
      115      120      125
Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Thr Met Met Ser Val
      130      135      140
Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu Asp Phe Arg
      145      150      155      160

```

Thr Pro Ala Lys Ala Lys Leu Ile Asn Ile Cys Ile Trp Val Leu Ala
 165 170 175
 Ser Gly Val Gly Val Pro Ile Met Val Met Ala Val Thr Gln Pro Arg
 180 185 190
 Asp Gly Ala Val Val Cys Met Leu Gln Phe Pro Ser Pro Ser Trp Tyr
 195 200 205
 Trp Asp Thr Val Thr Lys Ile Cys Val Phe Leu Phe Ala Phe Val Val
 210 215 220
 Pro Ile Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Leu Leu Arg Leu
 225 230 235 240
 Arg Ser Val Arg Leu Leu Ser Gly Ser Lys Glu Lys Asp Arg Ser Leu
 245 250 255
 Arg Arg Ile Thr Arg Met Val Leu Val Val Val Gly Ala Phe Val Val
 260 265 270
 Cys Trp Ala Pro Ile His Ile Phe Val Ile Val Trp Thr Leu Val Asp
 275 280 285
 Ile Asn Arg Arg Asp Pro Leu Val Val Ala Ala Leu His Leu Cys Ile
 290 295 300
 Ala Leu Gly Tyr Ala Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe
 305 310 315 320
 Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Gln Leu Cys Arg Thr Pro
 325 330 335
 Cys Gly Arg Gln Glu Pro Gly Ser Leu Arg Arg Pro Arg Gln Ala Thr
 340 345 350
 Thr Arg Glu Arg Val Thr Ala Cys Thr Pro Ser Asp Gly Pro Gly Gly
 355 360 365
 Gly Ala Ala Ala
 370

<210> 28

<211> 2219

<212> DNA

<213> Artificial Sequence

<220>

 <223> Description of Artificial Sequence:/note =
 synthetic construct

<400> 28

ctctaaaggc tgggtccctg cgcccagggc gcacggtgga gacggacacg gcggcgccat 60
 ggagctgggtg cctctgccc gtgcggagct gcagtcctcg cccctcgta acctctcgga 120
 cgcttttccc agcgcttcc ccagcgagg cgccaatgcg tcggggtcgc ggggagcccg 180
 tagtgctcgc tccctcgccc tagccatcgc catcacgcg ctctactcgg ctgtgtgcgc 240
 agtggggctt ctgggcaacg tgctcgatc gtttggcatc gtccggtaca ccaaattgaa 300
 gaccgccacc aacatctaca tctcaatct ggctttggct gatgcgctgg ccaccagcac 360
 gctgcccttc cagagcgcca agtacttgat ggaaacgtgg ccgtttggcg agctgctgtg 420
 caaggctgtg ctctccattg actactacaa catgttcaact agcatcttca cctcaccat 480
 gatgagcgtg gaccgctaca ttgctgtctg ccacctgtc aaagccctgg acttccggac 540
 accagccaag gccaagctga tcaatatatg catctgggtc ttggcttcag gtgtcggggt 600
 ccccatcatg gtcattggcag tgacccaacc ccgggatggg gcagtggtat gcatgctcca 660
 gttccccagt cccagctggt actgggacac tgtgaccaag atctgcgtgt tcctctttgc 720
 ctctgtgggt ccatcctca tcatcacggt gtgctatggc ctcatgctac tgccctgcg 780
 cagcgtgcgt ctgctgtccg gttccaagga gaaggaccgc agcctgcggc gcatcacgcg 840
 catggtgctg gtgggtgggg gcgccttcgt ggtgtgctgg gcgcccaccc acatcttcgt 900
 catcgtctgg acgctgggtg acatcaatcg gcgcgaccca cttgtgggtg ccgcactgca 960
 cctgtgcatt gcgctgggct acgccaacag cagcctcaac ccggttctct acgccttcct 1020
 ggacgagaac ttcaagcgct gttccgcca gctctgtcgc acgcccctgcg gccgccaaga 1080
 acccggcagt ctccgtcgtc cccgccaggc caccacgcgt gagcgtgtca ctgcctgcac 1140
 cccctccgac ggcccggggc gtggcgctgc cgctgacct acccgacct ccccttaaac 1200
 gccctccca agtgaagtga tccagaggcc acaccgagct ccctgggagg ctgtggccac 1260

```

caccaggaca gctagaattg ggctgcaca gaggggagggc ctctgtggg gacggggcct 1320
gaggggatcaa aggtccagg ttggaacggt gggggtgagg aagcagagct ggtgattcct 1380
aaactgtatc cattagtaag gcctctccaa tgggacagag cctccgcctt gagataacat 1440
cgggttcttg cctttttgaa caccagctc cagtccaaga cccaaggatt ccagctccag 1500
gaaccaggag gggcagtgat ggggtcgatg atttggtttg gctgagagtc ccagcatttg 1560
tgttatgggg aggatctctc atcttagaga agataagggg acagggcatt caggcaaggc 1620
agcttggggg ttggtcagga gataagcgcc cccttccctt ggggggagga taagtggggg 1680
atggtcaacg ttggagaaga gtcaaagttc tcaccacctt tctaactact cagctaaact 1740
cgttgaggct agggccaacg tgacttctct gtagagagga tacaagccgg gcctgatggg 1800
gcaggcctgt gtaatcccag tcatagtggg ggctgaggct ggaaaattaa ggaccaacag 1860
cctgggcaat ttagtgtctc aaaataaaat gtaaagaggg ctgggaatgt agctcagtgg 1920
taggggtgtt gtgtgaggct ctgggatcaa taagacaaaa caaccaacca accaaaaacc 1980
ttccaaacaa caaaaccaac cctcaaacca aaaaactatg tgggtgtctc tgagtctggt 2040
ttgaagagaa cccgcagccc tgtatccctg tggggctgtg gacagtgggc agaagcagag 2100
gctccctgga tcctgaacaa gggcccaaaa agcaagttct aaagggaccc ctgaaaccga 2160
gtaagccttt gtgtcaagaa gtgggagtag aaccagaaag gtggctgagt gctttagag 2219

```

<210> 29

<211> 380

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 29

```

Met Asp Ser Pro Ile Gln Ile Phe Arg Gly Glu Pro Gly Pro Thr Cys
1           5           10          15
Ala Pro Ser Ala Cys Leu Pro Pro Asn Ser Ser Ala Trp Phe Pro Gly
20          25          30
Trp Ala Glu Pro Asp Ser Asn Gly Ser Ala Gly Ser Glu Asp Ala Gln
35          40          45
Leu Glu Pro Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala
50          55          60
Val Tyr Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val
65          70          75          80
Met Phe Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile
85          90          95
Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala Leu Val Thr Thr Thr Met
100         105         110
Pro Phe Gln Ser Thr Val Tyr Leu Met Asn Ser Trp Pro Phe Gly Asp
115         120         125
Val Leu Cys Lys Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr
130         135         140
Ser Ile Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val
145         150         155         160
Cys His Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Leu Lys Ala Lys
165         170         175
Ile Ile Asn Ile Cys Ile Trp Leu Leu Ser Ser Ser Val Gly Ile Ser
180         185         190
Ala Ile Val Leu Gly Gly Thr Lys Val Arg Glu Asp Val Asp Val Ile
195         200         205
Glu Cys Ser Leu Gln Phe Pro Asp Asp Asp Tyr Ser Trp Trp Asp Leu
210         215         220
Phe Met Lys Ile Cys Val Phe Ile Phe Ala Phe Val Ile Pro Val Leu
225         230         235         240
Ile Ile Ile Val Cys Tyr Thr Leu Met Ile Leu Arg Leu Lys Ser Val
245         250         255
Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile
260         265         270

```

Thr Arg Leu Val Leu Val Val Val Ala Val Phe Val Val Cys Trp Thr
 275 280 285
 Pro Ile His Ile Phe Ile Leu Val Glu Ala Leu Gly Ser Thr Ser His
 290 295 300
 Ser Thr Ala Ala Leu Ser Ser Tyr Tyr Phe Cys Ile Ala Leu Gly Tyr
 305 310 315 320
 Thr Asn Ser Ser Leu Asn Pro Ile Leu Tyr Ala Phe Leu Asp Glu Asn
 325 330 335
 Phe Lys Arg Cys Phe Arg Asp Phe Cys Phe Pro Leu Lys Met Arg Met
 340 345 350
 Glu Arg Gln Ser Thr Ser Arg Val Arg Asn Thr Val Gln Asp Pro Ala
 355 360 365
 Tyr Leu Arg Asp Ile Asp Gly Met Asn Lys Pro Val
 370 375 380

<210> 30

<211> 1154

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 30

atggactccc cgatccagat cttccgcggg gagccggggc ctacctgegc cccgagcgcc 60
 tgccctgcccc ccaacagcag cgcctggttt cccggctggg ccgagcccca cagcaacggc 120
 agcgccgggt cggaggacgc gcagctggag cccgcgcaca tctccccggc catcccggtc 180
 atcatcacgg cggctctactc cgtagtgttc gtcgtgggct tgggtgggcaa ctcgctggtc 240
 atgttcgtga tcatccgata cacaaagatg aagacagcaa ccaacattta catatttaac 300
 ctggcttttg cagatgcttt agttactaca accatgccct ttcagagtac ggtctacttg 360
 atgaattcct ggccttttgg ggatgtgctg tgcaagatag taatttccat tgattactac 420
 aacatgttca ccagcatctt caccttgacc atgatgagcg tggaccgcta cattgccgtg 480
 tgccaccccg tgaaggcttt ggacttccgc acacccttga aggcaaagat catcaatata 540
 tgcattctggc tgctgtcgtc atctgttggc atctctgcaa tagtcttgg aggcacaaaa 600
 gtcagggaag acgtcgatgt cattgagtgc tccttgagc tcccagatga tgactactcc 660
 tgggtgggacc tcttcatgaa gatctgcgtc ttcattcttg ccttcgtgat ccctgtcttc 720
 atcatcatcg tctgtacac cctgatgac ctgcgtctca agagcgtccg gctcctttct 780
 ggctcccag agaaagatcg caacctgcgt aggatcacca gactggctct ggtggtgggtg 840
 gcagtcttcg tctgtctgct gactccatt cacatattca tcttggtgga ggctctgggg 900
 agcacctccc acagcacagc tgctctctcc agctattact tctgcatcgc cttaggctat 960
 accaacagta gctgaatcc cattctctac gcctttcttg atgaaaactt caagcgggtgt 1020
 ttccgggact tctgctttcc actgaagatg aggatggagc ggcagagcac tagcagagtc 1080
 cgaaatacag ttcaggatcc tgcttacctg agggacatcg atgggatgaa taaaccagta 1140
 tgactagtgc tggg 1154

<210> 31

<211> 380

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 31

Met Glu Ser Pro Ile Gln Ile Phe Arg Gly Asp Pro Gly Pro Thr Cys
 1 5 10 15
 Ser Pro Ser Ala Cys Leu Leu Pro Asn Ser Ser Ser Trp Phe Pro Asn
 20 25 30

Trp Ala Glu Ser Asp Ser Asn Gly Ser Val Gly Ser Glu Asp Gln Gln
 35 40 45
 Leu Glu Ser Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala
 50 55 60
 Val Tyr Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val
 65 70 75 80
 Met Phe Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile
 85 90 95
 Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala Leu Val Thr Thr Thr Met
 100 105 110
 Pro Phe Gln Ser Ala Val Tyr Leu Met Asn Ser Trp Pro Phe Gly Asp
 115 120 125
 Val Leu Cys Lys Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr
 130 135 140
 Ser Ile Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val
 145 150 155 160
 Cys His Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Leu Lys Ala Lys
 165 170 175
 Ile Ile Asn Ile Cys Ile Trp Leu Leu Ala Ser Ser Val Gly Ile Ser
 180 185 190
 Ala Ile Val Leu Gly Gly Thr Lys Val Arg Glu Asp Val Asp Val Ile
 195 200 205
 Glu Cys Ser Leu Gln Phe Pro Asp Asp Glu Tyr Ser Trp Trp Asp Leu
 210 215 220
 Phe Met Lys Ile Cys Val Phe Val Phe Ala Phe Val Ile Pro Val Leu
 225 230 235 240
 Ile Ile Ile Val Cys Tyr Thr Leu Met Ile Leu Arg Leu Lys Ser Val
 245 250 255
 Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile
 260 265 270
 Thr Lys Leu Val Leu Val Val Val Ala Val Phe Ile Ile Cys Trp Thr
 275 280 285
 Pro Ile His Ile Phe Ile Leu Val Glu Ala Leu Gly Ser Thr Ser His
 290 295 300
 Ser Thr Ala Ala Leu Ser Ser Tyr Tyr Phe Cys Ile Ala Leu Gly Tyr
 305 310 315 320
 Thr Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe Leu Asp Glu Asn
 325 330 335
 Phe Lys Arg Cys Phe Arg Asp Phe Cys Phe Pro Ile Lys Met Arg Met
 340 345 350
 Glu Arg Gln Ser Thr Asn Arg Val Arg Asn Thr Val Gln Asp Pro Ala
 355 360 365
 Ser Met Arg Asp Val Gly Gly Met Asn Lys Pro Val
 370 375 380

<210> 32

<211> 1410

<212> DNA

<213> Artificial Sequence

<220>

 <223> Description of Artificial Sequence:/note =
 synthetic construct

<400> 32

gcgcaccttg ctgatcccaa acaggcagag cttcttccag tcttgaagg cacaaattga 60
 gcatcaggaa cgtggaccca tcagggtga acagctactc aggatctaaa gtggtgactt 120
 ggaaagctga cggtagcttg ggaaggagg tcgccaatca gcgatctgga gctgcagcgc 180
 tcaccatgga gtccccatt cagatcttcc gaggagatcc aggcctacc tgctctcca 240
 gtgcttgctt tctccccaac agcagctctt ggttcccaa ctgggcagaa tccgacagta 300

```

atggcagtgt gggctcagag gatcagcagc tggagtccgc gcacatctct ccggccatcc 360
ctgttatcat caccgctgtc tactctgtgg tatttgtggg gggcttagtg ggcaattctc 420
tggcatggtt tgtcatcatc cgatacacga agatgaagac cgcaaccaac atctacatat 480
ttaacctggc tttggcagat gctttgggta ctaccactat gccctttcag agtgctgtct 540
acttgatgaa ttcttggcct tttggagatg tgctatgcaa gattgtcatt tccattgact 600
actacaacat gtttaccagc atattcacct tgaccatgat gagtgtggac cgctacattg 660
ctgtgtgcca ccctgtgaaa gctttgggact tccgaacacc tttgaaagca aagatcatca 720
acatctgcat ttggctcctg gcatcatctg ttggtatata agcgatagtc cttggaggca 780
ccaaagtcat ggaagatgtg gatgtcattg aatgctcctt gcagtttcct gatgatgaat 840
attcctgggtg ggatctcttc atgaagatct gtgtcttctg ctttgccttt gtgatcccag 900
tcctcatcat cattgtctgc tacaccctga tgatcctgcg cctgaagagt gtccggctcc 960
tgtctggctc ccgagagaag gaccgaaatc tccgccgcat caccaagctg gtgctggtag 1020
tagttgcagt cttcatcatc tgttggacc ccatcaccat ctttatcctg gtggaggctc 1080
tgggaagcac ctcccacagc acagctgccc tctccagcta ttatttctgt attgccttgg 1140
gttataccaa cagcagcctg aatcctgttc tctatgcctt tctggatgaa aacttcaagc 1200
gggtgttttag ggacttctgc ttccctatta agatgcgaat ggagcgccag agcaccaata 1260
gagttagaaa cacagttcag gatcctgctt ccatgagaga tgtgggaggg atgaataagc 1320
cagtatgact agtcgtggaa atgtcttctt attgttctcc aggtagagaa gatttcaatg 1380
atcttgggtt aaccagatt acaactgcag                                     1410

```

<210> 33

<211> 380

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 33

```

Met Glu Ser Pro Ile Gln Ile Phe Arg Gly Glu Pro Gly Pro Thr Cys
1           5           10           15
Ala Pro Ser Ala Cys Leu Leu Pro Asn Ser Ser Ser Trp Phe Pro Asn
20           25           30
Trp Ala Glu Ser Asp Ser Asn Gly Ser Val Gly Ser Glu Asp Gln Gln
35           40           45
Leu Glu Pro Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala
50           55           60
Val Tyr Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val
65           70           75           80
Met Phe Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile
85           90           95
Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala Leu Val Thr Thr Thr Met
100          105          110
Pro Phe Gln Ser Ala Val Tyr Leu Met Asn Ser Trp Pro Phe Gly Asp
115          120          125
Val Leu Cys Lys Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr
130          135          140
Ser Ile Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val
145          150          155          160
Cys His Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Leu Lys Ala Lys
165          170          175
Ile Ile Asn Ile Cys Ile Trp Leu Leu Ala Ser Ser Val Gly Ile Ser
180          185          190
Ala Ile Val Leu Gly Gly Thr Lys Val Arg Glu Asp Val Asp Val Ile
195          200          205
Glu Cys Ser Leu Gln Phe Pro Asp Asp Glu Tyr Ser Trp Trp Asp Leu
210          215          220
Phe Met Lys Ile Cys Val Phe Val Phe Ala Phe Val Ile Pro Val Leu
225          230          235          240

```


Ile Ile Ile Val Cys Tyr Thr Leu Met Ile Leu Arg Leu Lys Ser Val
 245 250 255
 Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile
 260 265 270
 Thr Lys Leu Val Leu Val Val Val Ala Val Phe Ile Ile Cys Trp Thr
 275 280 285
 Pro Ile His Ile Phe Ile Leu Val Glu Ala Leu Gly Ser Thr Ser His
 290 295 300
 Ser Thr Ala Val Leu Ser Ser Tyr Tyr Phe Cys Ile Ala Leu Gly Tyr
 305 310 315 320
 Thr Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe Leu Asp Glu Asn
 325 330 335
 Phe Lys Arg Cys Phe Arg Asp Phe Tyr Phe Pro Ile Lys Met Arg Met
 340 345 350
 Glu Arg Gln Ser Thr Asn Arg Val Arg Asn Thr Val Gln Asp Pro Ala
 355 360 365
 Ser Met Arg Asp Val Gly Gly Met Asn Lys Pro Val
 370 375 380

<210> 34

<211> 2481

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note.=
 synthetic construct

<400> 34

```

tccagccctg cctgcacagg caaagtttgt ctctctgggt ccaatctgtc ctctgctcct 60
gcagcccggc aagtgccatc gctcctcggt ttccagctgc agcactcacc atggagtcct 120
ccatccagat tttccgcgga gagccaggcc ctacctgtgc tcccagtgtc tgcctactcc 180
ccaacagcag ctcttggttc cccaactggg ccgaatcgga cagcaatggc agtttgggct 240
ccgaggacca gcagctggag cccgcgcaca tctctccagc catccctgtt atcatcacg 300
ctgtctactc tgtggtggtt gtggtgggct tagtgggcaa ttccctggtc atgtttgtca 360
tcatccgata cacaagatg aagaccgcaa ccaacatcta catatttaac ctggctttgg 420
cagatgcttt gggtactacc actatgcctt tccagagtgc tgtctacttg atgaattctt 480
ggccttttgg agatgttctg tgcaagattg tcatttccat tgactactac aacatgttta 540
ccagcatatt caccttgacc atgatgagtg tggaccgcta cattgccgtg tgccaccctg 600
tgaaagcttt ggatttccga acacctttga aagcaaagat catcaacatc tgcatttggc 660
tactggcatc atctgttggg atatcagcga tagtccttgg aggcaccaa gtcagggaag 720
atgtggatgt cattgaatgc tccttgagc ttcttgatga tgaatattcc tgggtgggac 780
tcttcatgaa gatctgtgtc ttctgtttg cctttgttat ccctgtctta atcatcattg 840
tctgctacac cctgatgac ctgcgcttga agagtgtccg gctcctctcg ggctctcgag 900
agaaggaccg aaatctccgc cggatcacca agctgggtgt ggtagtgggt gcagtcttca 960
tcattctgtt gaccccatc cacatcttta tcttggtcga ggctctaggc agcacctctc 1020
acagcacagc tgtcctctct agctattact tctgcattgc cttgggttat accaacagca 1080
gcttgaatcc tgttctctat gcctttcttg atgaaaactt caagcgggtg tttagggact 1140
tctgcttccc cattaagatg cgaatggagc gccagagcac aaacagagtt agaaacacag 1200
ttcaggatcc tgcttccatg agggatgtgg gtgggatgaa taagccagta tgactagtca 1260
tggaatgtc ttctattgt tctccgggta gagaagagtt caatgatctt ggtttaacct 1320
agattaccac tgcagtctga agaggaaaga tgagggtatt aataacttag ccatgttatg 1380
caatctaaag gtgcagggca cattagtgat ctaggctgag taggggcagc aagtgtgaag 1440
aacagagcac atgtcctggc aacaatacac ctctttccta ggacagagga gaaggcaatc 1500
taacctcaac ccttcgataa acagacagca ctctttcttc tgggtcccctt gatttactgc 1560
acctccatct gcgtggcctt tctgtgtaac atagttccaa agctctagag aagaaaatga 1620
aagaaaaagt gcatttgatc caaaacttac tgggcaccca accttcgcgt taacacaggc 1680
aaaaccaact tcgtatacaa tgaggccata ccaaggggtca tatccaactc actttctgct 1740
ggatgatcatg tcttatgtgt gatgagagtg aaacttagga agaacctgga ctacagacct 1800
gacactgggg gagagcacca tattgacatt tgtgaacctt tttaaagttg tgggtgttct 1860

```

```

cgtctcacag tgtctaagtc cttgaaaaac tacagttgct tcttaaggtc tctgggtttt 1920
agcatgctat tcaggaagat aatcttctga gaaaacatga actgatatta aaagggttgaa 1980
gcttaataacc agcaaagtgt gtgtaatttc atctgtaaat agtgggtctgt atataaataa 2040
ggaccagggtt ttcctgtcca gcctgtacat ttctcaagga tgccgtagac acaccctgg 2100
aggcatggaa agttcatgct gggatatttt gcttactat aagctacttt cttgatttgg 2160
tcttggtgtg atttctacta gattactcaa acattattta ctctaact gatcataact 2220
tggtgttaac aattcccaa actttgaatt cattctaaag tgtagcatt gatcaaact 2280
actttgtggt agcatctgtt tgtaaacaca cacatattgc cagattctct actcaggtag 2340
aggaagtgc tttgatctg tacacctca aatgttatgc tctggcttc cacagaaagt 2400
ggaattgttt caaatgcat gctgaaaaag gaaataggat ttgagatggc ttagcacaat 2460
ttgcatggta ttgagtaaga g                                     2481

```

<210> 35

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 35

tttttccagt tccgtttatc c 21

<210> 36

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 36

tttatcgcca atccacatct 20

<210> 37

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 37

cccatagtaa cgccaatagg 20

<210> 38

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 38

aaatgtgagc gagtaacaac c 21

<210> 39

<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 39
accacagtcc atgccatcac 20

<210> 40
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 40
tccaccaccc tggtgctgta 20

<210> 41

<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 41
cacctaatac gactcactat agg 23

<210> 42
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 42
cattaaccct cactaaag 18

<210> 43
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 43
atatcgatat cgctagcttt taaaagaaaa gggggg 35

<210> 44

<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 44
taatcgatgc taagcaaaat tttgaatttt tgtaatttg 39

<210> 45
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 45
gaattaccta atgggaacat gg 22

<210> 46
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 46
gcagacgatg aacacagc 18

<210> 47
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 47
accacagcaa tcac 14

<210> 48
<211> 10472
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 48
aatgtagtct tatgcaatac tctttagtc ttgcaacatg gtaacgatga gttagcaaca 60
tgccttacaa ggagagaaaa agcaccgtgc atgccgattg gtggaagtaa ggtggtacga 120

tctgtgcctta	ttaggaaggc	aacagacggg	tctgacatgg	attggacgaa	ccactgaatt	180
gccgcattgc	agagatat	tatttaagt	cctagctcga	tacataaacg	ggctctctctg	240
gttagaccag	atctgagcct	gggagctctc	tggctaacta	gggaaccac	tgcttaagcc	300
tcaataaagc	ttgccttgag	tgcttcaagt	agtgtgtgcc	cgtctgttgt	gtgactctgg	360
taactagaga	tcctcagac	ccttttagtc	agtgtgaaa	atctctagca	gtggcgccccg	420
aacagggact	tgaagcgaa	agggaaacca	gaggagctct	ctcgacgcag	gactcggctt	480
gctgaagcgc	gcacggcaag	aggcgagggg	cggcgactgg	tgagtacgcc	aaaaattttg	540
actagcggag	gctagaagga	gagagatggg	tgcgagagcg	tcagtattaa	gcgggggaga	600
attagatcgc	gatgggaaaa	aattcgggta	aggccagggg	gaaagaaaaa	atataaatta	660
aaacatatag	tatgggcaag	cagggagcta	gaacgattcg	cagttaatcc	tggcctgtta	720
gaaacatcag	aaggctgtag	acaaatactg	ggacagctac	aaccatccct	tcagacagga	780
tcagaagaac	ttagatcatt	atataataca	gtagcaaccc	tctatttgtt	gcatcaaagg	840
atagagataa	aagacaccaa	ggaagcttta	gacaagatag	aggaagagca	aaacaaaagt	900
aagaccaccg	cacagcaagc	ggccgctgat	cttcagacct	ggaggaggag	atatgaggga	960
caattggaga	agtgaattat	ataaatataa	agtagtaaaa	attgaacat	taggagtagc	1020
accaccaag	gcaaagagaa	gagtgggtgca	gagagaaaaa	agagcagtgg	gaataggagc	1080
tttgttcctt	gggttccttg	gagcagcagg	aagcactatg	ggcgacgcgt	caatgacgct	1140
gacggtacag	gccagacaat	tattgtcttg	tatagtgcag	cagcagaaca	atttgcctgag	1200
ggctattgag	gcgcaacagc	atctgttgca	actcacagtc	tggggcatca	agcagctcca	1260
ggcaagaatc	atggctgtgg	aaagatacct	aaaggatcaa	cagctcctgg	ggatttgggg	1320
ttgctctgga	aaactcattt	gcaccactgc	tgtgccttgg	aatgctagtt	ggagtaataa	1380
atctctggaa	cagatttgga	atcacacgac	ctggatggag	tgggacagag	aaattaacaa	1440
ttacacaagc	ttaatacact	ccttaattga	agaatcgcaa	aaccagcaag	aaaagaatga	1500
acaagaatta	ttggaattag	ataaatgggc	aagtttgttg	aattgggtta	acataacaaa	1560
ttggctgtgg	tataataaat	tattcataat	gatagtagga	ggcttggtag	gtttaagaat	1620
agtttttgc	gtactttcta	tagtgaatag	agttaggcag	ggatattcac	cattatcgtt	1680
tcagaccacc	ctcccaaccc	cgaggggacc	cgacaggccc	gaaggaaatg	aagaagaagg	1740
tggagagaga	gacagagaca	gatccattcg	attagtgaac	ggatctcgac	ggatctgata	1800
agcttgggag	ttccgcgtta	cataacttac	ggtaaatggc	ccgcctggct	gaccgcccac	1860
cgacccccgc	ccattgacgt	caataatgac	gtatgttccc	atagtaacgc	caatagggac	1920
tttccattga	cgtaaatggg	tggagtattt	acggtaaaact	gccacttgg	cagtacatca	1980
agtgtatcat	atgccaaagta	cgccccctat	tgacgtcaat	gacggtaaat	ggccccgctg	2040
gcattatgcc	cagtacatga	ccttatggga	ctttcctact	tggcagtaca	tctacgtatt	2100
agtcactcgt	attaccatgg	tgatgcgggt	tggcagtagc	atcaatggg	gtggatagcg	2160
gtttgactca	cggggatttc	caagtctcca	ccccattgac	gtcaatggga	gtttgttttg	2220
gcacccaaat	caacggggact	ttccaaaatg	tcgtaacaac	tcgcccccat	tgacgcaaat	2280
gggcggtagg	cgtgtacggg	gggaggtcta	tataagcaga	gctcgtttag	tgaaccgtca	2340
gatcgcttgg	agacgccatc	cacgctgttt	tgacctccat	agaagacacc	gactctagag	2400
gatccactag	tcagtggtgg	tgggaattgat	cccttcacct	aatacgactc	actataggct	2460
agctcctcct	ctgctcgccc	aatccttcca	acccccctatg	gtggtatggc	tgacacagaa	2520
aatgtctgct	cctgtatggg	acatttgccc	ctcttctcca	aatataagac	aggatgaggc	2580
ctagcttttg	ctgctccaaa	gttttaaaag	aacacattgc	acggcattta	gggactctaa	2640
aggggtggag	aggaatgagg	gaattgcatc	atgccaaagg	tggctcctcat	ccatcactgc	2700
ttccaggggc	cagagtggct	tccaggaggt	attcttaca	aggaagcccc	atctgtagct	2760
aacactcaga	gcccattttc	ctgcgttaac	ccctcccgac	ctcatataca	ggagtaacat	2820
gatcagtgac	ctgggggagc	tggccaaact	gcgggacctg	cccaagctga	gggccttggt	2880
gctgctggac	aacccctgtg	ccgatgagac	tgactaccgc	caggaggccc	tgggtgcagat	2940
ggcacaccta	gagcgcctag	acaaagagta	ctatgaggac	gaggaccggg	cagaagctga	3000
ggagatccga	cagaggctga	aggaggaaca	ggagcaagaa	ctcgaccggg	accaagacat	3060
ggaaccgtac	ctcccgccaa	cttagtggtc	cctctagcct	gcagggacag	taaagggtgat	3120
ggcaggaagg	cagcccccg	aggtcaaagg	ctgggcacgc	gggaggagag	gccagagtca	3180
gaggctgcgg	gtatctcaga	tatgaaggaa	agatgagaga	ggctcaggaa	gaggtaagaa	3240
aagacacaag	agaccagaga	agggagaaga	attagagagg	gaggcagagg	accgctgtct	3300
ctacagacat	agctggtaga	gactgggagg	aagggatgaa	ccctgagcgc	atgaagggaa	3360
ggagggtggc	ggtggtatat	ggaggatgta	gctgggcccag	ggaaaagatc	ctgcactaaa	3420
aatctgaagc	taaaaataac	aggacacggg	gtggagaggc	gaaaggaggg	cagattgagg	3480
cagagagact	gagaggcctg	gggatgtggg	cattccggta	gggcacacag	ttcacttgct	3540
ttctcttttt	ccaggaggcc	aragatgctg	acctcaagaa	ctcataatac	cccagtgggg	3600
accaccgcat	tcataagcct	gttacaagaa	gtgggagatg	ttcctttttg	ttccagactg	3660
gaaatccatt	acatcccag	gctcaggttc	tgtgggtggc	atctctgtgt	ggcttgttct	3720
gtgggcctac	ctaaagtcc	aagcacagct	ctcaagcaga	tccgaggcga	ctaagatgct	3780

agtaggggtt	gtctggagag	aagagccgag	gaggtgggct	gtgatggatc	agttcagctt	3840
tcaataaaaa	aggcgttttt	atattctgtg	tcgagttcgt	gaacccctgt	ggtgggcttc	3900
tccatctgtc	tgggttagta	cctgccacta	tactggaata	aggggacgcc	tgcttccctc	3960
gagttggctg	gacaaggtta	tgagcatccg	tgtacttatg	gggttgccag	cttggctctg	4020
gatcgcccg	gcccttcccc	caccgcttcg	gttccccacc	accacccgcg	ctcgtaactg	4080
cgtctccg	tgagctctct	gactcatcgg	ggcccccg	tcacatgcgc	tcgctcggtc	4140
ctataggcgc	cgccccctgc	ccaccccccg	cccgcgctgg	gagccgcagc	cgccgccact	4200
cctgctctct	ctgcgcgcgc	gccgtcacca	ccgccaccgc	caccggctga	gtctgcagtc	4260
ctcgaggaac	tgaaaaacca	gaaagttaac	tggtaagttt	agtctttttg	tcttttattt	4320
caggtcccgg	atccggtggg	ggtgcaaatc	aaagaactgc	tcctcagtg	atgttgctt	4380
tacttctagg	cctgtacgga	agtgttactt	ctgctctaaa	agctgcggaa	ttgtacccgc	4440
ggccaagcta	agcttgatat	cgaattccgg	atgagcctct	gtgaactact	aaggtgggag	4500
ggggctatac	gcagaggaga	atgtcagatg	ctcagctcgg	tccccctcgc	ctgacgctcc	4560
tctctgtctc	agccaggact	ggtttctgta	agaaacagca	ggagctgtgg	cagcggcgaa	4620
aggaagcggc	tgaggcgctt	ggaacccgaa	aagtctcgg	gctcctggct	acctcgacaa	4680
gcgggtcccg	cccgcccgct	agtaccatgg	acagcagcgc	tgccccacg	aacgccagca	4740
attgcactga	tgccctggcg	tactcaagtt	gctccccagc	accagccccc	ggttccctgg	4800
tcaacttgct	ccacttagat	ggcaacctgt	ccgacccatg	cgggtccgaac	cgcaccgacc	4860
tgggccccgg	agacagcctg	tgccctccga	ccggcagtc	ctccatgatc	acggccatca	4920
cgatcatggc	cctctactcc	atcgtgtgcg	tgggtgggct	cttcggaaac	ttcctgggtc	4980
tgtatgtgat	tgtcagatac	accaagatga	agactgccac	caacatctac	atcttcaacc	5040
ttgctctggc	agatgcctta	gccaccagta	ccctgccctt	ccagagtgtg	aattacctaa	5100
tgggaacatg	gccatttgga	accatccttt	gcaagatagt	gatctccata	gattactata	5160
acatgttcac	cagcatatct	accctctgca	ccatgagtgt	tgatcgatac	attgcagtct	5220
gccaccctgt	caaggcctta	gatttccgta	ctccccgaaa	tgccaaaatt	atcaatgtct	5280
gcaactggat	cctctcttca	gccattgggt	ttcctgtaat	gttcatggct	acaacaaaat	5340
acaggcaagg	ttccatagat	tgtacactaa	cattctctca	tccaacctgg	tactgggaaa	5400
acctgctgaa	gatctgtggt	ttcatcttcg	ccttcattat	gccagtgtc	atcattaccg	5460
tgtgctatgg	actgatgatc	ttgcgcctca	agagtgtccg	catgctctct	ggctccaaag	5520
aaaaggacag	gaatcttcga	aggatcacca	ggatgggtg	ggtgggtggg	gctgtgttca	5580
tcgtctcggt	gactccatt	cacatttacg	tcattcattaa	agccttgggt	acaatcccag	5640
aaactacgtt	ccagactggt	tcttggcact	tctgcattgc	tctagggtac	acaaacagct	5700
gcctcaacc	agtcctttat	gcatttctgg	atgaaaactt	caaacgatgc	ttcagagagt	5760
tctgtatccc	aacctcttcc	aacattgagc	aacaaaactc	cactcgaatt	cgtcagaaca	5820
tctagagacc	acccctccac	ggccaataca	gtggatagaa	ctaatacatca	gctagaaaat	5880
ctggaagcag	aaactgctcc	ggtgcccgtc	gacccggg	gccgcttccc	tttagtgagg	5940
gttaatgaag	ggctcgagtc	tagagggcc	gctgttcgaa	ggtaagccta	tccctaacc	6000
tctctcggt	ctcgattcta	cgcgtaccg	ctagtaatga	gttggaatt	aattctgtgg	6060
aatgtgtgtc	agttagggtg	tggaaagtcc	ccaggctccc	caggcaggca	gaagtatgca	6120
aagcatgcat	ctcaattagt	cagcaaccag	gtgtggaaag	tcccaggct	ccccagcagg	6180
cagaagtatg	caaagcatgc	atctcaatta	gtcagcaacc	atagtcccgc	ccctaactcc	6240
gcccatccc	cccctaactc	cgcccagttc	cgcccattct	cgccccatg	gctgactaat	6300
tttttttatt	tatgcagagg	ccgaggccgc	ctctgcctct	gagctattcc	agaagtatg	6360
aggaggcttt	tttggaggcc	taggcttttg	caaaaagctc	ccgggagctt	gtatatccat	6420
tttcggatct	gatcagcag	tgttgacaat	taactcatcg	catagtatat	cggcatagta	6480
taatacgaca	aggtgaggaa	ctaaaccatg	gccaagcctt	tgtctcaaga	agaatccacc	6540
ctcattgaaa	gagcaacggc	tacaatcaac	agcatcccca	tctctgaaga	ctacagcgtc	6600
gccagcgcag	ctctctctag	cgacggccgc	atcttccact	gtgtcaatgt	atatcatttt	6660
actgggggac	cttgtgcaga	actcgtgggt	ctgggcactg	ctgctgctgc	ggcagctggc	6720
aacctgactt	gtatcgtcgc	gatcggaaat	gagaacaggg	gcatcttgag	cccctgcgga	6780
cggtgccgac	aggtgcttct	cgatctgcat	cctgggatca	aagccatagt	gaaggacagt	6840
gatggacagc	cgacggcagt	tgggattcgt	gaattgctgc	cctctgggtt	tgtgtgggag	6900
ggtaagcac	aattcgagct	cggtaccttt	aagaccaatg	acttacaagg	cagctgtaga	6960
tcttagccac	tttttaaaag	aaaagggggg	actggaaggg	ctaattcact	cccaacgaag	7020
acaagatctg	ctttttgctt	gtactgggtc	tctctgggtt	gaccagatct	gagcctggga	7080
gctctctggc	taactaggga	acccactgct	taagcctcaa	taaagcttgc	cttgagtgtc	7140
tcaagtatgt	tgtgcccgtc	tgttgtgtga	ctctggtaac	tagagatccc	tcagaccctt	7200
ttagtcatgt	tggaaaatct	ctagcagtag	tagttcatgt	catcttatta	ttcagtattt	7260
ataacttgca	aagaaatgaa	tatcagagag	tgagaggaac	ttgtttattg	cagcttataa	7320
tggttacaaa	taaagcaata	gcatacacia	tttcacaaa	aaagcatttt	tttctactga	7380
ttctagttgt	ggtttgtcca	aactcatcaa	tgtatcttat	catgtctggc	tctagctatc	7440

```

ccgcccctaa ctccgcccct cccgccccta actccgcccga gttccgcccga ttctccgccc 7500
catggctgac taatTTTTTT tatttatgca gaggcgagg cgcctcggc ctctgagcta 7560
ttccagaagt agtgaggagg cttttttgga ggcctaggga cgtaccaat tcgcccata 7620
gtgagtcgta ttacgcgcgc tccctggccg tcgttttaca acgtcgtgac tgggaaaacc 7680
ctggcgttac ccaacttaat cgccttgacg cacatcccc tttcgccagc tggcgtaata 7740
gcgaagaggc ccgcaccgat cgccttccc aacagttgag cagcctgaat ggcgatggg 7800
acgcgccttg tagcgcgca ttaagcgcg cggtgtggt gggtacgcgc agcgtgaccg 7860
ctacacttgc cagcgcccta ggcgcgctc ctttcgcttt cttcccttcc tttctcgcca 7920
cgttcgccgg ctttccccgt caagctctaa atcgggggct ccctttaggg ttccgattta 7980
gtgctttacg gcacctogac cccaaaaaac ttgattaggg tgatggttca cgtagtgggc 8040
catcgccctg atagacgggt tttcgccctt tgacgttggg gtccacgttc tttaatagt 8100
gactcttggt ccaactgga acaactca accctatctc ggtctattct tttgatttat 8160
aagggatttt ccgattttcg gctattggg taaaaaatga gctgatttaa caaaaattta 8220
acgcgaattt taacaaaata ttaacgctta caatttaggt ggcacttttc ggggaaatgt 8280
gcgcggaacc cctatttggt tatttttcta aatacattca aatatgtatc cgctcatgag 8340
acaataaccc tgataaatgc ttcaataata ttgaaaaagg aagagtatga gtattcaaca 8400
tttccgtgtc gcctttatct ccttttttgc ggcattttgc cttcctggtt ttgctcacc 8460
agaaacgctg gtgaaagtaa aagatgctga agatcagttg ggtgcacgag tgggttacat 8520
cgaactggat ctcaacagcg gtaagatcct tgagagtttt cggcccgagg aacgttttcc 8580
aatgatgagc acttttaag ttctgctatg tggcgcggtt ttatccgta ttgacgcgg 8640
gcaagagcaa ctcggtcgcc gcatacacta ttctcagaat gacttggttg agtactcacc 8700
agtcacagaa aagcatctta cggatggcat gacagtaaga gaattatgca gtgctgccat 8760
aaccatgagt gataacactg cggccaactt acttctgaca acgatcggag gaccgaagga 8820
gctaaccgct tttttgcaca acatggggga tcatgtaact cgccttgatc gttgggaacc 8880
ggagctgaat gaagccatac caaacgacga gcgtgacacc acgatgcctg tagcaatggc 8940
aacaacgctg cgcaaaactat taactggcga actacttact ctacttccc ggcaacaatt 9000
aatagactgg atggaggcgg ataaagtgc aggaccaett ctgcgctcgg cccttcggc 9060
tggtggtttt attgctgata aatctggagc cggtagcggt tatctcgcg gtatcattgc 9120
agcactgggg ccagatggta agcctcccg tatcgtagtt atctacacga cggggagtca 9180
ggcaactatg gatgaacgaa atagacagat cgctgagata ggtgcctcac tgattaagca 9240
ttggttaactg tcagaccaag tttactcata tatactttag attgatttaa aacttcattt 9300
ttaatttaaa aggatctagg tgaagatcct ttttgataat ctcatgacca aaatccctta 9360
acgtgagttt tcgttccact gagcgtcaga ccccgtagaa aagatcaaag gatcttcttg 9420
agatcccttt tttctgcgcg taatctgctg cttgcaaca aaaaaaccac cgctaccagc 9480
ggtggtttgt ttgccggatc aagagctacc aactcttttt ccgaaggtaa ctggcttcag 9540
cagagcgcat ataccaaata ctgttcttct agttagcggt tagttaggcc accacttcaa 9600
gaactctgta gcaccgccta catacctcgc tctgctaata ctgttaccag tggctgctgc 9660
cagtggcgat aagtcgtgtc ttaccgggtt ggactcaaga cgatagttac cggataaggc 9720
gcagcggtcg ggctgaacgg ggggttcgtg cacacagccc agcttggagc gaacgaccta 9780
caccgaactg agatacctac agcgtgagct atgagaaagc gccacgcttc ccgaaggagg 9840
aaaggcggac aggtatccgg taagcggcag ggtcggaaca ggagagcgca cgaggaggct 9900
tccaggggga aacgcctggt atctttatag tctgtcggg tttcgccacc tctgacttga 9960
gcgtcgattt ttgtgatgct cgtcaggggg cgggagccta tggaaaaacg ccagcaacgc 10020
ggccttttta cggttcctgg ccttttgcgt gccttttgct cacatgttct ttcctgcgtt 10080
atccctgat tctgtggata accgtattac cgcttttgag tgagctgata ccgctcgccg 10140
cagccgaacg accgagcgca gcgagtcagt gagcgaggaa gcggaagagc gcccaatacg 10200
caaaccgcct cttcccgcg gttggccgat tcattaatgc agctggcacg acaggtttcc 10260
cgactggaaa gcgggcagtg agcgcaacgc aattaatgtg agttagctca ctcattaggc 10320
accccaggct ttacacttta tgcttcggc tcgtatgttg tgtggaattg tgagcggata 10380
acaatttcac acaggaaaca gctatgacca tgattacgcc aagcgcgcaa ttaaccctca 10440
ctaaggga caaaagctgg agctgcaagc tt 10472

```

<210> 49

<211> 8889

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =

synthetic construct

<400> 49

```

ttggcccatg gcatacgttg tatccatata ataatatgta catttatatt ggctcatgtc 60
caacattacc gccatgttga cattgattat tgactagtta ttaatagtaa tcaattacgg 120
ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg gtaaatggcc 180
cgcttggtg accgccaac gacccccgcc cattgacgtc aataatgacg tatgttccca 240
tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta cggtaaactg 300
cccacttggt agtacatcaa gtgtatcata tgccaagtac gccccctatt gacgtcaatg 360
acggtaaatg gcccgcctgg cattatgccc agtacatgac cttatgggac tttcctactt 420
ggcagtagat ctacgtatta gtcacgtcta ttaccatggg gatgcggttt tggcagtaca 480
tcaatgggag tggatagcgg tttgactcac ggggatttcc aagtctccac cccattgacg 540
tcaatgggag tttgttttgg caccaaaatc aacgggactt tccaaaatgt cgtaacaact 600
ccgccccatt gtcgcaaatg ggcggtaggc gtgtacgggt ggaggtctat ataagcagag 660
ctcgtttagt gaaccgtcag atcgctgga gacgccatcc acgctgtttt gacctccata 720
gaagacaccg ggaccgatcc agcctcccct cgaagcttac atgtggtacc gagctcggat 780
cctgagaact tcaggttgag tctatgggac ccttgatgtt ttctttcccc ttcttttcta 840
tggttaagtt catgtcatag gaaggggaga agtaacaggg tacacatatt gaccaaata 900
gggtaatttt gcatttgtaa ttttaaaaaa tgctttcttc ttttaataata cttttttgtt 960
tatcttattt ctaatacttt ccctaacttc tttctttcag ggcaataatg atacaatgta 1020
tcatgcctct tttgaccatt ctaagaata acagtgtataa tttctgggtt aaggcaatag 1080
caatatttct gcataataat atttctgcat ataaattgta actgatgtaa gaggtttcat 1140
attgctaata gcagctacaa tccagctacc attctgcttt tattttatgg ttgggataag 1200
gctggattat tctgagtcca agctaggccc ttttgctaata catgttcata cctcttatct 1260
tctcccaca gctcctgggc aacgtgctgg tctgtgtgct ggcccatcac tttggcaaag 1320
cacgtgagat ctgaattcga gatctgccgc cgccatgggt gcgagagcgt cagtattaag 1380
cggggggaga ttagatcgat gggaaaaaat tcggttaagg ccagggggaa agaaaaata 1440
taaattaaaa catatagtat gggcaagcag ggagctagaa cgattcgcag ttaatcctgg 1500
cctgttagaa acatcagaag gctgtagaca aatactggga cagctacaac catcccttca 1560
gacaggatca gaagaactta gatcattata taatacagta gcaacctctt attgtgtgca 1620
tcaaaggata gagataaaaag acaccaagga agctttagac aagatagagg aagagcaaaa 1680
caaaagtaag aaaaaagcac agcaagcagc agctgacaca ggacacagca atcagggtcag 1740
ccaaaattac cctatagtgc agaacatcca ggggcaaata gtacatcagg ccatatcacc 1800
tagaacttta aatgcatggg taaaagtagt agaagagaag gctttcagcc cagaagtgat 1860
acccatgttt tcagcattat cagaaggagc caccaccaca gatttaaaac ccatgctaaa 1920
cacagtgggg ggacatcaag cagccatgca aatgttaaaa gagaccatca atgaggaagc 1980
tgcaaatgg gatagagtgc atccagtgc tgagggcct attgcaccag gccagatgag 2040
agaaccaagg ggaagtgaca tagcaggaac tactagtacc cttcaggaac aaataggatg 2100
gatgacacat aatccacctt tcccagtagg agaatctat aaaagatgga taatcctggg 2160
attaaataaa atagtaagaa tgtatagccc taccagcatt ctggacataa gacaaggacc 2220
aaaggaaccc tttagagact atgtagaccg attctataaa actctaagag ccgagcaagc 2280
ttcacaagag gtaaaaaatt ggatgacaga aacctgtgtg gtccaaaatg cgaaccaga 2340
ttgtaagact attttaaaag cattgggacc aggagcgaca ctagaagaaa tgatgacag 2400
atgtcagggg gtggggggac ccggccataa agcaagagtt ttggctgaag caatgagcca 2460
agtaacaaat ccagctacca taatgatata gaaaggcaat tttaggaacc aaagaaagac 2520
tgttaagtgt ttcaattgtg gcaaagaagg gcacatagcc aaaaattgca gggcccctag 2580
gaaaagggg tggtgaaat gtggaaagga aggacaccaa atgaaagatt gtactgagag 2640
acaggctaata tttttaggga agatctggcc tcccacaag ggaaggccag ggaattttct 2700
tcagagcaga ccagagccaa cagccccacc agaagagagc ttcagggttg ggaagagac 2760
aacaactccc tctcagaagc aggagccgat agacaaggaa ctgtatcctt tagcttccct 2820
cagatcactc tttggcagcg accctcgtc acaataaaga taggggggca attaaaggaa 2880
gctctattag atacaggagc agatgatata gtattagaag aaatgaattt gccaggaaga 2940
tggaaccaa aaatgatagg ggggaattgga ggttttatca aagtaagaca gtatgatcag 3000
atactcatag aaatctgcgg acataaagct ataggtacag tattagtagg acctacacct 3060
gtcaacataa ttggaagaaa tctgttgact cagattgggt gcactttaaa ttttccatt 3120
agtccatttg agactgtacc agtaaaatta aagccaggaa tggatggccc aaaagttaa 3180
caatggccat tgacagaaga aaaaataaaa gcattagtag aaatttgtac agaaatggaa 3240
aaggaaggaa aaatttcaa aattgggcct aaaaatccat acaatactcc agtatttgc 3300
ataaagaaaa aagacagtac taaatggaga aaatttagtag atttcagaga acttaataag 3360
agaactcaag atttctggga agttcaatta ggaataccac atcctgcagg gttaaaacag 3420
aaaaaatcag taacagtact ggatgtgggc gatgcatatt tttcagttcc cttagataaa 3480

```


gacttcagga	agtatactgc	atttaccata	cctagtataa	acaatgagac	accagggatt	3540
agatatcagt	acaatgtgct	tccacagggg	tggaaaggat	caccagcaat	attccagtg	3600
agcatgacaa	aaatcttaga	gccttttaga	aaacaaaatc	cagacatagt	catctatcaa	3660
tacatggatg	atthgtatgt	aggatctgac	ttagaaatag	ggcagcatag	aacaaaaata	3720
gaggaactga	gacaacatct	gttgaggtgg	ggattttacca	caccagacaa	aaaacatcag	3780
aaagaacctc	cattcccttg	gatgggttat	gaactccatc	ctgataaatg	gacagtacag	3840
cctatagtgc	tgccagaaaa	ggacagctgg	actgtcaatg	acatacagaa	attagtggga	3900
aaattgaatt	gggcaagtca	gattttatgca	gggattaaag	taaggcaatt	atgtaaactt	3960
cttaggggaa	ccaaagcact	aacagaagta	gtaccactaa	cagaagaagc	agagctagaa	4020
ctggcagaaa	acagggagat	tctaaaagaa	ccggtacatg	gagtgtatta	tgacccatca	4080
aaagacttaa	tagcagaaat	acagaagcag	gggcaaggcc	aatggacata	tcaaatttat	4140
caagagccat	ttaaaaatct	gaaaaacagga	aagtatgcaa	gaatgaagg	tgccacact	4200
aatgatgtga	aacaattaac	agaggcagta	caaaaaatag	ccacagaaag	catagtaata	4260
tggggaaaaga	ctcctaaatt	taaattaccc	atacaaaagg	aaacatggga	agcatgggtg	4320
acagagtatt	ggcaagccac	ctggattcct	gagtgggagt	ttgtcaatac	ccctccctta	4380
gtgaagtatt	ggtaccagtt	agagaaagaa	cccataatag	gagcagaaac	tttctatgta	4440
gatggggcag	ccaataggga	aactaaatta	ggaaaagcag	gatatgtaac	tgacagagga	4500
agacaaaaag	ttgtccccct	aacggacaca	acaaatcaga	agactgagtt	acaagcaatt	4560
catctagctt	tgccaggattc	gggattagaa	gtaaacatag	tgacagactc	acaatatgca	4620
ttgggaatca	ttcaagcaca	accagataag	agtgaatcag	agttagtcag	tcaaataata	4680
gagcagttaa	taaaaaagga	aaaagtctac	ctggcatggg	taccagcaca	caaaggaatt	4740
ggaggaatag	aacaagtaga	taaattgggtc	agtgtctgaa	tcaggaaaagt	actattttta	4800
gatggaatag	ataaggccca	agaagaacat	gagaaatata	acagtaattg	gagagcaatg	4860
gctagtgtat	ttaacctacc	acctgtagta	gcaaaagaaa	tagtagccag	ctgtgataaa	4920
tgctagctaa	aaggggaagc	catgcatgga	caagtagact	gtagcccagg	aatatggcag	4980
ctagattgta	cacattttaga	aggaaaagtt	atcttggtag	cagttcatgt	agccagtggg	5040
tatatagaag	cagaagtaat	tcacagcag	acagggcaag	aaacagcata	cttcctctta	5100
aaattagcag	gaagatggcc	agtaaaaaca	gtacatacag	acaatggcag	caatttcacc	5160
agactacag	ttaaggccgc	ctgttgggtg	gcggggatca	agcaggaatt	tgccattccc	5220
tacaatcccc	aaagtcaagg	agtaatagaa	tctatgaata	aagaattaaa	gaaaattata	5280
ggacaggtaa	gagatcaggc	tgaacatctt	aagacagcag	tacaaatggc	agtattcatc	5340
cacaatttta	aaagaaaagg	ggggattggg	gggtacagtg	caggggaaaag	aatagtagac	5400
ataatagcaa	cagacataca	aactaaagaa	ttacaaaaac	aaattacaaa	aattcaaaat	5460
tttcgggttt	attacaggga	cagcagagat	ccagtttgga	aaggaccagc	aaagctcctc	5520
tggaaaggtg	aaggggcagt	agtaatacaa	gataatagt	acataaaagt	agtgccaa	5580
agaaaagcaa	agatcatcag	ggattatgga	aaacagatgg	cagggtgatga	ttgtgtggca	5640
agtagacagg	atgaggatta	acacatggaa	ttccggagcg	gccgcaggag	ctttgttctc	5700
tgggttcttg	ggagcagcag	gaagcactat	gggcccagcg	tcaatgacgc	tgacggtaca	5760
ggccagacaa	ttattgtctg	gtatagtgca	gcagcagaac	aatttgctga	gggctattga	5820
ggcgcaacag	catctgttgc	aactcacagt	ctggggcatc	aagcagctcc	aggcaagaat	5880
cctggctgtg	gaaagatacc	taaaggatca	acagctcctg	gggattttgg	gttgcctctg	5940
aaaactcatt	tgccaccactg	ctgtgccttg	gaatgctagt	tggagtaata	aatctctgga	6000
acagatttgg	aatcacacga	cctggatgga	gtgggacaga	gaaattaaca	attacacaag	6060
cttcgcgga	attcacccca	ccagtgcagg	ctgcctatca	gaaagtgggtg	gctggtgtgg	6120
ctaattgccct	ggccccacaag	tatcactaag	ctgcctttct	tgctgtccaa	tttctattaa	6180
aggttccttt	gttccctaag	tccaactact	aaactggggg	atattatgaa	gggccttgag	6240
catctggatt	ctgcctaata	aaaaacattt	atthtcattg	caatgatgta	tttaaatat	6300
ttctgaatat	tttactaaaa	agggaatgtg	ggaggtcagt	gcatttaaaa	cataaagaaa	6360
tgaagagcta	gttcaaacct	tgggaaaata	cactatatct	taaactccat	gaaagaaggt	6420
gaggctgcaa	acagctaatg	cacattggca	acagccctg	atgcctatgc	cttattcatc	6480
cctcagaaaa	ggattcaagt	agaggcttga	tttgagggtt	aaagttttgc	tatgctgtat	6540
tttacattac	ttattgtttt	agctgtcctc	atgaatgtct	tttactacc	catttgctta	6600
tcctgcatct	ctcagccttg	actccactca	gttctcttgc	ttagagatac	cacctttccc	6660
ctgaagtgtt	ccttccatgt	tttacggcga	gatggtttct	cctgcctgg	ccactcagcc	6720
ttagttgtct	ctgttgtctt	atagaggtct	acttgaagaa	ggaaaaacag	ggggcatgg	6780
ttgactgtcc	tgtgagccct	tcttccctgc	ctccccact	cacagtgacc	cggaaatccct	6840
cgacatggca	gtctagcact	agtgcggccg	cagatctgct	tcctcgctca	ctgactcgct	6900
gcgctcggtc	gttcggctgc	ggcgagcgg	atcagctcac	tcaaaggcgg	taatacgggt	6960
atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	gcaaaaggcc	agcaaaaggc	7020
caggaaccgt	aaaaaggccg	cgttgctggc	gtttttccat	aggctccgcc	cccctgacga	7080
gcatcacaaa	aatcgacgct	caagtcagag	gtggcgaaac	ccgacaggac	tataaagata	7140

```

ccaggcggttt ccccttgga gctccctcgt gcgctctcct gttccgaccc tgccgcttac 7200
cggatacctg tccgcctttc tocccttcggg aagcggtggcg ctttctcata gctcacgctg 7260
taggtatctc agttcggtgt aggtcggtcg ctccaagctg ggctgtgtgc acgaaccccc 7320
cggttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca acccggttaag 7380
acacgactta tcgccactgg cagcagccac tggtaacagg attagcagag cgaggatagt 7440
aggcgggtgct acagagttct tgaagtgggt gcctaactac ggctacacta gaagaacagt 7500
atctgggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg 7560
atccggcaaaa caaacaccg ctggtagcgg tgggtttttt gtttgcaagc agcagattac 7620
gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacggggt ctgacgctca 7680
gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac 7740
ctagatcctt ttaaattaaa aatgaagttt taaatcaatc taaagtatat atgagtaaac 7800
ttgggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt 7860
tcgttcaccc atagttgcct gactcctcgt cgtgtagata actacgatac gggagggctt 7920
accatctggc cccagtctgc caatgatacc gcgagaccca cgctcacccg ctccagattt 7980
atcagcaata aaccagccag ccggaagggc cgagcgaga agtggtcctg caactttatc 8040
cgctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccagttaa 8100
tagtttgccg aacgttgttg ccattgctac aggcacgtg gtgtcacgct cgctggttg 8160
tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat ccccatggt 8220
gtgcaaaaaa gcggttagct ccttcgggtcc tccgatcgtt gtcagaagta agttggccgc 8280
agtgttatca ctcatggtta tggcagcact gcataattct cttactgtca tgccatccgt 8340
aagatgcttt tctgtgactg gtgagtactc aaccaagtca ttctgagaat agtgtagcg 8400
gcgaccgagt tgctcttgcc cggcgtaaat acgggataat accgcgccac atagcagaac 8460
tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc 8520
gctgttgaga tccagttcga tgtaaccac tcgtgcaccc aactgatctt cagcatcttt 8580
tactttcacc agcgtttctg ggtgagcaaa aaacaggaagg caaaatgccg caaaaaagg 8640
aataagggcg acacggaat gttgaatact catactcttc ctttttcaat attattgaag 8700
catttatcag ggttattgtc tcatgagcgg atacatattt gaatgtattt agaaaaataa 8760
acaaataggg gttccgcgca catctcccg aaaagtgcc cctgacggga tccctgagg 8820
gggccccat gggctagagg atccggcctc ggctctgca taaataaaaa aaattagtca 8880
gccatgagc

```

<210> 50

<211> 4180

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 50

```

aatgtagtct tatgcaatac tcttgtagtc ttgcaacatg gtaacgatga gttagcaaca 60
tgctttacaa ggagagaaaa agcaccgtgc atgccgattg gtggaagtaa ggtggtagca 120
tcgtgcctta ttaggaaggc aacagacggg tctgacatgg attggacgaa ccaactgaatt 180
ccgcattgca gagatattgt atttaagtgc ctgactcgat acaataaacg ccatttgacc 240
attcaccaca ttggtgtgca cctccaagct cgagctcggt tagtgaaccg tcagatcgcc 300
tgagagcgcc atccacgctg ttttgacctc catagaagac accgggaccg atccagcctc 360
ccctcgaagc tagtcgatta ggcattctct atggcaggaa gaagcgaga cagcgacgaa 420
gacctcctca aggcagtcag actcatcaag tttctctatc aaagcaaccc acctcccaat 480
cccgagggga cccgacaggc ccgaaggaaat agaagaagaa ggtggagaga gagacagaga 540
cagatccatt cgattagtga acggatcctt agcacttata tgggacgata tgggagcct 600
gtgcctcttc agctaccacc gcttgagaga cttactcttg attgtaacga ggattgtgga 660
acttctggga cgcagggggg ggggaagcct caaatattgg tggaatctcc tacaatattg 720
gagtcaggag ctaaaagaata gtgctgttag cttgctcaat gccacagcta tagcagtagc 780
tgaggggaca gatagggtta tagaagtagt acaagaagct tggcactggc cgctggttta 840
caacgtcgtg atctgagcct gggagatctc tggctaacta ggaacccac tgcttaagcc 900
tcaataaagc ttgccttgag tgcttcaagt agtggtgccc cgtctgttgt gtgactctgg 960
taactagaga tcaggaaaac cctggcgtaa cctgaactta tcgccttgca gcacatcccc 1020
ctttcgccag ctggcgtaat agcgaagagg cccgcaccca tcgcccttcc caacagttgc 1080

```

```

gcagcctgaa tggcgaatgg cgctgatgc ggtattttct ccttacgcat ctgtgcggtg 1140
tttcacaccg catacgtcaa agcaaccata gtacgcgccc tgtagcgggc cattaagcgc 1200
ggcgggtgtg gtggttacgc gcagcgtgac cgctacactt gccagcgccc tagcgccccg 1260
tcctttcgct ttcttccctt cctttctcgc caggttcgcc ggctttcccc gtcaagctct 1320
aaatcggggg ctcccttttag ggttccgatt tagtgcttta cggcacctcg accccaaaaa 1380
acttgatttg ggtgatgggt cacgtagtgg gccatcgccc tgatagacgg tttttcgccc 1440
tttgacgttg gagtccacgt tctttaatag tggactcttg ttccaaactg gaacaacact 1500
caaccctatc tcgggctatt cttttgattt ataagggatt ttgccgattt cggcctattg 1560
gttaaaaaat gagctgattt aacaaaaatt taacgcgaat ttaacaaaaa tattaacggt 1620
tacaatttta tgggtgcactc tcagtacaat ctgctctgat gccgcatagt taagccagcc 1680
ccgacaccgg ccaacaccgg ctgacgcgcc ctgacgggct tgtctgctcc cggcatccgc 1740
ttacagacaa gctgtgacgg tctccgggag ctgcatgtgt cagaggtttt caccgtcatc 1800
accgaaacgc gcgagacgaa agggcctcgt gatacgcta tttttatagg ttaatgtcat 1860
gataataatg gtttcttaga cgtcaggtgg cacttttcgg ggaaatgtgc gcggaacccc 1920
tatttgttta tttttctaaa tacattcaaa tatgtatccg ctcatgagac aataaccctg 1980
ataaatgctt caataatatt gaaaaaggaa gagtatgagt attcaacatt tccgtgtcgc 2040
ccttattccc ttttttcggg cattttgcct tcctgttttt gctcaccag aaacgctggg 2100
gaaagtaaaa gatgctgaag atcagttggg tgcacgagtg ggttacatcg aactggatct 2160
caacagcggg aagatccttg agagttttcg ccccgaaaga cgttttccaa tgatgagcac 2220
ttttaaggtt ctgctatgtg gcgcggtatt atcccgatg gacgcggggc aagagcaact 2280
cggctcgccg atacactatt ctccagaatga cttggttgag tactcaccag tcacagaaaa 2340
gcactcttaag gatggcctga cagtaagaga attatgcagt gctgccataa ccatgagtga 2400
taacactcgg gccaaactac ttctgacaac gatcggagga ccgaaggagc taaccgcttt 2460
tttgacacaac atgggggatc atgtaactcg ccttgatcgt tgggaaccgg agctgaatga 2520
agccatacca aacgacgagc gtgacaccac gatgcctgta gcaatggcaa caacgttgcg 2580
caaactatta actggcgaac tacttactct agcttcccgg caacaattaa tagactggat 2640
ggaggcggat aaagttgcag gaccacttct gcgctcggcc cttccggctg gctggtttat 2700
tgctgataaa tctggagccg gtgagcgtgg gtctcgcggt atcattgcag cactggggcc 2760
agatggtaag cctcccgta tcgtagttag ctacacgacg gggagtcagg caactatgga 2820
tgaacgaaat agacagatcg ctgagatagg tgcctcactg attaagcatt ggtaactgtc 2880
agaccaagtt tactcatata tactttagat tgatttaaaa cttcattttt aatttaaaag 2940
gatctaggtg aagatccttt ttgataatct catgaccaa atcccttaac gtgagttttc 3000
gttccactga gcgtcagacc ccgtagaaaa gatcaaagga tcttcttgag atcctttttt 3060
tctgcgcgta atctgctgct tgcaaacaaa aaaaccaccg ctaccagcgg tggtttgttt 3120
gccggatcaa gagctaccaa ctctttttcc gaaggtaact ggcttcagca gagcgcagat 3180
accaaataca gttcttctag tgtagccgta gttaggccac cacttcaaga actctgttagc 3240
accgcctaca tacctcgctc tgctaatact gttaccagtg gctgctgcca gtggcgataa 3300
gtcgtgtctt accgggttg actcaagacg atagttaccg gataaggcgc agcggtcggg 3360
ctgaacgggg ggttcgtgca cacagcccag cttggagcga acgacctaca ccgaactgag 3420
atacctacag cgtgagctat gagaaagcgc cacgcttccc gaaggagaa aggcggacag 3480
gtatccggta agcggcaggg tcggaacagg agagcgcacg agggagcttc cagggggaaa 3540
cgctggtat ctttatagtc ctgctcgggt tcgccacctc tgacttgagc gtcgattttt 3600
gtgatgctcg tcaggggggc ggagcctatg gaaaaacgcc agcaacgcgg cctttttacg 3660
gttccctggc ttttgcctgc ctttgcctca catgttcttt cctgcgttat ccctgattc 3720
tgttgataac cgtattaccg cctttgagtg agctgatacc gctcgcgcga gccgaacgac 3780
cgagcgcagc gagtcagtga gcgaggaagc ggaagagcgc ccaatacgca aaccgcctct 3840
ccccgcgcgt tggccgattc attaatgcag ctggcacgac aggtttcccg actggaagc 3900
gggcagtgag cgcaacgcaa ttaatgtgag ttagctcact cattaggcac ccaggcttt 3960
acactttatg cttccggctc gtatgttgtg tggaaattgt agcggataac aatttcacac 4020
aggaaacagc tatgacatga ttacgaattc gatgtacggg ccagatatac gcgtatctga 4080
ggggactagg gtgtgttttag gcgaaaagcg gggcttcggg tgtacgcggg taggagctcc 4140
ctcaggatat agtagtttcg cttttgcata gggaggggga 4180

```

<210> 51

<211> 5821

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: /note =
synthetic construct.

<400> 51

```

ttggcccatt gcatacgttg tatccatata ataatatgta catttatatt ggctcatgtc 60
caacattacc gccatgttga cattgattat tgactagtta ttaatagtaa tcaattacgg 120
ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg gtaaatggcc 180
cgcttggtcg accgcccac gacccccgcc cattgaögtc aataatgacg tatgttccca 240
tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta cggtaaactg 300
cccacttggc agtacatcaa gtgtatcata tgccaagtac gccccctatt gacgtcaatg 360
acggtaaatg gcccgctgg cattatgccc agtacatgac cttatgggac tttcctactt 420
ggcagtacat ctacgtatta gtcacgcta ttaccatggg gatgcgggtt tggcagtaca 480
tcaatgggog tggatagcgg tttgactcac ggggatttcc aagtctccac ccattgacg 540
tcaatgggag tttgttttgg caccaaaatc aacgggactt tccaaaatgt cgtaacaact 600
ccgccccatt gacgcaaatg ggcggtaggc gtgtacgggt ggagggtctat ataagcagag 660
ctcgtttagt gaaccgtcag atcgctcgga gacgccatcc acgctgtttt gacctccata 720
gaagacaccg ggaccgatcc agcctcccct cgaagcttac atgtggtacc gagctcggat 780
cctgagaact tcagggtagg tctatgggac ccttgatgtt ttttttcccc ttttttcta 840
tggttaagtt catgtcatag gaaggggaga agtaacaggg tacacatatt gaccaaacta 900
gggtaatttt gcatttgtaa ttttaaaaaa tgctttcttc ttttaataata cttttttgtt 960
tatcttattt ctaatacttt ccctaacttc tttctttcag ggcaataatg atacaatgta 1020
tcatgcctct ttgcaccatt ctaaagaata acagtgataa tttctgggtt aaggcaatag 1080
caatatttct gcatataaat atttctgcac ataatgtga actgatgtaa gaggtttcat 1140
attgctaata gcagctacaa tccagctacc attctgcttt tattttatgg ttgggataag 1200
gctggattat tctgagtcca agctaggccc ttttgctaata catgttcata cctcttatct 1260
tctcccaca gtcctggggc aacgtgctgg tctgtgtgct ggcccatcac tttggcaaag 1320
cacgtgagat ctgaattctg acactatgaa gtgccttttg tacttagcct ttttattcat 1380
tggggtgaat tgcaagttca ccatagtttt tccacacaac caaaaaggaa actggaaaaa 1440
tgttccctct aattaccatt attgcccgtc aagctcagat ttaaattggc ataatgactt 1500
aataggcaca gccttacaag tcaaaatgcc caagagtcac aaggctattc aagcagacgg 1560
ttggatgtgt catgcttcca aatgggtcac tacttgtgat ttccgctggg atggaccgaa 1620
gtatataaca cattccatcc gatccttcac tccatctgta gaacaatgca aggaaagcat 1680
tgaacaaacg aaacaaggaa ctgggctgaa tccaggett cctcctcaaa gttgtggata 1740
tgcaactgtg acggatgccg aagcagtgat tgtccagggt actcctcacc atgtgctggg 1800
tgatgaatac acaggagaat ggggtgattc acagtccatc aacggaaaaat gcagcaatta 1860
catatgcccc actgtccata actctacaac ctggcattct gactataagg tcaaagggct 1920
atgtgattct aacctcattt ccatggacat caccttcttc tcagaggacg gagagctatc 1980
atccctggga aaggagggca cagggttcag aagtaactac tttgcttatg aaactggagg 2040
caaggcctgc aaaatgcaat actgcaagca ttggggagtc agactcccat cagggtctctg 2100
gttcogagat gctgataagg atctctttgc tgcagccaga ttccctgaat gcccagaagg 2160
gtcaagtatc tctgctccat ctcagacctc agtgatgta agtctaattc aggacgttga 2220
gaggatcttg gattattccc tctgccaaaga aacctggagc aaaatcagag cgggtcttcc 2280
aatctctcca gtggatctca gctatcttgc tcttaaaaaa ccaggaaccg gtccctgcttt 2340
caccataatc aatggtaccc taaaataact tgagaccaga tacatcagag tcatatttgc 2400
tgctccaatc ctctcaagaa tggtcggaat gatcagtgga actaccacag aaagggaact 2460
gtgggatgac tgggcaccat atgaagacgt ggaattgga cccaatggag ttctgaggac 2520
cagttcagga tataagtttc ctttatacat gattggacat ggtatgttgg actccgatct 2580
tcatcttagc tcaaaggctc aggtgttcga acatcctcac attcaagacg ctgcttcgca 2640
acttctgat gatgagagtt ttttttttgg tgatactggg ctatccaaaa atccaatcga 2700
gctttagaaa ggttggttca gtagttggaa aagctctatt gcctcttttt tctttatcat 2760
agggttaatc attggactat tcttggttct ccgagttggg atccatcttt gcattaaatt 2820
aaagcacacc aagaaaagac agatttatac agacatagag atgaaccgac ttggaaagta 2880
actcaaatcc tgcacaacag attcttcatg tttggacca atcaacttgt gataccatgc 2940
tcaaagaggg ctcaattata tttgagtttt taatttttat gaaaaaaaacg 3000
gaattcacc caccagtgcg ggctgcctat cagaaagtgg tggctggtgt ggctaattgc 3060
ctggcccaca agtatcacta agctcgcttt cttgctgtcc aatttctatt aaaggttcct 3120
ttgttcctta agtccaacta ctaaactggg ggatattatg aagggccttg agcatctgga 3180
ttctgcctaa taaaaaacat ttattttcat tgcaatgatg tatttaaat atttctgaat 3240
attttactaa aaagggaatg tgggaggtca gtgcatttaa aacataaaga aatgaagagc 3300
tagttcaaac cttgggaaaa tacactatat cttaaactcc atgaaagaag gtgagggtgc 3360
aaacagctaa tgcacattgg caacagcccc tgatgcctat gccttattca tccctcagaa 3420
aaggattcaa gtagaggctt gatttggagg ttaaagtttt gctatgctgt attttaccat 3480
acttattgtt ttagctgtcc tcatgaatgt ctttctacta cccatttgc tctctgcac 3540

```

```

ctctcagcct tgactccact cagttctctt gcttagagat accaccttcc cctgaagtg 3600
ttccttccat gttttacggc gagatggttt ctctctgcct ggccactcag ccttagttgt 3660
ctctgttgct ttatagaggt ctacttgaag aaggaaaaac agggggcatg gtttgactgt 3720
cctgtgagcc cttcttccct gcctccccc ctcacagtga cccggaatcc ctgcacatgg 3780
cagttctagca ctagtgcggc cgcagatctg ctctctcgct cactgactcg ctgcgctcgg 3840
tcgttcggct gcggcgagcg gtatcagctc actcaaaggc ggtaatacgg ttatccacag 3900
aatcagggga taacgcagga aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc 3960
gtaaaaaggc cgcgttgctg cgcgttttcc ataggctccg ccccccctgac gagcatcaca 4020
aaaatcgacg ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt 4080
ttccccctgg aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc 4140
tgtccgcctt tctcccttcg ggaagcgtgg cgctttctca tagctcacgc ttaggtatc 4200
tcagttcggg gtaggctcgt cgctccaagc tgggctgtgt gcacgaacct ccggttcagc 4260
ccgaccgctg cgctttatcc ggtaactatc gtcttgagtc caaccggta agacacgact 4320
tatcgccact ggcagcagcc actggtaaca ggattagcag agcgaggat gtaggcgggtg 4380
ctacagagtt cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttggtg 4440
tctgcgctct gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca 4500
aacaaccac cgtggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa 4560
aaaaaggatc tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaaacg 4620
aaaactcacg ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc 4680
ttttaatta aaaatgaagt tttaaatcaa cttaaagtat atatgagtaa acttggctcg 4740
acagttacca atgcttaatc agtgaggcac ctatctcagc gatctgtcta ttcggttcgt 4800
ccatagttgc ctgactcccc gtctgttaga taactacgat acgggagggc ttaccatctg 4860
gccccagtgc tgcaatgata ccgcgagacc cagctcacc ggctccagat ttatcagcaa 4920
taaaccagcc agccggaagg gccgagcgca gaagtgttcc tgcaacttta tccgcctcca 4980
tccagctctat taattgttgc cgggaagcta gagtaagtag ttccgaggtt aatagtttgc 5040
gcaacgttgt tgccattgct acaggcatcg tgggtgtcacg ctctctcgtt ggtaggtt 5100
cattcagctc cggttcccaa cgatcaaggc gattacatg atcccccatg ttgtgcaaaa 5160
aagcgttag cctcttcggt cctccgacgt ttgtcagaag taagttggcc gcagtgttat 5220
cactcatggt tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct 5280
tttctgtgac tgggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga 5340
gttgctcttg cccggcgctc atacgggata ataccgccc acatagcaga actttaaaag 5400
tgctcatcat tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga 5460
gatccagttc gatgtaacct actcgtgcac ccaactgatc ttcagcatct tttactttca 5520
ccagcgtttc tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg 5580
cgacacggaa atgttgaata ctcatctct tccttttca atattattga agcatttatc 5640
agggttattg ttcctatgac ggatacatct ttgaatgtat ttagaaaaat aaacaaatag 5700
gggttcgcg cacttttccc cgaaaagtgc cacctgacgg gatccctga gggggcccc 5760
atgggctaga ggatccggcc tcggcctctg cataaataaa aaaaattagt cagccatgag 5820
c 5821

```

<210> 52

<211> 1986

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 52

```

tcctcctctg ctgcaccaat ccttccaacc ccctatggtg gtatggctga cacagaaaat 60
gtctgtcctt gtatgggaca tttgcccctc ttctccaaat ataagacagg atgaggccta 120
gcttttgctg ctccaaagt tttaaagaac acattgcacg gcatttaggg actctaaagg 180
gtggaggagg aatgaggga ttgcatcatg ccaaggctgg tcctcatcca tcaactgttc 240
cagggccag agtggttcc aggaggtatt cttacaaagg aagcccgatc ttagtagtaac 300
actcagagcc cattttcctg cgtaaacccc tcccgacctc atatacagga gtaacatgat 360
cagtgcagct ggggctcgtg ccaaactgcg ggacctgccc aagctgagg ccttgggtgc 420
gctggacaac cctgtgctg atgagactga ctaccgccag gaggccctgg tgcagatggc 480
acacctagag cgcttagaca aagagtacta tgaggacgag gaccgggcag aagctgagga 540

```

```

gatccgacag aggctgaagg aggaacagga gcaagaactc gacccggacc aagacatgga 600
accgtacctc ccgccaactt agtggtcct ctagcctgca gggacagtaa aggtgatggc 660
aggaaggcag ccccgagg agtcaaaggctg ggcacgcggg aggagaggcc agagtacagag 720
gctgcgggta tctcagatat gaaggaaaga tgagagaggc tcaggaagag gtaagaaaag 780
acacaagaga ccagagaagg gagaagaatt agagaggagg gcagaggacc gctgtctcta 840
cagacatagc tggtagagac tgggaggaag ggatgaaccc tgagcgcagt aaggggaagga 900
ggtggctggt ggtatatgga ggatgtagct gggccaggga aaagatcctg cactaaaaat 960
ctgaagctaa aaataacagg acacgggggtg gagaggcgaa aggagggcag attgaggcag 1020
agagactgag aggcctgggg atgtgggcat tccggtaggg cacacagttc acttgtcttc 1080
tctttttcca ggaggccara gatgctgacc tcaagaactc ataatacccc agtggggacc 1140
accgcattca tagccctgtt acaagaagtg ggagatgttc ctttttgtcc cagactggaa 1200
atccattaca tcccagggtc caggttctgt ggtggtcac tctgtgtggc ttgttctgtg 1260
ggcctacctt aagtcctaag cacagctctc aagcagatcc gaggcgacta agatgctagt 1320
aggggttggtc tggagagaag agccgaggag gtgggctgtg atggatcagt tcagctttca 1380
aataaaaagg cgtttttata ttctgtgtcg agttcgtgaa cccctgtggt gggcttctcc 1440
atctgtctgg gttagtacct gccactatac tgggaataagg ggacgcctgc ttccctcgag 1500
ttggctggac aagggtatga gcatccgtgt acttatgggg ttgccagctt ggtcctggat 1560
cgcccgggcc cttccccccac ccgttcgggt cccaccacc acccgcgctc gtacgtgcgt 1620
ctccgcctgc agctcttgac tcatcggggc ccccggtca catgcgctcg ctcggtctta 1680
taggcgcgcg cccctgccca ccccccgcgc gcctggggag ccgcagccgc cgccactcct 1740
gtctctctcg cgcgcgcgc gtcaccacc ccaccgccac cggtgagtc tgcagtcctc 1800
gaggaactga aaaaccagaa agttaactgg taagtttagt ctttttgtct tttatttcag 1860
gtcccgatc cggtggtggt gcaaatacaa gaactgtctc tcagtggatg ttgcctttac 1920
ttctaggcct gtacggaagt gttacttctg ctctaaaagc tgcggaattg taccgcggc 1980
caagct 1986

```

<210> 53

<211> 1610

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
synthetic construct

<400> 53

```

cggatgagcc tctgtgaact actaagggtg gaggggggcta tacgcagagg agaatgtcag 60
atgtctagct cgggtccctc cgctgacgc tcctctctgt ctcagccagg actggtttct 120
gtaagaaaca gcaggagctg tggcagcggc gaaaggaagc ggctgaggcg cttggaaccc 180
gaaaagtctc ggtgctcctg gctacctcgc acagcgtgcc cgcccgccg tcagtaccat 240
ggacagcagc gctgccccca cgaacgccag caattgcact gatgccttgg cgtactcaag 300
ttgtccccca gcaccagcc ccggttcctg ggtcaacttg tccacttag atggcaacct 360
gtccgaccca tgccgtccga accgcaccga cctggggcgg agagacagcc tgtgcccctc 420
gacgggcagt cctcccatga tcacggccat cacgatcatg gccctctact ccatcgtgtg 480
cgtgggtggg ctcttcggaa acttcctggt catgtatgtg attgtcagat acaccaagat 540
gaagactgcc accaaccatct acattttcaa ccttgctctg gcagatgcct tagccaccag 600
taccctgccc ttccagagtg tgaattacct aatgggaaca tggccatttg gaaccatcct 660
ttgcaagata gtgatctcca tagattacta taacatgttc accagcatat tcaccctctg 720
caccatgagt gttgatcgat acattgcagt ctgccaccct gtcaaggcct tagatttccg 780
tactccccga aatgccaaaa ttatcaatgt ctgcaactgg atcctctctt cagccatttg 840
tcttctgtga atgttcattg ctacaacaaa atacaggcaa ggttccatag attgtacact 900
aacattctct catccaacct ggtactggga aaacctgctg aagatctgtg ttttcatctt 960
cgcttcatt atgccagtgc tcatcattac cgtgtgctat ggactgatga tcttgccct 1020
caagagtgtc cgcatgctct ctggctccaa agaaaaggac aggaatcttc gaaggatcac 1080
caggatggtg ctggtggtgg tggctgtgtt catcgtctgc tggactccca ttcacattta 1140
cgtcatcatt aaagccttgg ttacaatccc agaaactacg ttccagactg tttcttggca 1200
cttctgcatt gctctagggt acacaaacag ctgcctcaac ccagtccttt atgcatttct 1260
ggatgaaaac ttcaaacgat gtttcagaga gttctgtatc ccaacctctt ccaacattga 1320
gcaacaaaac tccactcgaa ttctgcagaa cactagagac caccctcca cggccaatac 1380
agtggataga actaatcatc agctagaaaa tctggaagca gaaactgctc cgttgcccta 1440
acagggtctc atgccattcc gaccttcacc aagcttagaa gccaccatgt atgtggaagc 1500

```

aggttgcttc aagaatgtgt aggaggctct aattctctag gaaagtgcct gcttttaggt 1560
 catccaacct ctttcctctc tggccactct gctctgcaca ttagaggccg 1610

<210> 54

<211> 1536

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/note =
 synthetic construct

<400> 54

atgaagtgcc tttgtactt agccttttta ttcattgggg tgaattgcaa gttcaccata 60
 gtttttccac acaacccaaa aggaaactgg aaaaatgttc cttctaatta ccattattgc 120
 ccgtcaagct cagatttaaa ttggcataat gacttaatag gcacagcctt acaagtcaaa 180
 atgccaaga gtcacaaggc tattcaagca gacggttggg tgtgtcatgc ttccaaatgg 240
 gtcactactt gtgatttccg ctggtatgga ccgaagtata taacacattc catccgatcc 300
 ttcactccat ctgtagaaca atgcaaggaa agcattgaac aaacgaaaca aggaacttgg 360
 ctgaatccag gcttccctcc tcaaagtgtg ggatatgcaa ctgtgacgga tgccgaagca 420
 gtgattgtcc aggtgactcc tcaccatgtg ctggttgatg aatacacagg agaatgggtt 480
 gattcacagt tcatcaacgg aaaatgcagc aattacatat gccccactgt ccataactct 540
 acaacctggc attctgacta taaggtaaaa gggctatgtg attctaacct catttccatg 600
 gacatcacct tcttctcaga ggacggagag ctatcatccc tgggaaagga gggcacaggg 660
 ttcagaagta actactttgc ttatgaaact ggaggcaagg cctgcaaaat gcaatactgc 720
 aagcattggg gagtccagact cccatcagggt gtctgggttc agatgggtga taaggatctc 780
 tttgctgcag ccagattccc tgaatgcccga gaagggtcaa gtatctctgc tccatctcag 840
 acctcagtgg atgtaagtct aattcaggac gttgagagga tcttggatta ttccctctgc 900
 caagaaacct ggagcaaaat cagagcgggt ctccaatct ctccagtggg tctcagctat 960
 cttgtccta aaaacccagg aaccggtcct gctttcacca taatcaatgg taacctaaaa 1020
 tactttgaga ccagatacat cagagtcgat attgctgctc caatcctctc aagaatgggtc 1080
 ggaatgatca gtggaactac cacagaaagg gaactgtggg atgactgggc accatatgaa 1140
 gacgtggaaa ttggacccaa tggagtcttg aggaccagtt caggatataa gtttccttta 1200
 tacatgattg gacatgggtat gttggactcc gatcttcac ttagctcaaa ggctcagggtg 1260
 ttccaacatc ctcacattca agacgctgct tcgcaacttc ctgatgatga gagtttattt 1320
 tttggtgata ctgggctatc caaaaatcca atcgagcttg tagaagggtg gttcagtagt 1380
 tggaaaagct ctattgcctc ttttttcttt atcatagggt taatcattgg actattcttg 1440
 gttctccgag ttggtatcca tctttgcatt aaattaaagc acaccaagaa aagacagatt 1500
 tatacagaca tagagatgaa ccgacttggg aagtaa 1536